



# **Cohere Medical Policy – Magnetic Resonance Imaging (MRI), Upper Extremity**

*Clinical Guidelines for Medical Necessity Review*

**Version:** 2  
**Effective Date:** August 5, 2024

# Important Notices

## Notices & Disclaimers:

**GUIDELINES ARE SOLELY FOR COHERE’S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION-MAKING OF END USERS.**

Cohere Health, Inc. (“**Cohere**”) has published these clinical guidelines to determine the medical necessity of services (the “**Guidelines**”) for informational purposes only, and solely for use by Cohere’s authorized “**End Users**”. These Guidelines (and any attachments or linked third-party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision-making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

©2024 Cohere Health, Inc. All Rights Reserved.

## Other Notices:

HCPCS® and CPT® copyright 2024 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

## Guideline Information:

**Specialty Area:** Diagnostic Imaging

**Guideline Name:** Cohere Medical Policy - Magnetic Resonance Imaging (MRI), Upper Extremity

**Date of last literature review:** 7/31/2024

**Document last updated:** 8/5/2024

**Type:**  Adult (18+ yo) |  Pediatric (0-17yo)

## **Table of Contents**

<b>Important Notices</b>	<b>2</b>
Table of Contents	3
<b>Medical Necessity Criteria</b>	<b>4</b>
<b>Service: Magnetic Resonance Imaging (MRI), Upper Extremity</b>	<b>4</b>
Recommended Clinical Approach	4
Medical Necessity Criteria	4
Indications	4
Non-Indications	8
Level of Care Criteria	9
Procedure Codes (CPT/HCPCS)	9
<b>Medical Evidence</b>	<b>10</b>
<b>References</b>	<b>11</b>
<b>Clinical Guideline Revision History/Information</b>	<b>14</b>

# Medical Necessity Criteria

**Service: Magnetic Resonance Imaging (MRI), Upper Extremity**

## Recommended Clinical Approach

Magnetic resonance imaging (MRI) is an advanced imaging modality used when further anatomic detail is required for diagnosis or treatment. It is segmented into joint and non-joint examinations and may be performed without or with contrast (IV or intra-articular). Metal hardware can limit certain exams and is generally inappropriate for imaging by 3 Tesla scanners. Alternate modalities may sometimes be more clinically appropriate based on clinician and supervising radiologist discussion.<sup>1-2</sup>

## Medical Necessity Criteria

### Indications

→ **Magnetic resonance imaging (MRI), upper extremity** is considered appropriate if **ALL** of the following is **TRUE**:

◆ **ANY** of the following:

- Plain radiographs are contraindicated or inconclusive; **OR**
- Ultrasound and CT/CTA are contraindicated or inconclusive (e.g., body habitus for ultrasound, anaphylactic reaction due to IV contrast reaction, pregnancy, pediatric); **AND**

◆ **ANY** of the following is **TRUE**:

- Trauma-related conditions as indicated by **ANY** of the following<sup>3-4</sup>:
  - Fracture (traumatic or stress), and **ANY** of the following is **TRUE**<sup>5</sup>:
    - ◆ Joint dislocation or instability; **OR**
    - ◆ Stress/insufficiency fracture (known) and follow-up imaging needed; **OR**

- ◆ Stress/insufficiency fracture (suspected) with negative radiographs; **OR**
  - Soft tissue injury or tear (e.g., tendon, ligament, muscle)<sup>3-4,6-7</sup>; **OR**
- Detection, screening, and surveillance of neoplasms, masses, and cysts of an upper extremity and **ANY** of the following is **TRUE**<sup>8</sup>:
  - Malignant or aggressive primary tumor<sup>9</sup>; **OR**
  - Evaluation of metastatic lesions of the upper extremity; **OR**
  - Bone tumor is suspected with **ANY** of the following<sup>10</sup>:
    - ◆ Initial radiographs are negative or do not explain symptoms; **OR**
    - ◆ Osteoid osteoma is suspected; **OR**
    - ◆ Lesion present on plain radiographs; **OR**
    - ◆ Indeterminate or aggressive appearance concerning malignancy; **OR**
    - ◆ “Incidental” osseous lesion on other imaging for unrelated indication; **OR**
  - Presence of a mass with **ANY** of the following<sup>2</sup>:
    - ◆ Absence of trauma; **OR**
    - ◆ Rapid growth; **OR**
    - ◆ Recurrence after prior surgery; **OR**
    - ◆ Non-diagnostic ultrasound or other inconclusive imaging; **OR**
  - Known malignancy and **ANY** of the following is required:
    - ◆ Monitor response to treatment; **OR**
    - ◆ Surveillance after treatment or surgery; **OR**
    - ◆ Non-diagnostic ultrasound or other inconclusive imaging; **OR**

- Palpable abnormality with a non-diagnostic radiograph or ultrasound; **OR**
- Infectious disorder including **ANY** of the following:
  - Osteomyelitis, suspected<sup>11</sup>; **OR**
  - Suspected septic arthritis or soft tissue infection is with **ANY** of the following initial radiographs<sup>11-12</sup>:
    - ◆ Normal; **OR**
    - ◆ Suggestive of joint effusion; **OR**
    - ◆ Suggestive of soft tissue swelling; **OR**
  - History of puncture wound with possible retained foreign body; **OR**
  - High clinical suspicion of necrotizing fasciitis; **OR**
- Vascular conditions, known or suspected, including **ANY** of the following:
  - Osteonecrosis or avascular necrosis, known or suspected, (e.g., suspected Kienböck's disease) with negative radiographs<sup>13</sup>; **OR**
  - Vascular compromise, suspected<sup>7</sup>; **OR**
  - Vascular malformation (with or without pain) and **ANY** of the following findings of physical deformity are suspected<sup>14</sup>:
    - ◆ Diffuse or focal enlargement; **OR**
    - ◆ Discoloration; **OR**
    - ◆ Soft-tissue mass; **OR**
    - ◆ Ulceration; **OR**
  - Vascular bruit or thrill; **OR**
- For evaluation of **ANY** of the following uncategorized/miscellaneous symptoms when applicable:
  - Marrow abnormalities<sup>8,13</sup>; **OR**
  - Pain or weakness of an upper extremity, and **ALL** of the following are **TRUE**<sup>7,15</sup>:

- ◆ Failure of conservative management (e.g., rest, analgesics, physical therapy, oral or injectable corticosteroids) must be documented for a period of greater than 3 months; **AND**
- ◆ Concern for acute rupture or high-grade tear of tendon, ligament, or soft tissue injury based on **ALL** of the following:
  - Clinical history; **AND**
  - Physical exam; **AND**
  - Prior radiographs excluding chronic pain or suspicion of a chronic or low-grade injury; **OR**
- Neurological symptoms or deficits with **ANY** of the following<sup>16</sup>:
  - ◆ Peripheral nerve sheath tumor suspected with **ANY** of the following:
    - Enlarging mass; **OR**
    - New or worsening localized pain; **OR**
    - Recurrence after prior resection; **OR**
  - ◆ Localized EMG abnormality; **OR**
  - ◆ Persistent symptoms or suspected nerve entrapment as confirmed by abnormal EMG; **OR**
  - ◆ Trauma or injury with suspected nerve injury or laceration; **OR**
- Initial diagnosis or follow-up of autoimmune, collagen vascular diseases, or inflammatory conditions (e.g., inflammatory arthritis)<sup>12</sup>; **OR**
- Synovial-related disorders (e.g., synovitis, bursitis, metaplasia, and neoplasia)<sup>16</sup>; **OR**
- Repeat imaging of a specific area or structure using the same imaging modality (in the absence of an existing

follow-up guideline) is considered appropriate when **ALL** of the following is **TRUE**:

- There is documented clinical necessity; **AND**
- Prior imaging results of the specific area or structure, obtained using the same imaging modality, must be documented and available for comparison; **AND**
- **ANY** of the following is **TRUE**:
  - ◆ A change in clinical status, such as worsening symptoms or the emergence of new symptoms, that may influence the treatment approach; **OR**
  - ◆ The requirement for interval reassessment, which may alter the treatment plan; **OR**
  - ◆ One-time follow-up of a prior indeterminate finding to assess for interval change; **OR**
  - ◆ The need for re-imaging either before or after performing an invasive procedure.

### **Non-Indications**

→ **Magnetic resonance imaging (MRI), upper extremity** may not be considered appropriate if **ANY** of the following is **TRUE**:

- ◆ The patient has undergone advanced imaging of the same body part within 3 months without undergoing treatment or developing new or worsening symptoms; **OR**
- ◆ If contrast is used, history of anaphylactic allergic reaction to gadolinium contrast media with detailed guidelines for use in patients with renal insufficiency; **OR**
- ◆ The patient has metallic clips on vascular aneurysms; **OR**
- ◆ Metallic foreign body in orbits/other critical area(s) or within the field of view and obscuring area of concern; **OR**
- ◆ Incompatible implanted devices (e.g., pacemakers, defibrillators, cardiac valves, spinal cord stimulators).



\*NOTE: MRI in patients with claustrophobia should be requested at the discretion of the ordering provider.

\*\*NOTE: MRI in pregnant patients should be requested at the discretion of the ordering provider and obstetric care provider.

### **Level of Care Criteria**

Inpatient or Outpatient

### **Procedure Codes (CPT/HCPCS)**

<b>CPT/HCPCS Code</b>	<b>Code Description</b>
73218	Magnetic resonance imaging (MRI) (e.g., proton), upper extremity, other than joint; without contrast material(s)
73219	Magnetic resonance imaging (MRI) (e.g., proton), upper extremity, other than joint; with contrast material(s)
73220	Magnetic resonance imaging (MRI) (e.g., proton), upper extremity, other than joint; without contrast material(s), followed by contrast material(s) and further sequences
73221	Magnetic resonance imaging (MRI) (e.g., proton), any joint of upper extremity; without contrast material(s)
73222	Magnetic resonance imaging (MRI) (e.g., proton), any joint of upper extremity; with contrast material(s)
73223	Magnetic resonance imaging (MRI) (e.g., proton), any joint of upper extremity; without contrast material(s), followed by contrast material(s) and further sequences.

# Medical Evidence

DeFrance et al. (2021) performed a study using magnetic resonance imaging (MRI) for diagnosing upper extremity conditions and assessed how MRI findings influenced patient management. Findings from 187 patients who had an MRI were analyzed on the usefulness of imaging and how they influenced treatment decisions. Imaging was ordered to assess for suspected occult scaphoid fractures, ulnar wrist pain, collateral ligament injuries of the metacarpophalangeal joint, and masses. Surgeons concurred with radiologists' interpretations in 88% of cases. Overall, surgeons noted MRI findings as helpful in 92% of cases.<sup>17</sup>

Cortes et al. (2019) conducted a small prospective study on patients with suspected cuff tendinopathy. Fifty-one patients were included and underwent magnetic resonance imaging (MRI). Ninety percent (n=46) did not require surgical intervention; individuals who underwent surgery (10%) within an average of 68.3 days post-imaging. A significant proportion (over 90%) underwent premature MRI, which illustrates early MRI utilization in patients with atraumatic shoulder pain whose condition may have improved with conservative treatment first.<sup>18</sup>

Rubin (2019) analyzed MRI and ultrasound findings of patients with rheumatoid arthritis in the hands and wrists. Advanced imaging modalities allow for visualization of synovitis and active soft-tissue inflammation, which are early indicators of potential structural damage. MRI can also identify osteitis, a crucial prognostic marker for disease aggressiveness. Findings include the distinct definitions (specifically synovitis, osteitis, and erosions) that enhance clinical assessment and imaging interpretation.<sup>19</sup>

# References

1. Expert Panel on Musculoskeletal Imaging, Roberts CC, Metter DF, et al. ACR appropriateness criteria - imaging after shoulder arthroplasty: 2021 update. *J Am Coll Radiol*. 2022 May;19(5S):S53-S66. doi: 10.1016/j.jacr.2022.02.019. PMID: 35550805.
2. Expert Panel on Musculoskeletal Imaging, Garner HW, Wessell DE, et al. ACR appropriateness criteria - soft tissue masses: 2022 update. *J Am Coll Radiol*. 2023 May;20(5S):S234-S245. doi: 10.1016/j.jacr.2023.02.009. PMID: 37236746.
3. Expert Panel on Musculoskeletal Imaging, Torabi M, Lenchik L, et al. ACR appropriateness criteria - acute hand and wrist trauma. *J Am Coll Radiol*. 2019 May;16(5S):S7-S17. doi: 10.1016/j.jacr.2019.02.029. PMID: 31054760.
4. Expert Panel on Musculoskeletal Imaging, Amini B, Beckmann NM, et al. ACR appropriateness criteria - shoulder pain-traumatic. *J Am Coll Radiol*. 2018 May;15(5S):S171-S188. doi: 10.1016/j.jacr.2018.03.013. PMID: 29724420.
5. Expert Panel on Musculoskeletal Imaging, Bencardino JT, Stone TJ, et al. ACR appropriateness criteria - stress (fatigue/insufficiency) fracture, including sacrum, excluding other vertebrae. *J Am Coll Radiol*. 2017 May;14(5S):S293-S306. doi: 10.1016/j.jacr.2017.02.035. PMID: 28473086.
6. Expert Panel on Musculoskeletal Imaging, Nacey N, Fox MG, et al. ACR appropriateness criteria - chronic shoulder pain: 2022 update. *J Am Coll Radiol*. 2023 May;20(5S):S49-S69. doi: 10.1016/j.jacr.2023.02.017. PMID: 37236752.
7. Expert Panel on Musculoskeletal Imaging, Stensby JD, Fox MG, et al. ACR appropriateness criteria - chronic hand and wrist pain. Updated 2023. Accessed July 31, 2024. <https://acsearch.acr.org/docs/69427/Narrative/>.
8. American College of Radiology (ACR). ACR practice parameter for performing and interpreting magnetic resonance imaging (MRI) - resolution 8. Updated 2022. Accessed July 31, 2024. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-Perf-Interpret.pdf>.

9. Expert Panel on Musculoskeletal Imaging, Stanborough R, Demertzis JL, et al. ACR appropriateness criteria – malignant or aggressive primary musculoskeletal tumor–staging and surveillance: 2022 update. *J Am Coll Radiol*. 2022 Nov;19(11S):S374–S389. doi: 10.1016/j.jacr.2022.09.015. PMID: 36436964.
10. Expert Panel on Musculoskeletal Imaging, Bestic JM, Wessell DE, et al. ACR appropriateness criteria – primary bone tumors. *J Am Coll Radiol*. 2020 May;17(5S):S226–S238. doi: 10.1016/j.jacr.2020.01.038. PMID: 32370967.
11. Expert Panel on Musculoskeletal Imaging, Pierce JL, Perry MT, et al. ACR appropriateness criteria – suspected osteomyelitis, septic arthritis, or soft tissue infection (excluding spine and diabetic foot): 2022 update. *J Am Coll Radiol*. 2022 Nov;19(11S):S473–S487. doi: 10.1016/j.jacr.2022.09.013. PMID: 36436971.
12. Expert Panel on Musculoskeletal Imaging, Subhas N, Wu F, et al. ACR appropriateness criteria – chronic extremity joint pain–suspected inflammatory arthritis, crystalline arthritis, or erosive osteoarthritis: 2022 update. *J Am Coll Radiol*. 2023 May;20(5S):S20–S32. doi: 10.1016/j.jacr.2023.02.020. PMID: 37236743.
13. Expert Panel on Musculoskeletal Imaging, Ha AS, Chang EY, et al. ACR appropriateness criteria – osteonecrosis: 2022 update. *J Am Coll Radiol*. 2022 Nov;19(11S): S409–S416. doi: 10.1016/j.jacr.2022.09.009. PMID: 36436966.
14. Expert Panel on Vascular Imaging, Obara P, McCool J, et al. ACR appropriateness criteria – clinically suspected vascular malformation of the extremities. *J Am Coll Radiol*. 2019 Nov;16(11S):S340–S347. doi: 10.1016/j.jacr.2019.05.013. PMID: 31685102.
15. Expert Panel on Musculoskeletal Imaging, Thomas JM, Chang EY, et al. ACR appropriateness criteria – chronic elbow pain. *J Am Coll Radiol*. 2022 Nov;19(11S):S256–S265. doi: 10.1016/j.jacr.2022.09.022. PMID: 36436956.
16. American College of Radiology (ACR), Society of Pediatric Radiology (SPR), Society of Skeletal Radiology (SSR). ACR–SPR–SSR practice parameter for the performance and interpretation of magnetic resonance imaging (MRI) of the shoulder – resolution 32. Published

2020. Accessed July 31, 2024.

<https://www.acr.org/-/media/ACR/Files/Practice-Parameters/mr-shldr.pdf>.

17. DeFrance MJ, Lutsky KF, Ilyas AM, et al. MRI utility in hand surgery. *Hand (N Y)*. 2021 May;16(3):298–302. doi: 10.1177/1558944719861725. PMID: 31288574; PMCID: PMC8120576.
18. Cortes A, Quinlan NJ, Nazal MR, Upadhyaya S, Alpaugh K, Martin SD. A value-based care analysis of magnetic resonance imaging in patients with suspected rotator cuff tendinopathy and the implicated role of conservative management. *J Shoulder Elbow Surg*. 2019 Nov;28(11):2153–2160. doi: 10.1016/j.jse.2019.04.003. Epub 2019 Jul 4. PMID: 31281001.
19. Rubin DA. MRI and ultrasound of the hands and wrists in rheumatoid arthritis: Imaging findings. *Skeletal Radiol*. 2019 May;48(5):677–695. doi: 10.1007/s00256-019-03179-z. PMID: 30796506.

# Clinical Guideline Revision History/Information

Original Date: April 4, 2022		
Review History		
Version 2	8/5/2024	Annual review and policy restructure.