



Cohere Medical Policy – Computed Tomography Angiography (CTA), Lower Extremity

Clinical Guidelines for Medical Necessity Review

Version: 3
Effective Date: October 30, 2024

Important Notices

Notices & Disclaimers:

GUIDELINES ARE SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION-MAKING OF END USERS.

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine the medical necessity of services (the "**Guidelines**") for informational purposes only, and solely for use by Cohere's authorized "**End Users**". These Guidelines (and any attachments or linked third-party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision-making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

©2024 Cohere Health, Inc. All Rights Reserved.

Other Notices:

HCPCS® and CPT® copyright 2024 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

Guideline Information:

Specialty Area: Diagnostic Imaging

Guideline Name: Cohere Medical Policy – Computed Tomography Angiography (CTA), Lower Extremity

Date of last literature review: 8/12/2024

Document last updated: 10/30/2024

Type: ☒ Adult (18+ yo) | ☒ Pediatric (0-17 yo)

Table of Contents

Important Notices	2
Table of Contents	3
Medical Necessity Criteria	4
Service: Computed Tomography Angiography (CTA), Lower Extremity	4
Recommended Clinical Approach	4
Medical Necessity Criteria	4
Indications	4
Non-Indications	8
Disclaimer on Radiation Exposure in Pediatric Population	9
Level of Care Criteria	9
Procedure Codes (CPT/HCPCS)	10
Medical Evidence	11
References	12
Clinical Guideline Revision History/Information	14

Medical Necessity Criteria

Service: Computed Tomography Angiography (CTA), Lower Extremity

Recommended Clinical Approach

Lower extremity computed tomography angiography (CTA) is indicated for peripheral artery disease (PAD), trauma, assessment of vascular anatomy and congenital malformations, vasculitis, and surgical planning.¹ After the intravenous injection of an iodinated contrast medium, CTA employs a thin-section CT scan timed to capture peak arterial and/or venous enhancement, depending on the targeted vascular structures. The resulting volumetric data set is analyzed using primary transverse reconstructions along with multiplanar reformations and 3-D renderings.¹

Medical Necessity Criteria

Indications

- **Computed tomography angiography (CTA), lower extremity** is considered appropriate if **ANY** of the following is **TRUE**:
- ◆ Neoplastic conditions (including masses or mass-like conditions) when the arterial blood supply needs to be evaluated (e.g., for treatment planning, treatment response, or prognostication); **OR**
 - ◆ Neoplastic invasion of arteries or veins; **OR**
 - ◆ Trauma-related conditions as indicated by **ANY** of the following²:
 - Expanding hematoma³; **OR**
 - Major blunt trauma and the patient is hemodynamically stable⁴; **OR**
 - Neurologic deficit of lower extremity in association with trauma⁵; **OR**
 - Known or suspected knee dislocation^{*6}; **OR**
 - Vascular trauma to a lower extremity⁷; **OR**
 - ◆ Vascular conditions, known or suspected, including **ANY** of the following:
 - Aneurysm, seen on ultrasound or where ultrasound is nondiagnostic; **OR**
 - Intramural hematoma; **OR**

- Dissection; **OR**
- Critical limb ischemia strongly suspected (e.g. sudden onset of a cold leg with pain, gangrene, rest pain)⁸; **OR**
- Lower extremity ischemic symptoms when **ALL** of the following are **TRUE**:
 - **ANY** of the following:
 - ◆ Leg pain worsens with activity and is relieved with rest (claudication); **OR**
 - ◆ Non-healing lower extremity ulcers; **AND**
 - **ALL** of the following:
 - ◆ Limitation of performance of daily activities; **AND**
 - ◆ Expected mobility after treatment warrants revascularization; **AND**
 - ◆ Revascularization is planned⁹; **AND**
 - ◆ Abnormal ABI as evidenced by **ANY** of the following:
 - ABI is inconclusive or nondiagnostic; **OR**
 - ABI less than 0.9 or greater than 1.4 on at least one leg; **OR**
 - ABI less than 1.1 in patients with risk factors for atherosclerosis (e.g., personal history of diabetes or known cardiac disease)¹⁰; **AND**
 - ◆ Either low concern for aortic and iliac artery disease or aorta and iliac arteries previously imaged; **OR**
- Determination of hemorrhage source (including non-surgical, spontaneous)³; **OR**
- Localization and characterization of vascular malformation or fistula (e.g., assessing treatment response, treatment planning) with **ANY** of the following:
 - Duplex ultrasound indeterminate or nondiagnostic; **OR**
 - High flow lesion suspected clinically or by imaging; **OR**
 - Preoperative planning; **OR**

- Vasculitis, initial evaluation, when **ANY** of the following is **TRUE**⁷:
 - Biopsy proven; **OR**
 - Rheumatologic panel work-up including but not limited to erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) is suggestive of vasculitis; **OR**
 - The requesting clinician specializes in rheumatology and the outcome of the imaging is expected to change management and/or treatment plan; **OR**
- Popliteal entrapment syndrome, when ultrasound and MRI/MRA are indeterminate or contraindicated⁷; **OR**
- ◆ Pre and post-intervention evaluation when **ANY** of the following is **TRUE**:
 - Postoperative evaluation of the effectiveness of arterial or venous reconstruction or bypass; **OR**
 - Characterization of normal and variant vascular anatomy; **OR**
 - Determination of the patency, location, or integrity of grafts and other vascular devices (e.g., stents); **OR**
 - Planning autografts for musculoskeletal reconstruction; **OR**
 - Treatment of popliteal entrapment syndrome; **OR**
- ◆ Hemodialysis access evaluation, when duplex ultrasound is inconclusive and fistulogram cannot be performed, for **ANY** of the following conditions:
 - Arteriovenous fistula (AVF) stenosis; **OR**
 - Occlusion; **OR**
 - Pseudoaneurysm; **OR**
 - Steal syndrome (cool and painful extremity); **OR**
- ◆ Repeat imaging (defined as repeat request following recent imaging of the same anatomic region with the same modality), in the absence of established guidelines, will be considered reasonable and necessary if **ANY** of the following is **TRUE**:
 - New or worsening symptoms, such that repeat imaging would influence treatment; **OR**
 - One-time clarifying follow-up of a prior indeterminate finding; **OR**

- In the absence of change in symptoms, there is an established need for monitoring which would influence management.

→ **Computed tomography venography (CTV), lower extremity** is

considered appropriate for **ANY** of the following is **TRUE**:

- ◆ Neoplastic conditions (including masses or mass-like conditions) when the arterial blood supply needs to be evaluated (e.g., for treatment planning, treatment-response, or prognostication); **OR**
- ◆ Neoplastic invasion of arteries or veins; **OR**
- ◆ Evaluation for known venous leg ulcer after ultrasound has been completed; **OR**
- ◆ Known or suspected acute or chronic deep venous thrombosis, when results would change management and ultrasound has been completed¹¹; **OR**
- ◆ Known severe post-thrombotic changes incompletely evaluated by ultrasound¹¹; **OR**
- ◆ Evidence of severe venous reflux disease and **ALL** of the following¹¹:
 - Duplex ultrasound evaluation indeterminate, incomplete, or non-diagnostic; **AND**
 - Surgical or endovascular intervention planned; **OR**
- ◆ Pre and post-intervention evaluation when **ANY** of the following is **TRUE**:
 - Postoperative evaluation of the effectiveness of arterial or venous reconstruction or bypass; **OR**
 - Characterization of normal and variant vascular anatomy; **OR**
 - Determination of the patency, location, or integrity of grafts and other vascular devices (e.g. stents); **OR**
 - Planning autografts for musculoskeletal reconstruction; **OR**
 - Treatment of popliteal entrapment syndrome; **OR**
- ◆ Repeat imaging (defined as repeat request following recent imaging of the same anatomic region with the same modality), in the absence of established guidelines, will be considered reasonable and necessary if **ANY** of the following is **TRUE**:
 - New or worsening symptoms, such that repeat imaging would influence treatment; **OR**

- One-time clarifying follow-up of a prior indeterminate finding; **OR**
- In the absence of change in symptoms, there is an established need for monitoring which would influence management.

Non-Indications

→ **Computed tomography angiography (CTA)/computed tomography venography (CTA/CTV), lower extremity with contrast** is not considered appropriate if **ANY** of the following is **TRUE**¹²:

- ◆ The patient has undergone advanced imaging of the same body part within 3 months without undergoing treatment or developing new or worsening symptoms¹³; **OR**
- ◆ The request is for CT/CTA with contrast, and the patient has a history of anaphylactic allergic reaction to iodinated contrast media; **OR**
- ◆ Evaluation of lower extremity arterial perfusion, such as for claudication, when there may be a concern for aorta or iliac disease and aorta and iliac have not been imaged.

*NOTE: The referring professional and radiologist should discuss the risks and benefits of contrast media administration, including possible prophylaxis, in patients with chronic or worsening kidney disease or severe renal failure.

**NOTE: CT in pregnant patients should be requested at the discretion of the ordering provider and obstetric care provider.

***NOTE: CT in patients with claustrophobia should be requested at the discretion of the ordering provider.

Disclaimer on Radiation Exposure in Pediatric Population

Due to the heightened sensitivity of pediatric patients to ionizing radiation, minimizing exposure is paramount. At Cohere, we are dedicated to ensuring that every patient, including the pediatric population, has access to appropriate imaging following accepted guidelines. Radiation risk is dependent mainly on the patient's age at exposure, the organs exposed, and the patient's sex, though there are other variables. The following technical guidelines are provided to ensure safe and effective imaging practices:

Radiation Dose Optimization: Adhere to the lowest effective dose principle for pediatric imaging. Ensure that imaging protocols are specifically tailored for pediatric patients to limit radiation exposure.¹⁴⁻¹⁵

Alternative Modalities: Prioritize non-ionizing imaging options such as ultrasound or MRI when clinically feasible, as they are less likely to expose the patient to ionizing radiation. For instance, MRI or ultrasound should be considered if they are more likely to provide an accurate diagnosis than CT, fluoroscopy, or radiography.¹⁴⁻¹⁵

Cumulative Dose Monitoring: Implement systems to track cumulative radiation exposure in pediatric patients, particularly for those requiring multiple imaging studies. Regularly reassess the necessity of repeat imaging based on clinical evaluation.¹⁴⁻¹⁵

CT Imaging Considerations: When CT is deemed the best method for achieving a correct diagnosis, use the lowest possible radiation dose that still yields reliable diagnostic images.¹⁴⁻¹⁵

Cohere Imaging Gently Guideline

The purpose of this guideline is to act as a potential override when clinically indicated to adhere to Imaging Gently and Imaging Wisely guidelines and As Low As Reasonably Possible (ALARA) principles.

Level of Care Criteria

Inpatient and Outpatient

Procedure Codes (CPT/HCPCS)

CPT/HCPCS Code	Code Description
73706	Computed tomographic angiography (CTA), lower extremity; with contrast material(s), including non-contrast images, if performed, and image postprocessing

Medical Evidence

Jayaraj et al. (2024) evaluate using CT venography (CTV) to diagnose chronic iliac venous obstruction. Advantages of CTV include acquisition and postprocessing functionalities, including multiplanar reconstruction assessment and image processing techniques (e.g., volume rendering and maximum intensity projection). These serve to ascertain and categorize the features of the lesion. CTV also enables the detection of extravascular structures potentially implicated in compression and obstruction. The identification of congenital anatomical variances was present in 20% of patients with these conditions.¹⁶

Kakkos et al. (2021) discuss the efficacy of CTV as a reliable diagnostic tool for proximal DVT in patients suspected of DVT and pulmonary embolism (PE), exhibiting sensitivity and specificity akin to ultrasound. CTV presents clear advantages over ultrasound in assessing pelvic veins and the inferior vena cava (IVC) while also identifying concurrent medical conditions manifesting as pain and swelling. The superior spatial resolution of CTV facilitates precise vessel measurements and treatment planning when intervention is warranted. Limitations of CTV include the necessity of iodine contrast administration and exposure to radiation – a concern, especially in younger patients.¹⁷

Cook et al. (2016) review CTA of the lower extremities for conditions including peripheral artery disease (PAD), trauma, evaluation of variant anatomy and congenital malformations, vasculitis, and pre-surgical planning. CTA is a crucial and adaptable noninvasive technique with various diagnostic and procedural applications. While it is commonly employed for patients with PAD or lower extremity trauma, the utility of CTA extends to investigating non-ischemic conditions and congenital vascular abnormalities. Tailoring CT scanner protocols to specific clinical inquiries, including adjusting bolus timing and employing multiphasic imaging, is essential. Additionally, the utilization of three-dimensional postprocessing techniques is vital for enhancing visualization and interpretation of the detailed imaging data obtained from these examinations.¹⁸

References

1. American College of Radiology (ACR), North American Society for Cardiovascular Imaging (NASCI), Society of Interventional Radiology (SIR), Society for Pediatric Radiology (SPR). ACR–NASCI–SIR–SPR practice parameter for the performance and interpretation of body computed tomography (CTA) – resolution 47. Updated 2021. Accessed January 16, 2024. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/body-cta.pdf>.
2. Brian R, Bennett DJ, Kim WC, et al. Computed tomography angiography is associated with low added utility for detecting clinically relevant vascular injuries among patients with extremity trauma. *Trauma Surg Acute Care Open*. 2021 Dec 20;6(1):e000828. doi: 10.1136/tsaco-2021-000828. PMID: 34993352; PMCID: PMC8689162.
3. Fox N, Rajani RR, Bokhari F, et al. Evaluation and management of penetrating lower extremity arterial trauma: An Eastern Association for the Surgery of Trauma practice management guideline. *J Trauma Acute Care Surg*. 2012 Nov;73(5 Suppl 4):S315–20. doi: 10.1097/TA.0b013e31827018e4. PMID: 23114487.
4. Expert Panel on Major Trauma Imaging, Shyu JY, Khurana B, et al. ACR appropriateness criteria – major blunt trauma. *J Am Coll Radiol*. 2020 May;17(5S):S160–S174. doi: 10.1016/j.jacr.2020.01.024. PMID: 32370960.
5. Dreizin D, Smith EB, Champ K, et al. Roles of trauma CT and CTA in salvaging the threatened or mangled extremity. *Radiographics*. 2022 Mar–Apr;42(2):E50–E67. doi: 10.1148/rg.210092. PMID: 35230918; PMCID: PMC8906352.
6. Expert Panel on Musculoskeletal Imaging, Taljanovic MS, Chang EY, et al. ACR appropriateness criteria – acute trauma to the knee. *J Am Coll Radiol*. 2020 May;17(5S):S12–S25. doi: 10.1016/j.jacr.2020.01.041. PMID: 32370956.
7. Expert Panels on Vascular Imaging, Francois CJ, Skulborstad EP, et al. ACR appropriateness criteria – nonatherosclerotic peripheral arterial disease. *J Am Coll Radiol*. 2019 May;16(5S):S174–S183. doi: 10.1016/j.jacr.2019.02.026. PMID: 31054743.
8. Expert Panel on Vascular Imaging, Browne WF, Sung J, et al. ACR appropriateness criteria – sudden onset of cold, painful leg: 2023 update. *J Am Coll Radiol*. 2023 Nov;20(11S):S565–S573. doi: 10.1016/j.jacr.2023.08.012. PMID: 38040470.

9. Expert Panel on Vascular Imaging, Azene EM, Steigner ML, Aghayev A, et al. ACR appropriateness criteria – lower extremity arterial claudication–imaging assessment for revascularization: 2022 update. *J Am Coll Radiol*. 2022 Nov;19(11S):S364–S373. doi: 10.1016/j.jacr.2022.09.002. PMID: 36436963; PMCID: PMC9876734.
10. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC guideline on the management of patients with lower extremity peripheral artery disease: A report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2017 Mar 21;135(12):e726–e779. doi: 10.1161/CIR.0000000000000471. PMID: 27840333; PMCID: PMC5477786.
11. Expert Panels on Interventional Radiology and Vascular Imaging, Rochon PJ, Reghunathan A, et al. ACR appropriateness criteria – lower extremity chronic venous disease. *J Am Coll Radiol*. 2023 Nov;20(11S):S481–S500. doi: 10.1016/j.jacr.2023.08.011. PMID: 38040466.
12. Davenport MS, Perazella MA, Yee J, et al. Use of intravenous iodinated contrast media in patients with kidney disease: Consensus statements from the American College of Radiology and the National Kidney Foundation. *Radiology*. 2020;294(3):660–668. doi: 10.1148/radiol.2019192094. PMID: 33015613; PMCID: PMC7525144.
13. Wasser EJ, Prevedello LM, Sodickson A, Mar W, Khorasani R. Impact of a real-time computerized duplicate alert system on the utilization of computed tomography. *JAMA Intern Med*. 2013;173(11):1024–1026. doi: 10.1001/jamainternmed.2013.543. PMID: 23609029.
14. The Image Gently Alliance. Procedures – image gently and CT scans. Updated 2014. Accessed June 26, 2024. <https://www.imagegently.org/Procedures/Computed-Tomography>.
15. National Cancer Institute. Radiation risks and pediatric computed tomography (CT): A guide for health care. Updated September 4, 2018. Accessed June 26, 2024. <https://www.cancer.gov/about-cancer/causes-prevention/risk/radiation/pediatric-ct-scans>.
16. Jayaraj A, Rossi FH, Lurie F, et al. Diagnosis of chronic iliac venous obstruction. *J Vasc Surg Venous Lymphat Disord*. 2024 Jan 18:101744. doi: 10.1016/j.jvsv.2023.101744. PMID: 38242206.
17. Kakkos SK, Gohel M, Baekgaard N, et al. Editor's choice – European Society for Vascular Surgery (ESVS) 2021 clinical practice guidelines on the management of venous thrombosis. *Eur J Vasc Endovasc Surg*. 2021 Jan;61(1):9–82. doi: 10.1016/j.ejvs.2020.09.023. PMID: 33334670.
18. Cook TS. Computed tomography angiography of the lower extremities. *Radiol Clin North Am*. 2016 Jan;54(1):115–30. doi: 10.1016/j.rcl.2015.08.001. PMID: 26654395.

Clinical Guideline Revision History/Information

Original Date: April 22, 2022		
Review History		
Version 2	8/13/2024	Annual review and policy restructure.
Version 3	10/30/2024	Edited repeat imaging criteria language.