



Cohere Medical Policy – Hammertoe, Claw Toe, or Mallet Toe Surgical Treatment (With or Without Fusion)

Clinical Guidelines for Medical Necessity Review

Version: 2
Effective Date: September 20, 2024

Important Notices

Notices & Disclaimers:

GUIDELINES SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION MAKING OF END USERS.

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine medical necessity of services (the "**Guidelines**") for informational purposes only, and solely for use by Cohere's authorized "**End Users**". These Guidelines (and any attachments or linked third party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

©2024 Cohere Health, Inc. All Rights Reserved.

Other Notices:

HCPCS® and CPT® copyright 2024 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

Guideline Information:

Specialty Area: Diseases & Disorders of the Musculoskeletal System

Guideline Name: Cohere Medical Policy - Hammertoe, Claw Toe, or Mallet Toe Surgical Treatment with or without Fusion

Literature review current through: 7/18/2024

Document last updated: 9/20/2024

Type: ☒ Adult (18+ yo) | ☒ Pediatric (0-17yo)

Table of Contents

Important Notices	2
Medical Necessity Criteria	4
Service: Hammertoe, Claw Toe, or Mallet Toe Surgical Treatment (With or Without Fusion)	4
General Guidelines	4
Medical Necessity Criteria	4
Indications	4
Non-Indications	5
Level of Care Criteria	5
Procedure Codes (CPT/HCPCS)	5
Medical Evidence	7
References	8
Clinical Guideline Revision History/Information	9

Medical Necessity Criteria

Service: Hammertoe, Claw Toe, or Mallet Toe Surgical Treatment (With or Without Fusion)

General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach:** Surgery may be appropriate for a hammertoe, claw toe, or mallet toe deformity that is not alleviated by conservative management. Surgical treatment may consist of arthrodesis, arthroplasty, amputation, or tendon release/transfer. Surgical repair for cosmetic purposes only is not recommended.¹⁻³
- **Exclusions:** None.

Medical Necessity Criteria

Indications

→ **Hammertoe, claw toe, or mallet toe surgical treatment (with or without fusion)** is considered appropriate if **ALL** of the following are **TRUE**:

- ◆ Radiographic confirmation of hammertoe or mallet toe deformity³; **AND**
- ◆ The patient has **ANY** of the following clinical presentations or positive findings:
 - Pain; **OR**
 - Difficulty walking; **OR**
 - The toe bends downwards or appears clawlike; **OR**
 - Balance may be affected; **OR**
 - Inability to flex or wiggle toes; **OR**
 - Callosities on toe; **OR**
 - Crossing over of lesser toes; **AND**
- ◆ Failure of conservative management for greater than 3 months, including **ALL** of the following¹⁻⁴:
 - Trial of **ANY** of the following:
 - Taping; **OR**
 - Splinting; **OR**

- Toe spacer; **OR**
- Toe sleeve; **OR**
- Padding; **AND**
- Shoe modifications; **AND**
- Oral steroids, topical or oral anti-inflammatory medications, or oral analgesics; **AND**
- **ANY** of the following:
 - Corticosteroid injection if medically appropriate; **OR**
 - Corticosteroid injection is contraindicated; **AND**
- ◆ The patient has **ANY** of the following⁵:
 - Bursitis; **OR**
 - Ankylosis of the proximal interphalangeal (PIP) or distal interphalangeal (DIP) joint; **OR**
 - Interdigital neuroma from the deformity; **OR**
 - Lateral metatarsophalangeal (MTP) capsular tear caused by the deformity; **OR**
 - Subluxation or dislocation of the MTP joint from the deformity; **OR**
 - MTP plantar plate tear; **OR**
 - Synovitis/capsulitis of the MTP joint; **OR**
 - Ulceration at the apex of the deformity.

Non-Indications

→ **Hammertoe, claw toe, or mallet toe surgical treatment (with or without fusion)** is not considered appropriate if **ANY** of the following is **TRUE**:

- ◆ There is poor circulation; **OR**
- ◆ The patient has an active, uncontrolled infection (does not apply to chronic infections that are managed).

Level of Care Criteria

Outpatient

Procedure Codes (CPT/HCPCS)

CPT/HCPCS Code	Code Description
28010	Tenotomy of toe tendon, accessed through the skin
28232	Incision to lengthen toe tendon, open procedure

28285	Correction, hammertoe (eg, interphalangeal fusion, partial or total phalangectomy)
28286	Correction, cock-up fifth toe, with plastic closure
28308	Osteotomy, with or without lengthening, shortening, or angular correction, metatarsal; other than first metatarsal
28312	Osteotomy, shortening, angular or rotational correction; other phalanges, any toe
28313	Reconstruction of soft tissue angular deformity of toe
28899	Unlisted procedure of foot and toes.

Medical Evidence

Thomas et al. (2009) developed a Clinical Practice Guideline for the American College of Foot and Ankle Surgeons for the diagnosis and treatment of forefoot disorders: digital deformities. The guideline consists of multiple pathways which include digital deformities, central metatarsalgia, Morton's neuroma, tailor's bunion, and trauma. They state that trauma may be an etiology of digital deformity; however, congenital or acquired deformities are more common. Examinations are generally performed sitting or standing, and gait analysis is stated to be beneficial. Regarding clinical maneuvers, the push-up test is effective at determining whether the deformity can be reduced. The metaphalangeal joint drawer test assists in confirming sagittal and transverse instability and the potential for plantar plate pathology.¹

Shirzad and colleagues performed a systematic review of the literature in 2011 regarding lesser toe deformities. The group stated that in mallet toe, good results had been reported with distal interphalangeal (DIP) joint resection arthroplasty or fusion with a 72% fusion rate in one study of 50 patients with 97% of patients reporting satisfactory pain relief. Flexor digitorum longus (FDL) tendon transfer study patients reported 90% satisfaction in metatarsophalangeal (MTP) joint instability or subluxation correction. Other studies in this area reported complete correction of subluxation in only 54% of patients, and in the largest published series of FDL tendon transfers, 34 of 38 patients were satisfied with the procedure. In proximal interphalangeal resection arthroplasty, fusion rates of 81 to 100% were reported, with additional studies reporting high success rates.²

Mizel and Yodlowski (1995) discuss disorders of the lesser metatarsophalangeal joints including, claw toe, hammer toe, plantar keratoses, Freiberg's infraction, and cock-up fifth toe. Claw toe often results from hyperextension at the metatarsophalangeal joint with flexion deformities at the interphalangeal joints. Along with hammer toe, these conditions are usually acquired and progressive, often involving multiple toes. Freiberg's infraction patients present with pain at the MTP joint that is usually exacerbated by activity. Magnetic resonance imaging results frequently appear as suspected avascular necrosis.³

References

1. Thomas JL, Blitch EL, Chaney DM, et al. Diagnosis and treatment of forefoot disorders – section 1: Digital deformities. *J Foot Ankle Surg.* 2009 Mar-Apr;48(2):230–8. doi: 10.1053/j.jfas.2008.12.003. PMID: 19232978.
2. Shirzad K, Kiesau CD, DeOrio JK, et al.. Lesser toe deformities. *J Am Acad Orthop Surg.* 2011 Aug;19(8):505–14. doi: 10.5435/00124635-201108000-00006. PMID: 21807918.
3. American College of Foot and Ankle Surgeons (ACFAS). ACFAS position statement on cosmetic surgery. <https://www.acfas.org>. Published February 2004. Updated February 2020.
4. Murphy GA. Lesser toe abnormalities. In: Azar, F, Beaty JH. Campbell's Operative Orthopaedics. 14th ed. Elsevier; 2021:4227–4283.e3.
5. Mizel MS, Yodlowski, ML. Disorders of the lesser metatarsophalangeal joints. *J Am Acad Orthop Surg.* 1995 May;3(3):166–173. doi: 10.5435/00124635-199505000-00006. PMID: 10790665.

Clinical Guideline Revision History/Information

Original Date: April 12, 2024		
Review History		
Version 2	9/20/2024	Updated language regarding conservative treatment.