



Cohere Medical Policy – Magnetic Resonance Angiography (MRA), Abdomen/Pelvis

Clinical Guidelines for Medical Necessity Review

Version: 3
Effective Date: October 30, 2024

Important Notices

Notices & Disclaimers:

GUIDELINES ARE SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION-MAKING OF END USERS.

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine the medical necessity of services (the "**Guidelines**") for informational purposes only, and solely for use by Cohere's authorized "**End Users**". These Guidelines (and any attachments or linked third-party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision-making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

©2024 Cohere Health, Inc. All Rights Reserved.

Other Notices:

HCPCS® and CPT® copyright 2024 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

Guideline Information:

Specialty Area: Diagnostic Imaging

Guideline Name: Cohere Medical Policy - Magnetic Resonance Angiography (MRA), Abdomen/Pelvis

Date of last literature review: 6/18/2024

Document last updated: 10/30/2024

Type: Adult (18+ yo) | Pediatric (0-17yo)

Table of Contents

Important Notices	2
Table of Contents	3
Medical Necessity Criteria	4
Service: Magnetic Resonance Angiography (MRA), Abdomen/Pelvis	4
Recommended Clinical Approach	4
Medical Necessity Criteria	4
Indications	4
Non-Indications	8
Level of Care Criteria	9
Procedure Codes (CPT/HCPCS)	9
Medical Evidence	10
References	11
Clinical Guideline Revision History/Information	13

Medical Necessity Criteria

Service: Magnetic Resonance Angiography (MRA), Abdomen/Pelvis

Recommended Clinical Approach

Magnetic resonance angiography (MRA) visualizes the blood vessels in the abdomen and pelvis. It aids in diagnosing and evaluating vascular conditions such as aneurysms, stenosis, occlusions, and vascular malformations. Unlike CT angiography, MRA does not use ionizing radiation yet provides detailed images of blood vessels and surrounding tissues. Magnetic resonance venography (MRV) of the abdomen and pelvis is a non-invasive imaging technique that uses magnetic resonance imaging (MRI) to visualize the veins in these regions. This method is particularly useful for evaluating venous disorders without exposing patients to ionizing radiation. MRA and MRV are less invasive than conventional X-ray digital subtraction angiography.¹

Medical Necessity Criteria

Indications

- **Magnetic resonance angiography (MRA), abdomen/pelvis** is considered appropriate if **ALL** of the following is **TRUE**¹⁻²:
- ◆ Ultrasound and CTA are contraindicated or inconclusive (e.g., body habitus for ultrasound, anaphylactic reaction due to IV contrast reaction, pregnancy, pediatric); **AND**
 - ◆ **ANY** of the following is **TRUE**:
 - Vascular conditions, known or suspected, including **ANY** of the following:
 - Suspected renal artery stenosis when surgical intervention is planned if diagnosed, including **ANY** of the following³:
 - ◆ Previous imaging (including ultrasound, captopril scintigraphy) indicates small kidney or unequal kidney sizes⁴⁻⁵; **OR**
 - ◆ Early-onset hypertension (age less than 35, diastolic greater than 110 mmHg)⁶; **OR**
 - ◆ Late-onset hypertension (age greater than 50)⁴⁻⁶; **OR**

- ◆ Renal artery bruit⁴⁻⁶; **OR**
- ◆ Malignant or accelerated hypertension⁶; **OR**
- ◆ Sudden development or worsening of hypertension⁶; **OR**
- ◆ Deterioration of renal function in response to angiotensin-converting enzyme inhibitors⁶; **OR**
- ◆ Generalized arteriosclerotic occlusive disease with hypertension⁶; **OR**
- ◆ Hypertension resistant to medication, and the patient must be currently taking **ALL** of the following at maximally tolerated doses⁶⁻⁷;
 - Long-acting calcium channel blocker; **AND**
 - Long-acting ace inhibitor or angiotensin receptor blocker (ARB); **AND**
 - Diuretic (e.g., loop or thiazide); **OR**
- Mesenteric ischemia or ischemic enteritis/colitis
- Vasculitis
- Vascular injury secondary to trauma; **OR**
- Thromboembolic Disease; **OR**
- Unrepaired aortic aneurysm, initial evaluation; **OR**
- Unrepaired aortic aneurysm, follow-up evaluation is based on aneurysm size when **ANY** of the following is **TRUE**⁸:
 - ◆ 3-3.9 cm, every 3 years; **OR**
 - ◆ 4-4.9 cm, annually; **OR**
 - ◆ 5-5.4 cm, every 6 months; **OR**
- Preoperative, postoperative, or pre-treatment evaluation for **ANY** of the following:
 - Evaluation post-endovascular repair (EVAR) or open repair of abdominal aortic aneurysm (AAA) and **ANY** of the following is **TRUE**^{4-5,8}:
 - ◆ Follow-up within one month; **OR**
 - ◆ Follow-up within one year; **OR**
 - ◆ Type II endoleak, every six months for up to two years after diagnosis: **OR**
 - ◆ Type II endoleak, annual followup; **OR**

- Planning for vascular surgery, interventional procedure; **OR**
- Other procedures involving arteries (e.g. Inferior epigastric arteries for breast reconstruction, UPJ obstruction, solid organ transplant); **OR**
- Retroperitoneal hematoma or hemorrhage; **OR**
- Known or suspected syndromes with increased risk of vascular anomalies including **ANY**, but not limited to, the following:
 - As a one-time screening for syndromes with a vascular component (e.g., fibromuscular dysplasia, Ehlers-Danlos syndrome, Marfan syndrome, neurofibromatosis, William’s syndrome, tuberous sclerosis); **OR**
 - Loeys Dietz (every two years for screening; surveillance as indicated depending on abnormalities found); **OR**
 - Other syndromes not otherwise specified, follow-up as clinical documentation supports; **OR**
- Repeat imaging (defined as repeat request following recent imaging of the same anatomic region with the same modality), in the absence of established guidelines, will be considered reasonable and necessary if **ANY** of the following is **TRUE**:
 - New or worsening symptoms, such that repeat imaging would influence treatment; **OR**
 - One-time clarifying follow-up of a prior indeterminate finding; **OR**
 - In the absence of change in symptoms, there is an established need for monitoring which would influence management.

→ **Magnetic resonance venography (MRV), abdomen/pelvis** is considered appropriate if **ALL** of the following are **TRUE**:

- ◆ Ultrasound and CT venography (CTV) are contraindicated or inconclusive (e.g., body habitus for ultrasound, anaphylactic reaction due to IV contrast reaction, pregnancy, pediatric); **AND**
- ◆ **ANY** of the following is **TRUE**:
 - Vascular conditions, known or suspected, including **ANY** of the following:
 - Diffuse unexplained lower extremity edema with negative or inconclusive ultrasound; **OR**
 - May-Thurner syndrome (iliac vein compression syndrome including pelvic CT venography)⁹; **OR**
 - Retroperitoneal hematoma or hemorrhage; **OR**
 - Large vein injury secondary to trauma; **OR**
 - Large vein thrombosis of the major abdominal or pelvic veins, including IVC, iliac, renal, portal, hepatic, and mesenteric veins, when Doppler ultrasound is inconclusive or indicates presence or complications; **OR**
 - Vascular invasion or displacement by tumor; **OR**
 - Detailed evaluation for Pelvic Congestion Syndrome; **OR**
 - Vascular mapping for organ donation; **OR**
 - Initial diagnostic, one-time pre or one-time post-treatment evaluation for treatment planning or evidence of clinical concern for **ANY** of the following:
 - Anastomotic integrity or stent patency; **OR**
 - Portal venous system (hepatic portal system) after Doppler ultrasound has been performed; **OR**
 - Vascular malformation; **OR**
 - Vascular mapping for organ donation or before procedure/surgery for planning purposes (including TIPS); **OR**
 - Repeat imaging (defined as repeat request following recent imaging of the same anatomic region with the same modality), in the absence of established guidelines, will be considered reasonable and necessary if **ANY** of the following is **TRUE**:

- New or worsening symptoms, such that repeat imaging would influence treatment; **OR**
- One-time clarifying follow-up of a prior indeterminate finding; **OR**
- In the absence of change in symptoms, there is an established need for monitoring which would influence management.

Non-Indications

→ **Magnetic resonance angiography (MRA), abdomen/pelvis** is not considered appropriate if **ANY** of the following is **TRUE**:

- ◆ The patient has undergone advanced imaging of the same body part within 3 months without undergoing treatment or developing new or worsening symptoms¹²; **OR**
- ◆ The patient has undergone advanced imaging of the same body part and for the same indication within 3 months, without being on treatment; **OR**
- ◆ If contrast is used, history of anaphylactic allergic reaction to gadolinium contrast media with detailed guidelines for use in patients with renal insufficiency; **OR**
- ◆ The patient has metallic clips on vascular aneurysms; **OR**
- ◆ Incompatible implantable devices (e.g., pacemakers, defibrillators, cardiac valves); **OR**
- ◆ Metallic foreign body in orbits/other critical area(s) or within the field of view and obscuring area of concern.

*NOTE: MRI in patients with claustrophobia should be requested at the discretion of the ordering provider.

**NOTE: MRI in pregnant patients should be requested at the discretion of the ordering provider and obstetric care provider.

Level of Care Criteria

Inpatient or Outpatient

Procedure Codes (CPT/HCPCS)

CPT/HCPCS Code	Code Description
72198	Magnetic resonance angiography (MRA) of pelvis, with contrast material
74185	Magnetic resonance angiography (MRA) of abdomen, with contrast material
C8900	Magnetic resonance angiography with contrast, abdomen
C8901	Magnetic resonance angiography without contrast, abdomen
C8902	Magnetic resonance angiography without contrast followed by with contrast, abdomen
C8918	Magnetic resonance angiography with contrast, pelvis
C8919	Magnetic resonance angiography without contrast, pelvis
C8920	Magnetic resonance angiography without contrast followed by with contrast, pelvis

Medical Evidence

Roditi et al. (2022) performed a review on abdominal and pelvic magnetic resonance angiography (MRA). The topics discussed include MRA for assessing renal vasculature in potential kidney donors and hypertensive patients, hepatic and mesenteric MRA for evaluating liver donors, individuals with portal hypertension, and those with chronic mesenteric ischemia. Pelvic MRA is also mentioned for pre-treatment planning in uterine fibroid embolization and patients with pelvic congestion syndrome. Abdominal wall MRA is also highlighted for planning breast reconstructive surgery.¹⁰

Chaikof et al. (2018) discuss updates to practice guidelines published by the Society for Vascular Surgery on the care of patients with an abdominal aortic aneurysm. Recommendations include surveillance imaging at 12-month intervals for AAA between 4.0 to 4.9 cm in diameter and utilizing the Vascular Quality Initiative mortality risk score for decision-making in aneurysm repair. Endovascular repair is also preferred for ruptured aneurysms. Color duplex ultrasound for postoperative surveillance after endovascular repair without complications is also recommended. Overall, the focus is to enhance decision-making and perioperative outcomes.⁸

Zucker et al. (2016) review noninvasive diagnostic imaging for assessing venous compression syndromes, including magnetic resonance venography (MRV). While the exam typically takes longer than CT scans, MRV offers the advantage of reducing ionizing radiation risks. Optimal timing is more easily achieved for venous contrast. Additionally, MRI enables non-contrast exams, which are safer for patients with renal insufficiency, who face a higher risk of nephrogenic systemic fibrosis with gadolinium contrast.¹¹

References

1. American College of Radiology (ACR). ACR–NASCI–SPR practice parameter for the performance of body magnetic resonance angiography (MRA). Published 2020. Accessed May 10, 2024. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Body-MR A.pdf>.
2. François CJ. Abdominal magnetic resonance angiography. *Nephrol Dial Transplant*. 2006 Apr;21(4):1104–8. doi: 10.1093/ndt/gfk062. PMID: 16431890.
3. Suvila K, Langén V, Cheng S, et al. Age of hypertension onset: Overview of research and how to apply in practice. *Curr Hypertens Rep*. 2020 Aug 27;22(9):68. doi: 10.1007/s11906-020-01071-z. PMID: 32852608; PMCID: PMC7452883.
4. Centers for Medicare and Medicaid Services (CMS). Local coverage determination: Magnetic resonance angiography (MRA)(L33633). Revision Effective Date October 1, 2019. Accessed May 10, 2024. <https://www.cms.gov/medicare-coverage-database/search.aspx>.
5. Centers for Medicare and Medicaid Services (CMS). Local coverage determination: Magnetic resonance angiography (MRA)(L34865). Revision Effective Date July 1, 2020. Accessed May 10, 2024. <https://www.cms.gov/medicare-coverage-database/search.aspx>.
6. Expert Panels on Urologic Imaging and Vascular Imaging, Harvin HJ, Verma N, et al. ACR appropriateness criteria - renovascular hypertension. *J Am Coll Radiol*. 2017 Nov;14(11S):S540–S549. doi: 10.1016/j.jacr.2017.08.040. PMID: 29101991.
7. Unger T, Borghi C, Charchar F, et al. 2020 International Society of Hypertension global hypertension practice guidelines. *Hypertension*. 2020 Jun;75(6):1334–1357. doi: 10.1161/HYPERTENSIONAHA.120.15026. PMID: 32370572.
8. Chaikof EL, Dalman RL, Eskandari MK, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *J Vasc Surg*. 2018 Jan;67(1):2–77.e2. doi: 10.1016/j.jvs.2017.10.044. PMID: 29268916.
9. Ibrahim W, Al Safran Z, Hasan H, et al. Endovascular management of may–thurner syndrome. *Ann Vasc Dis*. 2012;5(2):217–21. doi: 10.3400/avd.cr.12.00007. PMID: 23555515; PMCID: PMC3595863.
10. Roditi G, Wieben O, Prince MR, et al. MR angiography series: Abdominal and pelvic MR angiography. *Radiographics*. 2022 May–Jun;42(3):E94–E95. doi: 10.1148/rg.210224. PMID: 35245106.

11. Zucker EJ, Ganguli S, Ghoshhajra BB, et al. Imaging of venous compression syndromes. *Cardiovasc Diagn Ther*. 2016 Dec;6(6):519–532. doi: 10.21037/cdt.2016.11.19. PMID: 28123973; PMCID: PMC5220205.
12. Wasser EJ, Prevedello LM, Sodickson A, Mar W, Khorasani R. Impact of a real-time computerized duplicate alert system on the utilization of computed tomography. *JAMA Intern Med*. 2013;173(11):1024–1026. doi: 10.1001/jamainternmed.2013.543. PMID: 23609029.

Clinical Guideline Revision History/Information

Original Date: March 18, 2022		
Review History		
Version 2	8/2/2024	Annual review and policy restructure.
Version 3	10/30/2024	Edited repeat imaging criteria language.