



Cohere Medical Policy – Hip Arthroscopy

Clinical Guidelines for Medical Necessity Review

Version: 4
Revision Date: April 17, 2025

Important Notices

Notices & Disclaimers:

GUIDELINES SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION MAKING OF END USERS.

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine medical necessity of services (the "**Guidelines**") for informational purposes only, and solely for use by Cohere's authorized "**End Users**". These Guidelines (and any attachments or linked third party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

©2025 Cohere Health, Inc. All Rights Reserved.

Other Notices:

HCPCS® and CPT® copyright 2025 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

Guideline Information:

Specialty Area: Disorders of the Musculoskeletal System

Guideline Name: Cohere Medical Policy - Hip Arthroscopy

Date of last literature review: 1/16/2025

Document last updated: 4/16/2025

Type: ☒ Adult (18+ yo) | ☒ Pediatric (0-17 yo)

Table of Contents

Important Notices	2
Table of Contents	3
Medical Necessity Criteria	4
Service: Hip Arthroscopy	4
Recommended Clinical Approach:	4
Medical Necessity Criteria	4
Indications	4
Non-Indications	8
Level of Care Criteria	8
Procedure Codes (CPT/HCPCS)	8
Medical Evidence	10
References	12
Clinical Guideline Revision History/Information	17

Medical Necessity Criteria

Service: Hip Arthroscopy

Recommended Clinical Approach:

Hip arthroscopy is a minimally invasive procedure that offers visualization of the hip joint through an instrument known as an arthroscope. Hip arthroscopy is useful for clinically complex cases with an unclear diagnosis and can also be used to collect biopsies to assess for synovial disease. Injuries to the hip may also be repaired during arthroscopy, including labral tears and acute trauma resulting in loose bodies within the hip joint. The torn labrum may be repaired by removing labral pieces and then suturing the tear.

Femoroacetabular impingement (FAI) is another common indication for hip arthroscopy. It causes pain and limits activity because of the abnormal shape of the bones of the hip, which developed improperly during childhood skeletal growth. In general, arthroscopy is well-tolerated, safe, and leads to improved function and quality of life.¹

Medical Necessity Criteria

Indications

→ **Hip arthroscopy** is considered appropriate if **ANY** of the following is **TRUE**¹:

- ◆ **Diagnostic hip arthroscopy** is considered appropriate if the patient has **ALL** of the following^{1,2}:
 - A source of hip pain with an unclear diagnosis; **AND**
 - **ANY** of the following is **TRUE**:
 - Loose bodies; **OR**
 - Chondral lesion; **OR**
 - Synovial disease; **OR**
 - Adhesive capsulitis; **AND**
 - **ANY** of the following:
 - Acute pathology or acute injury (infection, trauma, gluteal tear) which is exempt from conservative care; **OR**

- Failure of conservative management for greater than 3 months, including **ALL** of the following^{1,2}:
 - ◆ Anti-inflammatory medications, non-opioid analgesics, or prescription medications (e.g., oral steroids, neuropathic pain medications) if not contraindicated; **AND**
 - ◆ Physical therapy or physician-directed exercise program; **AND**
 - ◆ **ANY** of the following:
 - Corticosteroid injection if medically appropriate; **OR**
 - Documentation that corticosteroid injection is contraindicated; **OR**
- ◆ **Arthroscopic labral debridement/repair** is considered appropriate with **ALL** of the following:
 - Labral tearing is visible on advanced imaging¹⁻⁴; **AND**
 - **ANY** of the following:
 - Acute traumatic tear; **OR**
 - Failure of conservative management for greater than 3 months, including **ALL** of the following:
 - ◆ Anti-inflammatory medications, non-opioid analgesics, or prescription medications (e.g., oral steroids, neuropathic pain medications) if not contraindicated; **AND**
 - ◆ Physical therapy or physician-directed exercise program; **AND**
 - ◆ **ANY** of the following:
 - Corticosteroid injection if medically appropriate; **OR**
 - Documentation that corticosteroid injection is contraindicated; **OR**
- ◆ **Trochanteric bursectomy** is considered appropriate when the patient has **ALL** of the following^{1,2}:
 - **ANY** of the following:
 - Trochanteric bursitis (greater trochanteric pain syndrome); **OR**
 - Greater trochanteric pain syndrome
 - Gluteal tendinopathy; **AND**

- Failure of conservative management for greater than 3 months, including **ALL** of the following:
 - Anti-inflammatory medications, non-opioid analgesics, or prescription medications (e.g., oral steroids, neuropathic pain medications) if not contraindicated; **AND**
 - Physical therapy or physician-directed exercise program; **AND**
 - **ANY** of the following:
 - ◆ Corticosteroid injection if medically appropriate; **OR**
 - ◆ Documentation that corticosteroid injection is contraindicated; **OR**
- ◆ **Gluteal repair** is considered appropriate when the patient has imaging findings of **ANY** of the following^{1,2}:
 - Gluteal tear; **OR**
 - **ALL** of the following:
 - Gluteal tendinopathy; **AND**
 - Failure of conservative management for greater than 3 months, including **ALL** of the following:
 - ◆ Anti-inflammatory medications, non-opioid analgesics, or prescription medications (e.g., oral steroids, neuropathic pain medications) if not contraindicated; **AND**
 - ◆ Physical therapy or physician-directed exercise program; **AND**
 - ◆ **ANY** of the following:
 - Corticosteroid injection if medically appropriate; **OR**
 - Documentation that corticosteroid injection is contraindicated; **OR**
- ◆ **Arthroscopic osteochondroplasty** is considered appropriate with **ALL** of the following^{1,2,5}:
 - Positive impingement sign with pain (hip is flexed to 90 degrees, adducted, and internally rotated); **AND**
 - Moderate to severe persistent hip or groin pain that limits activity and is worse with hip flexion; **AND**
 - Advanced imaging demonstrates **ANY** of the following:

- Femoroacetabular impingement (FAI) impingement with evidence of CAM impingement (alpha angle greater than 50 degrees); **OR**
- Pincer impingement (coxa profunda or acetabular retroversion); **OR**
- Subspine impingement⁶; **OR**
- Ischiofemoral impingement^{6,7}; **OR**
- Residual impingement remains after first arthroscopy¹⁸; **AND**
- Failure of conservative management for greater than 3 months, including **ALL** of the following:
 - Anti-inflammatory medications, non-opioid analgesics, or prescription medications (e.g., oral steroids, neuropathic pain medications) if not contraindicated; **AND**
 - Physical therapy or physician-directed exercise program; **AND**
 - **ANY** of the following:
 - ◆ Corticosteroid injection if medically appropriate; **OR**
 - ◆ Documentation that corticosteroid injection is contraindicated; **OR**
- ◆ **Other arthroscopic intervention of the hip** is considered appropriate for **ANY** of the following^{1,2}:
 - Ligamentum teres disorder⁹; **OR**
 - Proximal hamstring injury indicated by **ANY** of the following⁹⁻¹¹:
 - 2-tendon injury with greater than 2 centimeters of retraction; **OR**
 - 3-tendon injury; **OR**
 - Acute infection/joint sepsis¹; **OR**
 - Acute trauma with associated findings on imaging (e.g. loose body noted after hip dislocation); **OR**
 - Acute gluteal tear (partial or full-thickness) within three months of injury¹²; **OR**
 - **ALL** of the following:
 - **ANY** of the following:
 - ◆ Sciatic nerve entrapment¹³; **OR**

- ◆ Psoas tendon disorder; **OR**
- ◆ Atraumatic instability that is felt to be resolvable with arthroscopic capsular plication^{14,15}; **AND**
- Failure of conservative management for greater than 3 months, including **ALL** of the following:
 - ◆ Anti-inflammatory medications, non-opioid analgesics, or prescription medications (e.g., oral steroids, neuropathic pain medications) if not contraindicated; **AND**
 - ◆ Physical therapy or physician-directed exercise program; **AND**
 - ◆ **ANY** of the following:
 - Corticosteroid injection if medically appropriate; **OR**
 - Documentation that corticosteroid injection is contraindicated.

Non-Indications

- **Hip arthroscopy** is not considered appropriate when **ANY** of the following is **TRUE**^{1,2}:
- ◆ Ankylosis of the hip; **OR**
 - ◆ Advanced hip osteoarthritis (Tönnis grade 2 or 3).

Level of Care Criteria

Outpatient

Procedure Codes (CPT/HCPCS)

CPT/HCPCS Code	Code Description/Definition
27299	Unlisted procedure, pelvis or hip joint
29860	Diagnostic arthroscopy of hip joint; Diagnostic arthroscopy of hip joint with synovial biopsy
29861	Surgical arthroscopy of hip with removal of foreign body; Surgical arthroscopy of hip with removal of loose body

29862	Surgical arthroscopy of hip with debridement of articular cartilage; Surgical arthroscopy of hip with debridement of articular cartilage, abrasion arthroplasty, and resection of labrum; Surgical arthroscopy of hip with debridement of articular cartilage, and abrasion arthroplasty; Surgical arthroscopy of hip with shaving of articular cartilage, abrasion arthroplasty, and resection of labrum
29863	Surgical arthroscopy of hip with synovectomy
29914	Surgical arthroscopy of hip with femoroplasty; Surgical arthroscopy of hip with femoroplasty for cam lesion
29915	Surgical arthroscopy of hip with acetabuloplasty; Surgical arthroscopy of hip with acetabuloplasty for pincer lesion
29916	Surgical arthroscopy of hip with labral repair
29999	Unlisted arthroscopic procedure

Medical Evidence

As compared to traditional open procedures, hip arthroscopy is minimally invasive, decreases the risk of neurovascular injury, and requires a shorter recovery time. A 2023 systematic review of patients aged 50 years and older saw these benefits persist among an older population, albeit alongside a pronounced risk for revision surgery and subsequent total hip arthroplasty (THA). The systematic review consisted of 6,696 patients across 17 studies who underwent primary hip arthroscopy between 2015 and 2021. Up to 10.8% of patients underwent surgical revision, while up to 34% experienced ultimate conversion to THA. The authors concluded that, although postoperative patient-reported outcomes were significantly improved as compared to baseline, careful patient selection is particularly important for patients of advanced age in order to appropriately weigh the very real risk of eventual revision or conversion to THA.¹⁶

Conducted in 2018, the UK FASHIoN study was an assessor-blinded, randomized controlled trial involving 348 patients with femoroacetabular hip impingement syndrome (FAI), 171 of whom underwent hip arthroscopy, while the other 177 were treated with conservative care alone. The authors found that hip arthroscopy conferred a significant improvement in quality of life at one year after surgical intervention. They noted that these results solidified the use of arthroscopy to treat FAI, which has grown in clinical popularity in recent years.¹⁷

A 2021 systematic review addressed the role of hip arthroscopy in the management of gluteal tendinopathy. The authors evaluated 27 studies, including 6 randomized controlled trials, with an aggregate of 1103 patients. Bursectomy was felt to be a valuable treatment option for patients who failed conservative measures. Surgical repair of partial-thickness and full-thickness gluteal tears was encouraged to be considered early in the course of injury due to the lack of data regarding nonoperative treatment of grade 3 and grade 4 (partial; full) tendinopathy.¹²

The American Association of Orthopaedic Surgeons (AAOS) has issued position statements pertaining to hip arthroscopy. Information statement 1047, published in 2016, acknowledges the increased patient safety risks conferred by tobacco use – including increased ventilatory support, myocardial infarction, cardiac arrest, cerebrovascular accident, sepsis, and death.¹⁸ The AAOS states that patients who are active smokers may reduce these risks through cessation of smoking prior to surgery; they also note the special role orthopaedic surgeons play in counseling patients on the benefits of reduced or eliminated tobacco use before surgery. Importantly, unconfirmed cessation is not endorsed as a hard stop to surgery; rather, the surgeon’s unique role as an advocate for preoperative smoking cessation is emphasized. Statements 1040 and 1184 discuss the impact of obesity on musculoskeletal conditions.^{19,20} Patients with morbid obesity (BMI of 40 or above) are encouraged to participate in a weight loss program, obtain weight reduction resources through their physician, rectify nutritional deficiencies, and consider a delay in surgical treatment if it would facilitate participation in weight loss interventions that may improve surgical outcomes. Statement 1040 notes that individuals with obesity face an increased risk for sports injuries, and that when such injuries are treated arthroscopically, the procedure may be more technically difficult because of the loss of superficial landmarks. Questions remain as to whether functional results are affected by obesity. Further, the authors note the risks associated with general anesthesia for patients with obesity and emphasize the importance of adequate patient positioning and padding to avoid pressure ulcers, nerve palsies, and compartment syndromes, which are more common among obese patients. In general, obesity is associated with greater risk of premature complications and mortality during the perioperative period. The AAOS endorses compassionate, risk-informed patient counseling for obese patients who are considering surgery. Careful screening and appropriate referral to nutrition or endocrine care is also endorsed. Statement 1184 reinforces the risks associated with obesity in the setting of orthopaedic care and similarly encourages adequate patient counseling prior to surgery.

Social determinants of health remain an important area of ongoing orthopaedic surgery research, with recent literature raising questions regarding the healthcare disparities that may be potentiated by care limitations based on obesity and smoking status/nicotine dependence.²¹⁻²³ Other ongoing research interrogates the impacts that biological sex, race,

and socioeconomic status have on hip arthroscopy utilization and outcomes.²⁴⁻³³

References

1. Phillips BB, Mihalko MJ. Arthroscopy of the lower extremity. Azar FM, Beaty JH, editors. In: *Campbell's Operative Orthopaedics*. 14th ed. Philadelphia, PA: Elsevier; 2021:2576–2662.e8
2. Ross JR, Larson CM, Bedi A. Indications for hip arthroscopy. *Sports Health*. 2017 Sep/Oct;9(5):402–413. doi: 10.1177/1941738117712675. PMID: 28678628; PMCID: PMC5582699
3. Groh MM, Herrera J. A comprehensive review of hip labral tears. *Curr Rev Musculoskelet Med*. 2009 Jun;2(2):105–17. doi: 10.1007/s12178-009-9052-9. PMID: 19468871; PMCID: PMC2697339
4. Woyski D, Mather RC. Surgical treatment of labral tears: Debridement, repair, reconstruction. *Curr Rev Musculoskelet Med*. 2019 Sep;12(3):291–299. doi: 10.1007/s12178-019-09575-1. PMID: 31346980; PMCID: PMC6684681
5. Wilson AS, Cui Q. Current concepts in management of femoroacetabular impingement. *World J Orthop*. 2012 Dec 18;3(12):204–11. doi: 10.5312/wjo.v3.i12.204. PMID: 23362464; PMCID: PMC3557322
6. Ryan MK. Addressing Extra-articular impingement in the athlete. *Oper Tech Sports Med*. 2024 Sep 1;32(3):151109
7. Wu WT, Chang KV, Mezian K, Naňka O, Ricci V, et al. Ischiofemoral impingement syndrome: Clinical and imaging/guidance issues with special focus on ultrasonography. *Diagn*. 2022 Dec 31;13(1):139
8. Gwathmey FW, Jones KS, Thomas Byrd JW. Revision hip arthroscopy: Findings and outcomes. *J Hip Preserv Surg*. 2017 Dec;4(4):318–23
9. Fletcher AN, Cheah JW, Nho SJ, Mather RC. Proximal hamstring injuries. *Clin Sports Med*. 2021 Apr 1;40(2):339–61
10. Ahmad CS, Redler LH, Ciccotti MG, Maffulli N, Longo UG, Bradley J. Evaluation and management of hamstring injuries. *Amer J Sports Med*. 2013 Dec;41(12):2933–47
11. Allahabadi S, Salazar LM, Obioha OA, Fenn TW, Chahla J, Nho SJ. Hamstring injuries: A current concepts review: Evaluation, nonoperative treatment, and surgical decision making. *Amer J Sports Med*. 2024 Mar;52(3):832–44

12. Ladurner A, Fitzpatrick J, O'Donnell JM. Treatment of gluteal tendinopathy: A systematic review and stage-adjusted treatment recommendation. *Ortho J Sports Med*. 2021 Jul 29;9(7):23259671211016850
13. Perets I, Rybalko D, Mu BH, Friedman A, Morgenstern DR, Domb BG. Hip arthroscopy: Extra-articular procedures. *Hip Int*. 2019 Jul;29(4):346–54
14. Kalisvaart MM, Safran MR. Hip instability treated with arthroscopic capsular plication. *Knee Surg, Sports Traumatol, Arthrosc*. 2017 Jan;25:24–30
15. Maldonado DR, Chen JW, Yelton MJ, Rosinsky PJ, Shapira J, Brayboy C, Lall AC, Domb BG. Achieving successful outcomes of hip arthroscopy in the setting of generalized ligamentous laxity with labral preservation and appropriate capsular management: a propensity matched controlled study. *Amer J Sports Med*. 2020 Jun;48(7):1625–35
16. Shanmugaraj A, Kumar MV, Al Naji O, Simunovic N, Philippon MJ, Ayeni OR. Hip arthroscopy improves outcomes with moderate conversion to total hip arthroplasty rates in patients aged 50 years or older: a systematic review. *Arthrosc*. 2023 Jun 1;39(6):1539–51
17. Griffin DR, Dickenson EJ, Wall PD, et al. Hip arthroscopy versus best conservative care for the treatment of femoroacetabular impingement syndrome (UK FASHIoN): a multicentre randomised controlled trial. *Lancet*. 2018 Jun 2;391(10136):2225–35. DOI: 10.1016/S0140-6736(18)31202-9
18. American Academy of Orthopaedic Surgeons. Information Statement 1047: Tobacco use and orthopaedic surgery. Published February 2016. Accessed January 3, 2025. <https://www.aaos.org/globalassets/about/bylaws-library/information-statements/1047-tobacco-use-and-orthopaedic-surgery-3.pdf>
19. American Academy of Orthopaedic Surgeons. Information Statement 1040: Obesity and musculoskeletal care. Published June 2022. Accessed January 3, 2025. <https://www.aaos.org/globalassets/about/bylaws-library/information-statements/1040-obesity-and-musculoskeletal-care.pdf>
20. American Academy of Orthopaedic Surgeons. Information Statement 1184: The impact of obesity on bone and joint health. Published March 2015. Accessed January 3, 2025. <https://www.aaos.org/contentassets/1cd7f41417ec4dd4b5c4c48532183b96/1184-the-impact-of-obesity-on-bone-and-joint-health1.pdf>

21. Anaspure OS, Patel S, Baumann AN, Lenz T, Pascual-Leone N, Anastasio AT, Lau BC. Examining the role of smoking on clinical outcomes after arthroscopic surgery of the hip: A systematic review and meta-analysis. *Euro J of Ortho Surg & Traumatol*. 2025 Dec;35(1):1-7
22. Lee MS, Jimenez AE, Owens JS, Curley AJ, Paraschos OA, Maldonado DR, Lall AC, Domb BG. Comparison of outcomes between nonsmokers and patients who discontinued smoking 1 month before primary hip arthroscopy: A propensity-matched study with minimum 2-year follow-up. *Ortho J Sports Med*. 2022 Jun 8;10(6):23259671221097372
23. Emara AK, Grits D, Samuel LT, Acuña AJ, Rosneck JT, Kamath AF. Hip arthroscopy in smokers: A systematic review of patient-reported outcomes and complications in 18,585 cases. *Amer J Sports Med*. 2021 Mar;49(4):1101-8
24. Niehaus R, Zingg PO, Hoch A, Luttenberger M, Stefan R. Hip arthroscopy versus total hip arthroplasty—A study on patients with obesity above 40 years of age. *Clin Obesity*. 2023 Oct;13(5):e12590
25. Kuroda Y, Hashimoto S, Saito M, Hayashi S, Nakano N, Matsushita T, Niikura T, Kuroda R, Matsumoto T. Obesity is associated with less favorable outcomes following hip arthroscopic surgery: A systematic review and meta-analysis. *Knee Surg, Sports Traumatol, Arthrosc*. 2021 May;29:1483-93
26. Gupta A, Redmond JM, Hammarstedt JE, Stake CE, Domb BG. Does obesity affect outcomes in hip arthroscopy? A matched-pair controlled study with minimum 2-year follow-up. *Amer J of Sports Med*. 2015 Apr;43(4):965-71
27. Lee JS, Rachala RR, Gillinov SM, Siddiq BS, Dowley KS, Cherian NJ, Martin SD. Relationship between neighborhood-level socioeconomic status and functional outcomes after hip arthroscopy. *Amer J of Sports Med*. 2024 Oct;52(12):3054-64
28. Amen TB, Chatterjee A, Rudisill SS, Joseph GP, Nwachukwu BU, Ode GE, Williams III RJ. National patterns in utilization of knee and hip arthroscopy: An analysis of racial, ethnic, and geographic disparities in the United States. *Ortho J Sports Med*. 2023 Aug 16;11(8):23259671231187447
29. Buerba RA, Dalton J, Sadhwani S, Schulz W, Atte AC, Vyas D. Hip arthroscopy utilization disparities and complications amongst ethnic groups. *INQUIRY: J Health Car*. 2024 Oct;61:00469580241282644

30. Gillinov SM, LaPorte ZL, Lee JS, Siddiq BS, Dowley KS, Cherian NJ, Eberlin CT, Kucharik MP, Martin SD. Social determinants of health disparities increase 5-year revision rates but not postoperative complications after primary hip arthroscopy. *Arthrosc*. 2024 Aug 20
31. Saks BR, Ouyang VW, Domb ES, Jimenez AE, Maldonado DR, Lall AC, Domb BG. Equality in hip arthroscopy outcomes can be achieved regardless of patient socioeconomic status. *Amer J Sports Med*. 2021 Dec;49(14):3915-24
32. Parvaresh K, Rasio JP, Wichman D, Chahla J, Nho SJ. The influence of body mass index on outcomes after hip arthroscopy for femoroacetabular impingement syndrome: Five-year results in 140 patients. *Amer J Sports Med*. 2021 Jan;49(1):90-6
33. McCormack TJ, Vopat ML, Rooker J, Tarakemeh A, Baker J, Templeton KJ, Mulcahey MK, Mullen SM, Schroepfel JP, Vopat BG. Sex-based differences in outcomes after hip arthroscopic surgery for femoroacetabular impingement: A systematic review. *Ortho J Sports Med*. 2022 Nov 24;10(11):23259671221137857
34. Kyin C, Maldonado DR, Go CC, Shapira J, Lall AC, Domb BG. Mid-to long-term outcomes of hip arthroscopy: A systematic review. *Arthros*. 2021 Mar 1;37(3):1011-25
35. Snaebjörnsson T, Anari SS, Lindman I, Desai N, Ståhlman A, Ayeni OR, Öhlin A. Most elite athletes who underwent hip arthroscopy for femoroacetabular impingement syndrome did not return to the same level of sport, but the majority were satisfied with the outcome of surgery. *Arthrosc, Sports Med, and Rehab*. 2022 Jun 1;4(3):e899-906
36. Nawabi DH, Bedi A, Tibor LM, Magennis E, Kelly BT. The demographic characteristics of high-level and recreational athletes undergoing hip arthroscopy for femoroacetabular impingement: A sports-specific analysis. *Arthrosc*. 2014 Mar 1;30(3):398-405
37. National Institute for Health and Care Excellence (NICE). Osteoarthritis in over 16s: Diagnosis and management [NG226]. Published October 19, 2022. Accessed July 1, 2024. www.nice.org.uk/guidance/ng226
38. Briggs KK, Bolia IK. Hip arthroscopy: An evidence-based approach. *Lancet*. 2018 Jun 2;391(10136):2189-90. DOI: 10.1016/S0140-6736(18)31218-2
39. Zhu Y, Su P, Xu T, Zhang L, Fu W. Conservative therapy versus arthroscopic surgery of femoroacetabular impingement syndrome (FAI): A systematic review and meta-analysis. *J Orthop Surg Res*. 2022 Jun 3;17(1):296. <https://doi.org/10.1186/s13018-022-03187-1>

40. Mok TN, He QY, Teng Q, et al. Arthroscopic hip surgery versus conservative therapy on femoroacetabular impingement syndrome: A meta-analysis of RCTs. *Orthopaedic Surg.* 2021 Aug;13(6):1755–64. <https://doi.org/10.1111/os.13099>
41. Mansell NS, Rhon DI, Meyer J, Slevin JM, Marchant BG. Arthroscopic surgery or physical therapy for patients with femoroacetabular impingement syndrome: A randomized controlled trial with 2-year follow-up. *Am J Sports Med.* 2018 May;46(6):1306–14. <https://doi.org/10.1177/0363546517751912>
42. Anzillotti G, Iacomella A, Grancagnolo M, et al. Conservative vs. surgical management for femoro-acetabular impingement: a systematic review of clinical evidence. *J Clin Med* 2022 Oct 2;11(19):5852. <https://doi.org/10.3390/jcm11195852>
43. Shahpari O, Mortazavi J, Ebrahimzadeh MH, Bagheri F, Mousavian A. Role of hip arthroscopy in the treatment of avascular necrosis of the hip: A systematic review. *Arch Bone and Joint Surg.* 2022 Jun;10(6):480
44. Yee C, Wong M, Cohen D, et al. Labral tears and chondral lesions are common comorbidities identified during endoscopic repair of gluteal tendon tears for greater trochanteric pain syndrome: a systematic review. *Arthrosc.* 2023 Mar 1;39(3):856–64. <https://doi.org/10.1016/j.arthro.2022.06.031>
45. Suarez-Ahedo C, Camacho-Galindo J, López-Reyes A, et al. A comprehensive review of hip arthroscopy techniques and outcomes. *SAGE Open Med.* 2024 Jan;12:20503121231222212. <https://doi.org/10.1177/20503121231222212>
46. Awad MA, Bajwa AK, Slaunwhite E, Logan KJ, Wong IH. Indications for hip arthroscopy in pediatric patients: A systematic review. *J Hip Preserv Surg.* 2019 Dec;6(4):304–15
47. Patricio Cordeiro TT, Rocha EA, Scattone Silva R. Effects of exercise-based interventions on gluteal tendinopathy. Systematic review with meta-analysis. *Sci Reports.* 2024 Feb 9;14(1):3343
48. Cargnelli S, Catapano M, Peterson D, et al. Efficacy of hip arthroscopy for the management of septic arthritis: A systematic review. *Arthrosc.* 2015 Jul 1;31(7):1358–70
49. Balato G, de Matteo V, Ascione T, et al. Management of septic arthritis of the hip joint in adults. A systematic review of the literature. *BMC MSK Dis.* 2021 Nov;22:1–3

50.Knapik DM, Farivar D, Kunze KN, Gilat R, Nho SJ, Chahla J. Indications and outcomes after ligamentum teres reconstruction: A systematic review. *Arthrosc, Sports Med, and Rehab*. 2021 Jun 1;3(3):e939–49

Clinical Guideline Revision History/Information

Original Date: May 22, 2020		
Review History		
Version 2	12/29/2023	
Version 3	9/20/2024	Updated language regarding conservative treatment
Version 4	4/17/2025	<p>Annual policy review & restructure:</p> <ul style="list-style-type: none">• Adjusted Recommended Clinical Approach to current format• Added indications for gluteal repair and hip bursectomy based on physician feedback• Removed weight reduction requirement in accordance with professional society guidelines• Added indications: Acute infection/joint sepsis; acute trauma with findings documented on imaging (e.g. loose body noted after hip dislocation); acute gluteal tear within 3 months of injury; proximal hamstring injury; extra-articular impingement; sciatic nerve entrapment; ligamentum teres disorder; capsular plication• Updated references

		<ul style="list-style-type: none">• Updated medical evidence section• Conservative care language modified to reflect non-opioid pain control.• Modified steroid injection language for clarity.
--	--	---