



Cohere Medical Policy - Computed Tomography (CT), Face/Sinus

Clinical Guidelines for Medical Necessity Review

Version: 2
Effective Date: August 20, 2024

Important Notices

Notices & Disclaimers:

GUIDELINES ARE SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION-MAKING OF END USERS.

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine the medical necessity of services (the "**Guidelines**") for informational purposes only, and solely for use by Cohere's authorized "**End Users**". These Guidelines (and any attachments or linked third-party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision-making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

©2024 Cohere Health, Inc. All Rights Reserved.

Other Notices:

HCPCS® and CPT® copyright 2024 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

Guideline Information:

Specialty Area: Diagnostic Imaging

Guideline Name: Cohere Medical Policy - Computed Tomography (CT), Face/Sinus

Date of last literature review: 8/16/2024

Document last updated: 8/19/2024

Type: Adult (18+ yo) | Pediatric (0-17 yo)

Table of Contents

Important Notices	2
Table of Contents	3
Medical Necessity Criteria	4
Service: Computed Tomography (CT), Face/Sinus	4
Recommended Clinical Approach	4
Medical Necessity Criteria	4
Indications	4
Non-Indications	7
Disclaimer on Radiation Exposure in Pediatric Population	7
Level of Care Criteria	8
Procedure Codes (CPT/HCPCS)	8
Medical Evidence	9
References	10
Clinical Guideline Revision History/Information	12

Medical Necessity Criteria

Service: Computed Tomography (CT), Face/Sinus

Recommended Clinical Approach

Computed tomography (CT) of the face/sinus is an advanced imaging modality best utilized per institutional ENT, oncologic, infectious, disease, and radiologic protocols. The specialty guidelines may be consulted before ordering and can direct appropriate contrast utilization patterns.

Medical Necessity Criteria

Indications

→ **Computed tomography (CT), face/sinus (including soft tissues of the extracranial head and neck)** is considered appropriate for **ANY** of the following:

- ◆ Conditions, known or suspected, including **ANY** of the following¹:
 - Anatomic abnormalities (e.g. deviated septum), suspected as a cause of patient symptoms and surgical management is being considered; **OR**
 - Bell's palsy or other facial nerve abnormalities requiring evaluation of the extracranial portion of the nerve (MRI is contraindicated or cannot be performed); **OR**
 - Congenital conditions and craniofacial abnormalities²; **OR**
 - Sinusitis when, if infective, and **ANY** of the following is **TRUE**²⁻⁵:
 - Four or more acute episodes per year and surgery/biologic therapy are contemplated; **OR**
 - Not resolving despite 2 courses of antibiotics; **OR**
 - A complication is suspected (e.g., abscess formation, involvement of adjacent structures such as orbits, cavernous sinus, or intracranial); **OR**
 - The patient is immunocompromised, and invasive fungal sinusitis is suspected; **OR**
 - Allergic fungal sinusitis (AFS) suspected, with failed medical treatment or surgery is contemplated; **OR**

- Chronic rhinosinusitis, symptomatic (discharge, congestion, anosmia, pain), severity staging or restaging when management change is contemplated; **OR**
- Suspected osteonecrosis when the patient is on bisphosphonates or has had radiation; **OR**
- Osteomyelitis; **OR**
- Odontogenic infections with suspected complications (e.g., abscess formation, facial swelling, nerve, sinus involvement); **OR**
- Unexplained facial swelling (e.g. over the mandible); **OR**
- Foreign-body (suspected), clinically or seen on prior imaging; **OR**
- Neoplastic conditions for initial staging, treatment planning, response assessment, and surveillance; **OR**
- Lymphadenopathy with failure of conservative management (e.g., antibiotics, anti-inflammatory) that is documented for a period of greater than two weeks for reactive adenopathy; **OR**
- Sinonasal polyposis detected on nasal endoscopy with **ALL** of the following^{3,6}:
 - The patient is symptomatic; **AND**
 - No relief with appropriate medical therapy (e.g., corticosteroids, antihistamines, antibiotics); **AND**
 - Surgical intervention or biologic therapy is being contemplated; **OR**
- Known sinonasal polyposis with complications suspected, (e.g., involvement of the orbits); **OR**
- Non-infectious inflammatory involvement of the sinus is suspected based on clinical history and symptoms (e.g., history of granulomatosis with polyangiitis)¹; **OR**
- Salivary stones, suspected¹; **OR**
- Salivary gland inflammation (sialadenitis)¹; **OR**
- Trigeminal neuralgia with **ANY** of the following²:
 - MRI is contraindicated or cannot be performed; **OR**
 - Atypical features present (e.g., symptoms outside of typical short-duration trigeminal nerve distribution pain); **OR**
 - Age less than 40; **OR**

- Failure of conservative management with failure of at least two concurrent agents (e.g., gabapentin, duloxetine) and surgery is being considered; **OR**
 - Vascular malformations (e.g., arterio-venous malformations)^{1,8}; **OR**
- ◆ For evaluation of **ANY** of the following symptoms when applicable:
 - Anosmia with **ANY** of the following⁹:
 - Persistent anosmia with nondiagnostic endoscopy; **OR**
 - Abnormal endoscopy with further evaluation needed; **OR**
 - Known or suspected neoplasm; **OR**
 - History of head or facial trauma; **OR**
 - Cerebrospinal fluid (CSF) leak, confirmed on testing or strong clinical history (e.g., prior trauma or CSF leak that increases after Valsalva maneuvers)³; **OR**
 - Epistaxis with failure of conservative management (e.g., nasal packing/tampon, cautery, etc); **OR**
 - Epistaxis with detection of mass, polyp, or other pathology on examination that requires further evaluation¹; **OR**
- ◆ Preoperative, postoperative, and pre-treatment evaluation for surgery, radiation, or chemotherapy; **OR**
- ◆ Maxillo-facial trauma based on history, swelling, prior imaging, and need for further evaluation; **OR**
- ◆ Repeat imaging of a specific area or structure using the same imaging modality (in the absence of an existing follow-up guideline) is considered appropriate when **ALL** of the following is **TRUE**:
 - There is documented clinical necessity; **AND**
 - Prior imaging results of the specific area or structure, obtained using the same imaging modality, must be documented and available for comparison; **AND**
 - **ANY** of the following is **TRUE**:
 - A change in clinical status (e.g., worsening symptoms or the emergence of new symptoms, that may influence the treatment approach); **OR**
 - The requirement for interval reassessment, which may alter the treatment plan; **OR**

- One-time follow-up of a prior indeterminate finding to assess for interval change; **OR**
- The need for re-imaging either before or after performing an invasive procedure.

Non-Indications

→ **Computed tomography (CT), face/sinus (including soft tissues of the extracranial head and neck) with contrast** is not considered appropriate if **ANY** of the following is **TRUE**¹⁰:

- ◆ The patient has undergone advanced imaging of the same body part within 3 months without undergoing treatment or developing new or worsening symptoms; **OR**
- ◆ If contrast is used, history of anaphylactic allergic reaction to iodinated contrast media.

*NOTE: The referring professional and radiologist should discuss the risks and benefits of contrast media administration, including possible prophylaxis, in patients with chronic or worsening kidney disease or severe renal failure.

**NOTE: CT in pregnant patients should be requested at the discretion of the ordering provider and obstetric care provider.

***NOTE: CT in patients with claustrophobia should be requested at the discretion of the ordering provider.

Disclaimer on Radiation Exposure in Pediatric Population

Due to the heightened sensitivity of pediatric patients to ionizing radiation, minimizing exposure is paramount. At Cohere, we are dedicated to ensuring that every patient, including the pediatric population, has access to appropriate imaging following accepted guidelines. Radiation risk is dependent mainly on the patient's age at exposure, the organs exposed, and the patient's sex, though there are other variables. The following technical guidelines are provided to ensure safe and effective imaging practices:

Radiation Dose Optimization: Adhere to the lowest effective dose principle for pediatric imaging. Ensure that imaging protocols are specifically tailored for pediatric patients to limit radiation exposure.¹¹⁻¹²

Alternative Modalities: Prioritize non-ionizing imaging options such as ultrasound or MRI when clinically feasible, as they are less likely to expose the

patient to ionizing radiation. For instance, MRI or ultrasound should be considered if they are more likely to provide an accurate diagnosis than CT, fluoroscopy, or radiography.¹¹⁻¹²

Cumulative Dose Monitoring: Implement systems to track cumulative radiation exposure in pediatric patients, particularly for those requiring multiple imaging studies. Regularly reassess the necessity of repeat imaging based on clinical evaluation.¹¹⁻¹²

CT Imaging Considerations: When CT is deemed the best method for achieving a correct diagnosis, use the lowest possible radiation dose that still yields reliable diagnostic images.¹¹⁻¹²

Cohere Imaging Gently Guideline

The purpose of this guideline is to act as a potential override when clinically indicated to adhere to Imaging Gently and Imaging Wisely guidelines and As Low As Reasonably Possible (ALARA) principles.

Level of Care Criteria

Inpatient and Outpatient

Procedure Codes (CPT/HCPCS)

CPT/HCPCS Code	Code Description
70486	Computed tomography (CT), maxillofacial area; without contrast material
70487	Computed tomography (CT), maxillofacial area; with contrast material(s)
70488	Computed tomography (CT), maxillofacial area; without contrast material, followed by contrast material(s) and further sections
76380	Computed tomography, limited or localized follow-up study

Medical Evidence

Bedernik et al. (2022) conducted a randomized control trial (RCT) to assess image quality by comparing single-energy computed tomography (SECT) with automated tube voltage adaptation (TVA) to dual-energy CT (DECT) weighted average images. A total of 80 patients underwent SECT or radiation dose-matched DECT. The effective radiation dose (ED) showed no significant difference between the SECT and DECT study groups. Compared to the SECT group, the DECT group exhibited significantly higher contrast-to-noise ratio differences (CNRD) for jugular veins relative to fatty tissue and muscle tissue relative to fatty tissue. However, the CNRD for jugular veins relative to muscle tissue was comparable between groups. Image artifacts were also less pronounced, and overall diagnostic acceptability was higher in the DECT group. Overall, DECT-weighted average images demonstrate superior objective and subjective image quality compared to SECT performed with TVA in head and neck imaging.¹³

Smith-Bindman et al. (2020) performed an RCT to study the efficacy of interventions to lower the amount of radiation patients are exposed to. The RCT included 864,080 adults at 100 facilities who underwent a CT scan, including CT Head (n = 1,156,657 scans). The study included two primary measures: the percentage of high-dose CT scans and the average effective dose administered at the facility level. The study's secondary measure included the doses received by specific organs. Outcomes were assessed with respect to the impact of the interventions and outcomes post-intervention. Data were contrasted with pre-intervention data, utilizing hierarchical generalized linear models that accounted for temporal patterns and patient attributes. In conclusion, data regarding CT radiation dosage and practical recommendations may improve quality, including significant dose reductions, especially for organ-specific doses.¹⁴

Aulino et al. (2018) report on a clinical trial that focused on an assessment tool for the late effect continuum of lymphedema and fibrosis (LEF) for patients with head and neck cancer (HNC) undergoing CT. The tool evaluates areas of soft tissue damage resulting from tumors, surgical interventions, or radiation therapy. The tool analyzed CT scans taken before and after treatment in 10 patients with HNC. The finalized tool comprised 11 elements, including the assessment of fat stranding at six specific sites, epiglottic thickness measurement, and measurement of prevertebral soft tissue thickness at C3. A total of 176 CT scans from the 20 patients (with a range of 4-14 scans per patient) were evaluated. The final version of the LEF assessment tool (CT-LEFAT) offers a standardized approach to assess critical sites affected by soft tissue damage. Studies continue to evaluate reliability and validity.¹⁵

References

1. Expert Panel on Neurologic Imaging, Sharma A, Kirsch CFE, et al. ACR appropriateness criteria – hearing loss and/or vertigo. Published 2018. Accessed August 17, 2024. <https://acsearch.acr.org/docs/69488/Narrative/>.
2. American College of Radiology (ACR), American Society of Neuroradiology (ASNR), Society for Pediatric Radiology (SPR). ACR–ASNR–SPR practice parameter for the performance of computed tomography (CT) of the extracranial head and neck – resolution 5. Updated 2021. Accessed August 14, 2024. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CT-Head-Neck.pdf>.
3. Expert Panel on Neurological Imaging, Hagiwara M, Policeni B, et al. ACR appropriateness criteria – sinonasal disease: 2021 update. *J Am Coll Radiol*. 2022 May;19(5S):S175–S193. doi: 10.1016/j.jacr.2022.02.011. PMID: 35550800.
4. Chow AW, Benninger MS, Brook I, et al. IDSA clinical practice guideline for acute bacterial rhinosinusitis in children and adults. *Clin Infect Dis*. 2012 Apr;54(8):e72–e112. doi: 10.1093/cid/cir1043. PMID: 22438350.
5. Gregurić T, Prokopakis E, Vlastos I, et al. Imaging in chronic rhinosinusitis: A systematic review of MRI and CT diagnostic accuracy and reliability in severity staging. *J Neuroradiol*. 2021 Jun;48(4):277–281. doi: 10.1016/j.neurad.2021.01.010. PMID: 33539844.
6. Kim SL, Rank MA, Peters AT. The chronic rhinosinusitis practice parameter. *Ann Allergy Asthma Immunol*. 2023 Sep;131(3):307–310. doi: 10.1016/j.anai.2022.12.022. PMID: 37667905.
7. Krafft RM. Trigeminal neuralgia. *Am Fam Physician*. 2008 May 1;77(9):1291–6. PMID: 18540495.
8. Expert Panel on Neurologic Imaging, Ledbetter LN, Burns J, et al. ACR appropriateness criteria – cerebrovascular diseases, aneurysm, vascular malformation, and subarachnoid hemorrhage. *J Am Coll Radiol*. 2021 Nov;18(11S):S283–S304. doi: 10.1016/j.jacr.2021.08.012. PMID: 34794589.
9. Expert Panel on Neurological Imaging, Rath TJ, Policeni B, et al. ACR appropriateness criteria – cranial neuropathy. Updated 2022. Accessed August 6, 2024. <https://acsearch.acr.org/docs/69509/Narrative/>.
10. Davenport MS, Perazella MA, Yee J, et al. Use of intravenous iodinated contrast media in patients with kidney disease: Consensus statements from the American College of Radiology and the National Kidney Foundation. *Radiology*. 2020;294(3):660–668. doi: 10.1148/radiol.2019192094. PMID: 33015613; PMCID: PMC7525144.
11. The Image Gently Alliance. Procedures – image gentle and CT scans.

Updated 2014. Accessed August 17, 2024.

<https://www.imagegently.org/Procedures/Computed-Tomography>.

12. National Cancer Institute. Radiation risks and pediatric computed tomography (CT): A guide for health care. Updated September 4, 2018. Accessed August 16, 2024.
<https://www.cancer.gov/about-cancer/causes-prevention/risk/radiation/pediatric-ct-scans>.
13. Bedernik A, Wuest W, May MS, et al. Image quality comparison of single-energy and dual-energy computed tomography for head and neck patients: A prospective randomized study. *Eur Radiol*. 2022 Nov;32(11):7700-7709. doi: 10.1007/s00330-022-08689-4. PMID: 35441839; PMCID: PMC9668949.
14. Smith-Bindman R, Chu P, Wang Y, et al. Comparison of the effectiveness of single-component and multicomponent interventions for reducing radiation doses in patients undergoing computed tomography: A randomized clinical trial. *JAMA Intern Med*. 2020 May 1;180(5):666-675. doi: 10.1001/jamainternmed.2020.0064. PMID: 32227142; PMCID: PMC7105953.
15. Aulino JM, Wulff-Burchfield EM, Dietrich MS, et al. Evaluation of CT changes in the head and neck after cancer treatment: Development of a measurement tool. *Lymphat Res Biol*. 2018 Feb;16(1):69-74. doi: 10.1089/lrb.2017.0024. PMID: 29432066; PMCID: PMC5810430.

Clinical Guideline Revision History/Information

Original Date: May 6, 2022		
Review History		
Version 2	8/20/2024	Annual review and policy restructure.