



Cohere Medical Policy – Magnetic Resonance Imaging (MRI), Cardiac

Clinical Guidelines for Medical Necessity Review

Version: 4
Effective Date: October 30, 2024

Important Notices

Notices & Disclaimers:

GUIDELINES ARE SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION-MAKING OF END USERS.

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine the medical necessity of services (the "**Guidelines**") for informational purposes only, and solely for use by Cohere's authorized "**End Users**". These Guidelines (and any attachments or linked third-party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision-making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

©2024 Cohere Health, Inc. All Rights Reserved.

Other Notices:

HCPCS® and CPT® copyright 2024 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

Guideline Information:

Specialty Area: Diagnostic Imaging

Guideline Name: Cohere Medical Policy - Magnetic Resonance Imaging (MRI) - Cardiac

Date of last literature review: 7/24/2024

Document last updated: 10/30/2024

Type: ☒ Adult (18+ yo) | ☒ Pediatric (0-17 yo)

Table of Contents

Important Notices	2
Table of Contents	3
Medical Necessity Criteria	4
Service: Magnetic Resonance Imaging (MRI), Cardiac	4
Recommended Clinical Approach	4
Medical Necessity Criteria	4
Indications	4
Non-Indications	5
Level of Care Criteria	6
Procedure Codes (CPT/HCPCS)	6
Medical Evidence	8
References	8
Clinical Guideline Revision History/Information	12

Medical Necessity Criteria

Service: Magnetic Resonance Imaging (MRI), Cardiac

Recommended Clinical Approach

Cardiac magnetic resonance imaging (MRI) offers exquisite anatomic detail and can also provide valuable functional information through various sequences. While not a first-line imaging modality, it proves highly useful when structural abnormalities (congenital or acquired) or functional deficiencies require further investigation of the heart and pericardial structures.

Medical Necessity Criteria

Indications

→ **Cardiac magnetic resonance imaging (MRI)** is considered appropriate if **ALL** of the following are **TRUE**:

- ◆ First-line cardiac imaging modality such as transthoracic echo is inconclusive/non-diagnostic, and further imaging is indicated for diagnostic and therapeutic purposes; **AND**
- ◆ **ANY** of the following is **TRUE**:
 - Neoplastic conditions (including masses or mass-like conditions) such as cardiac or para-cardiac mass¹; **OR**
 - Infection or an infectious disorder, including suspected infective endocarditis, myocarditis, or complications not diagnosable by other imaging modalities²; **OR**
 - Cardiovascular conditions, known or suspected, including **ANY** of the following:
 - Chest pain of suspected cardiac origin³⁻⁴; **OR**
 - Dyspnea (or other ischemic equivalent) of suspected cardiac origin⁵⁻⁶; **OR**
 - Suspected pulmonary hypertension⁷; **OR**
 - Presyncope or syncope of suspected cardiac origin⁸; **OR**
 - Intermediate to high probability of coronary artery disease⁹⁻¹⁰; **OR**
 - Cardiac valve dysfunction, stenosis, or regurgitation; **OR**

- For evaluation of **ANY** of the following uncategorized/miscellaneous symptoms when applicable:
 - Suspected hypertrophic cardiomyopathy¹¹; **OR**
 - Suspected infiltrative cardiomyopathy (e.g., amyloidosis, hemochromatosis)¹¹; **OR**
 - Suspected or confirmed new-onset heart failure^{4,12}; **OR**
- Preoperative or pre-treatment evaluation (e.g., myocardial viability assessment, before cardiac ablation procedures for arrhythmias, etc); **OR**
- Known or suspected congenital heart disease¹³; **OR**
- Repeat imaging (defined as repeat request following recent imaging of the same anatomic region with the same modality), in the absence of established guidelines, will be considered reasonable and necessary if **ANY** of the following is **TRUE**:
 - New or worsening symptoms, such that repeat imaging would influence treatment; **OR**
 - One-time clarifying follow-up of a prior indeterminate finding; **OR**
 - In the absence of change in symptoms, there is an established need for monitoring which would influence management.

Non-Indications

→ **Cardiac magnetic resonance imaging (MRI)** is not considered appropriate if **ANY** of the following is **TRUE**:

- ◆ The patient has undergone advanced imaging of the same body part within 3 months without undergoing treatment or developing new or worsening symptoms¹⁴; **OR**
- ◆ If contrast is used, history of anaphylactic allergic reaction to gadolinium contrast media with detailed guidelines for use in patients with renal insufficiency; **OR**
- ◆ The patient has metallic clips on vascular aneurysms; **OR**
- ◆ Incompatible implantable devices (e.g., pacemakers, defibrillators, cardiac valves); **OR**
- ◆ Metallic foreign body in orbits/other critical area(s) or within the field of view and obscuring area of concern.

*NOTE: MRI in patients with claustrophobia should be requested at the discretion of the ordering provider.

**NOTE: MRI in pregnant patients should be requested at the discretion of the ordering provider and obstetric care provider

Level of Care Criteria

Inpatient or Outpatient

Procedure Codes (CPT/HCPCS)

CPT/HCPCS Code	Code Description
75557	Cardiac magnetic resonance imaging (MRI) without contrast material, for evaluation of morphology and function
75559	Cardiac magnetic resonance imaging (MRI) with stress imaging, without contrast material, for evaluation of morphology and function
75561	Cardiac magnetic resonance imaging (MRI) without contrast material, followed by contrast material and further sequences, for evaluation of morphology and function
75563	Cardiac magnetic resonance imaging (MRI) with stress imaging, without contrast material, followed by contrast material and further sequences, for evaluation of morphology and function
75565	Cardiac magnetic resonance imaging (MRI) for velocity flow mapping (List separately in addition to code for primary procedure)
C9762	Cardiac magnetic resonance imaging (MRI) for morphology and function, quantification of segmental dysfunction; with strain imaging
C9763	Cardiac magnetic resonance imaging (MRI) for morphology and function, quantification of segmental dysfunction; with stress imaging
S8042	Magnetic resonance imaging (MRI), low-field

Medical Evidence

Miller et al. (2023) CMR-IMPACT (Cardiac Magnetic Resonance Imaging Strategy for the Management of Patients with Acute Chest Pain and Detectable to Elevated Troponin) trial conducted from September 2013 to July 2018 at four U.S. tertiary care hospitals. The trial involved the management of patients with acute chest pain and detectable elevated troponin levels. The 312 participants were randomized into two care pathways: invasive-based (156 participants) and CMR-based (156 participants), with adjustments permitted based on the patient's condition. The primary outcome measured was a composite of death, myocardial infarction, and cardiac-related hospital readmissions or emergency visits. The study followed 312 participants (mean age 60.6 years, 59.9% women) over a median of 2.6 years. The authors conclude no significant difference between clinical and safety outcomes. Benefits include reducing the long-term utilization of invasive angiography, positive discharge outcomes, and enhanced therapeutic yield of angiography. (Clinicaltrials.gov Identifier NCT01931852).¹⁵

Alabed et al. (2020) performed a meta-analysis concerning patient mortality due to pulmonary arterial hypertension (PAH). A total of 1938 patients in 22 studies were included. Research indicates that CMR-derived metrics for right ventricular (RV) volume and function, rather than left ventricular (LV) measurements, predict clinical deterioration. This insight is pertinent for regulatory authorities seeking clinically relevant trial endpoints. Further, this meta-analysis reaffirms the prognostic significance of CMR metrics across a large patient cohort, enabling assessment of how changes in these metrics relate to clinical outcomes such as worsening health and mortality. The authors reaffirm CMR as a useful prognostic marker in PAH among a large cohort. The study confirms that RV function, RV, and left ventricular volumes predict mortality and clinical deterioration in PAH. The study underscores the rationale for using CMR as a meaningful clinical endpoint in trials testing PAH therapies.¹⁶

Kwong et al. (2019) conducted a retrospective study to evaluate the diagnostic and prognostic value. The study enrolled 2349 patients with chest pain syndrome at 13 centers in 11 states. The median follow-up was 5.4 years. A stress CMR showing no ischemia or LGE were linked to a very low incidence of adverse cardiac events in patients with stable intermediate-risk chest pain syndromes. Subsequent cardiac testing also reduced. (Clinicaltrials.gov Identifier NCT03192891).¹⁷

References

1. American College of Radiology (ACR), North American Society for Cardiovascular Imaging (NASCI), Society for Pediatric Radiology (SPR). ACR–NASCI–SPR practice parameter for the performance and interpretation of cardiac magnetic resonance imaging (MRI) (resolution 42). Published 2021. Accessed June 27, 2024. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-Cardiac.pdf>.
2. Expert Panel on Cardiac Imaging, Malik SB, Hsu JY, et al. ACR appropriateness criteria – infective endocarditis. *J Am Coll Radiol*. 2021 May;18(5S):S52–S61. doi: 10.1016/j.jacr.2021.01.010. PMID: 33958118.
3. Expert Panel on Cardiac Imaging, Batlle JC, Kirsch J, et al. ACR appropriateness criteria – chest pain, possible acute coronary syndrome. *J Am Coll Radiol*. 2020 May;17(5S):S55–S69. doi: 10.1016/j.jacr.2020.01.027. PMID: 32370978.
4. Doherty JU, Kort, Mehran R, Schoenhagen P, Soman P. 2019 appropriate use criteria for multimodality imaging in nonvalvular heart disease. Published January 7, 2019. Accessed December 4, 2023. <https://www.acc.org/latest-in-cardiology/ten-points-to-remember/2019/01/04/18/00/2019-auc-for-multimodality-imaging-in-nonvalvular-heart-disease>.
5. Writing Group Members, Doherty JU, Kort S, et al. ACC/AATS/AHA/ASE/ASNC/HRS/SCAI/SCCT/SCMR/STS 2019 appropriate use criteria for multimodality imaging in the assessment of cardiac structure and function in nonvalvular heart disease: A report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and the Society of Thoracic Surgeons. *J Am Soc Echocardiogr*. 2019 May;32(5):553–579. doi: 10.1016/j.echo.2019.01.008. PMID: 30744922.
6. Expert Panel on Cardiac Imaging, Bolen MA, Saeedan MNB, et al. ACR appropriateness criteria – dyspnea-suspected cardiac origin (ischemia already excluded): 2021 update. *J Am Coll Radiol*. 2022 May;19(5S):S37–S52. doi: 10.1016/j.jacr.2022.02.014. PMID: 35550804.
7. Expert Panel on Thoracic Imaging, Sirajuddin A, Mirmomen SM, et al. ACR

- appropriateness criteria – suspected pulmonary hypertension: 2022 update. *J Am Coll Radiol*. 2022 Nov;19(11S):S502–S512. doi: 10.1016/j.jacr.2022.09.018. PMID: 36436973.
8. Expert Panels on Cardiac Imaging and Neurological Imaging, Kligerman SJ, Bykowski J, et al. ACR appropriateness criteria – syncope. *J Am Coll Radiol*. 2021 May;18(5S):S229–S238. doi: 10.1016/j.jacr.2021.02.021. PMID: 33958116.
 9. Expert Panel on Cardiac Imaging, Shah AB, Kirsch J, et al. ACR appropriateness criteria – chronic chest pain, noncardiac etiology unlikely (low to intermediate probability of coronary artery disease). *J Am Coll Radiol*. 2018 Nov;15(11S):S283–S290. doi: 10.1016/j.jacr.2018.09.021. PMID: 30392597.
 10. Expert Panel on Cardiac Imaging, Litmanovich D, Hurwitz LM, et al. ACR appropriateness criteria – chronic chest pain, high probability of coronary artery disease: 2021 update. *J Am Coll Radiol*. 2022 May;19(5S):S1–S18. doi: 10.1016/j.jacr.2022.02.021. PMID: 35550795.
 11. Expert Panel on Cardiac Imaging, Rajiah P, Kirsch J, et al. ACR appropriateness criteria – nonischemic myocardial disease with clinical manifestations (ischemic cardiomyopathy already excluded). *J Am Coll Radiol*. 2021 May;18(5S):S83–S105. doi: 10.1016/j.jacr.2021.01.019. PMID: 33651982.
 12. Expert Panel on Cardiac Imaging, White RD, Kirsch J, et al. ACR appropriateness criteria – suspected new, onset and known nonacute heart failure. *J Am Coll Radiol*. 2018 Nov;15(11S):S418–S431. doi: 10.1016/j.jacr.2018.09.031. PMID: 30392610.
 13. Expert Panels on Cardiac Imaging and Pediatric Imaging, Krishnamurthy R, Suman G, et al. ACR appropriateness criteria – congenital or acquired heart disease. *J Am Coll Radiol*. 2023 Nov;20(11S):S351–S381. doi: 10.1016/j.jacr.2023.08.018. PMID: 38040460.
 14. Wasser EJ, Prevedello LM, Sodickson A, Mar W, Khorasani R. Impact of a real-time computerized duplicate alert system on the utilization of computed tomography. *JAMA Intern Med*. 2013;173(11):1024–1026. doi: 10.1001/jamainternmed.2013.543. PMID: 23609029.
 15. Miller CD, Mahler SA, Snively AC, et al. Cardiac magnetic resonance imaging versus invasive-based strategies in patients with chest pain and detectable to mildly elevated serum troponin: A randomized clinical trial. *Circ Cardiovasc Imaging*. 2023 Jun;16(6):e015063. doi: 10.1161/CIRCIMAGING.122.015063. PMID: 37339173; PMCID: PMC10287041.
 16. Alabed S, Shahin Y, Garg P, et al. Cardiac-MRI predicts clinical worsening

and mortality in pulmonary arterial hypertension: A systematic review and meta-analysis. *JACC Cardiovasc Imaging*. 2021 May;14(5):931-942. doi: 10.1016/j.jcmg.2020.08.013. PMID: 33008758; PMCID: PMC7525356.

17. Kwong RY, Ge Y, Steel K, et al. Cardiac magnetic resonance stress perfusion imaging for evaluation of patients with chest pain. *J Am Coll Cardiol*. 2019 Oct 8;74(14):1741-1755. doi: 10.1016/j.jacc.2019.07.074. PMID: 31582133; PMCID: PMC8109181.

Clinical Guideline Revision History/Information

Original Date: April 8, 2022		
Review History		
Version 2	12/1/2023	
Version 3	8/13/2024	Annual review and policy restructure
Version 4	10/30/2024	Edited repeated imaging criteria language.