



Cohere Medical Policy (Tennessee Medicaid) – Magnetic Resonance Imaging (MRI), Spine (Cervical, Thoracic, and Lumbar)

Clinical Guidelines for Medical Necessity Review

Version: 1
Effective Date: November 21, 2024

Important Notices

Notices & Disclaimers:

GUIDELINES ARE SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION-MAKING OF END USERS.

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine the medical necessity of services (the "**Guidelines**") for informational purposes only, and solely for use by Cohere's authorized "**End Users**". These Guidelines (and any attachments or linked third-party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision-making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

©2024 Cohere Health, Inc. All Rights Reserved.

Other Notices:

HCPCS® and CPT® copyright 2024 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

Guideline Information:

Specialty Area: Diagnostic Imaging

Guideline Name: Cohere Medical Policy (Tennessee Medicaid) - Magnetic Resonance Imaging (MRI), Spine (Cervical, Thoracic, and Lumbar)

Date of last literature review: 8/9/2024

Document last updated: 11/20/2024

Type: ☒ Adult (18+ yo) | ☒ Pediatric (0-17yo)

Table of Contents

Important Notices	2
Table of Contents	3
Medical Necessity Criteria	4
Service: Magnetic Resonance Imaging (MRI), Spine (Cervical, Thoracic, and Lumbar)	4
Recommended Clinical Approach	4
Medical Necessity Criteria	4
Indications	4
Non-Indications	8
Level of Care Criteria	9
Procedure Codes (CPT/HCPCS)	9
Medical Evidence	11
References	12
Clinical Guideline Revision History/Information	14

Medical Necessity Criteria

Service: Magnetic Resonance Imaging (MRI), Spine (Cervical, Thoracic, and Lumbar)

Recommended Clinical Approach

Magnetic resonance imaging (MRI) is a versatile imaging technique that operates on the interaction between radiofrequency electromagnetic fields and specific nuclei in the body, typically hydrogen nuclei, following exposure to a powerful magnetic field. This method allows for the discrimination between normal and abnormal tissues, offering a highly sensitive diagnostic tool for detecting diseases. The effectiveness of MRI stems from the notable contrast inherent in various tissues, both healthy and diseased, owing to differences in their magnetic relaxation properties. MRI of the spine is the preferred imaging modality for pain, radiculopathy, or neurological symptoms. This includes clinical suspicion of cancer, infection, autoimmune disease, persistent symptoms following six weeks of conservative management, or new or worsening symptoms with a history of spine surgery.¹

Contrast should be used at the discretion of the ordering clinician, with guidance from the radiologist as needed. Common indications for administering contrast for an MRI of the spine include infection, prior spine surgery, demyelinating diseases, or tumor. A detailed patient history and indication will ensure the appropriate region is covered when ordering an MRI of the spine. For many patients with neck or back pain, symptoms resolve after a trial of conservative treatment without performing imaging, especially patients with low back pain.

Medical Necessity Criteria

Indications

- **Magnetic resonance imaging (MRI), spine (cervical/thoracic/lumbar)** is considered appropriate if **ANY** of the following is **TRUE**:
- ◆ Neoplastic conditions (including masses or mass-like conditions) including **ANY** of the following¹⁻³:
 - Bone tumors; **OR**

- Intradural-extramedullary masses, including leptomeningeal disease; **OR**
 - Intramedullary masses; **OR**
 - Other extradural soft-tissue neoplasms of **ANY** of the following:
 - Connective tissues; **OR**
 - Muscles; **OR**
 - Regional nerves; **OR**
 - Initial diagnosis of suspected tumor or malignancy as indicated by **ANY** of the following:
 - Abnormal laboratory values; **OR**
 - Inconclusive or abnormal prior imaging; **OR**
 - Primary or metastatic lesion to the spinal cord (including the spinal canal or vertebral bodies) for **ANY** of the following scenarios:
 - Known tumor or malignancy with worsening symptoms or pain; **OR**
 - To monitor response to treatment; **OR**
- ◆ Concern for infection or an infectious disorder in the spine, with **ANY** of the following²⁻⁵:
- **ANY** of the following¹:
 - Discitis; **OR**
 - Epidural abscess; **OR**
 - Postoperative infections; **OR**
 - Surrounding soft-tissue infection; **OR**
 - Vertebral osteomyelitis; **OR**
 - Spinal cord infection and inflammation, including abscess¹; **OR**
 - Pain (including back and neck) with elevated laboratory markers suspicious of infection; **OR**
 - Follow-up to abnormal initial imaging with suspected infection; **OR**
 - Trauma-related conditions, and **ANY** of the following^{1,3,5-6}:
 - Follow-up to initial imaging (e.g., radiograph, CT) with positive findings; **OR**
 - High suspicion for **ANY** of the following injury types:
 - ◆ Fracture; **OR**
 - ◆ Ligamentous; **OR**

- ◆ Nerve; **OR**
- ◆ Spine; **OR**
- Neurological deficit (myelopathy) or radiculopathy, following traumatic event including accident, surgery, or intervention or with **ANY** of the following^{3,7}:
 - Bladder dysfunction; **OR**
 - Bowel dysfunction; **OR**
 - Fecal incontinence; **OR**
 - Loss of anal sphincter tone; **OR**
 - Physical exam finding of major muscle weakness; **OR**
 - Saddle anesthesia; **OR**
 - Severe sciatic/dermatomal sensory loss; **OR**
 - Urinary retention or overflow incontinence; **OR**
 - Weakness (bilateral or progressive) in the lower extremities; **OR**
- Persistent or worsening pain without acute findings on initial imaging (including patients who are elderly, osteoporotic, or have chronic steroid use); **OR**
- ◆ Vascular conditions, known or suspected, including **ANY** of the following¹:
 - Extraspinal vascular malformations; **OR**
 - Spinal cord infarction; **OR**
 - Spinal vascular malformations and/or the cause of occult subarachnoid hemorrhage; **OR**
- ◆ Autoimmune, collagen vascular diseases, or inflammatory conditions including **ANY** of the following¹:
 - Connective tissue disorders (systemic lupus erythematosus); **OR**
 - Muscular dystrophies and myopathies; **OR**
 - **ANY** of the following demyelinating diseases:
 - Acute disseminated encephalomyelitis; **OR**
 - Acute inflammatory demyelinating polyradiculopathy (Guillain-Barre syndrome); **OR**
 - Chronic inflammatory demyelinating polyradiculopathy (including relapsing polyneuropathy); **OR**

- Multiple sclerosis (MS) and its variants (cervical or thoracic); **OR**
- Myelin oligodendrocyte glycoprotein antibody-associated disease; **OR**
- Neuromyelitis optica spectrum disorder; **OR**
- ◆ For evaluation of **ANY** of the following uncategorized/miscellaneous symptoms when applicable:
 - Pain or radiculopathy with **ALL** of the following:
 - At least 8 weeks of conservative treatment; **AND**
 - No trauma or known malignancy with **ANY** of the following^{1,3}:
 - ◆ Discordant pain or radiculopathy to radiograph findings; **OR**
 - ◆ Neurological deficit (myelopathy) or severe radiculopathy with **ANY** of the following:
 - Abnormal EMG; **OR**
 - New or worsening symptoms; **OR**
 - Not previously imaged; **OR**
 - Degenerative conditions including **ANY** of the following¹:
 - Degenerative disc disease and its sequelae in the lumbar, thoracic, and cervical spine (including myelopathy); **OR**
 - Symptomatic radiculopathy; **OR**
 - Compression (suspected) in **ANY** of the following⁷:
 - Spinal cord; **OR**
 - Cauda equina; **OR**
 - Myelopathy (acute or progressive); **OR**
 - Nerve root; **OR**
 - **ANY** of the following¹:
 - Amyloid deposition in the spine; **OR**
 - Cerebrospinal fluid (CSF) leak (may include spontaneous intracranial hypotension); **OR**
 - Gout; **OR**
 - Spinal cord herniation⁸; **OR**
- ◆ Preoperative, postoperative, or pre-treatment evaluation for **ANY** of the following:
 - Planning for treatment fields for radiation therapy; **OR**
 - Postradiation changes (e.g., myelopathy); **OR**

- Epidural and subdural fluid collection¹; **OR**
- Follow-up of incidental or concerning findings seen on other imaging examinations¹; **OR**
- Pre-procedure assessment for vertebroplasty and kyphoplasty⁹; **OR**
- Postoperative fluid collections and soft-tissue changes (extradural and intradural)¹; **OR**
- Postoperative with new or worsening neurological symptoms²⁻³; **OR**
- ◆ **ANY** of the following congenital conditions^{1,10}:
 - Chiari malformation; **OR**
 - Scoliosis with **ANY** of the following¹¹:
 - Neurological symptoms; **OR**
 - Requiring preoperative assessment; **OR**
 - Worsening pain not previously imaged; **OR**
 - Syringohydromyelia (syrinx); **OR**
- ◆ Repeat imaging of a specific area or structure using the same imaging modality (in the absence of an existing follow-up guideline) is considered appropriate when **ALL** of the following are **TRUE**:
 - There is documented clinical necessity; **AND**
 - Prior imaging results of the specific area or structure, obtained using the same imaging modality, must be documented and available for comparison; **AND**
 - **ANY** of the following is **TRUE**:
 - A change in clinical status, such as worsening symptoms or the emergence of new symptoms, that may influence the treatment approach; **OR**
 - The requirement for interval reassessment, which may alter the treatment plan; **OR**
 - One-time follow-up of a prior indeterminate finding to assess for interval change; **OR**
 - The need for re-imaging either before or after performing an invasive procedure.

Non-Indications

→ **Magnetic resonance imaging (MRI), spine (cervical/thoracic/lumbar)** is not considered appropriate if **ANY** of the following is **TRUE**:

- ◆ The patient has undergone advanced imaging of the same body

part within 3 months without undergoing treatment or developing new or worsening symptoms; **OR**

- ◆ If contrast is used, history of anaphylactic allergic reaction to gadolinium contrast media with detailed guidelines for use in patients with renal insufficiency; **OR**
- ◆ The patient has metallic clips on vascular aneurysms; **OR**
- ◆ Incompatible implantable devices (e.g., pacemakers, defibrillators, cardiac valves); **OR**
- ◆ Metallic foreign body in orbits/other critical area(s) or within the field of view and obscuring area of concern.

*NOTE: MRI in patients with claustrophobia should be requested at the discretion of the ordering provider.

**NOTE: MRI in pregnant patients should be requested at the discretion of the ordering provider and obstetric care provider.

Level of Care Criteria

Inpatient or Outpatient

Procedure Codes (CPT/HCPCS)

CPT/HCPCS Code	Code Description
72141	Magnetic resonance imaging (MRI) (e.g., proton), spinal canal and contents, cervical; without contrast material
72142	Magnetic resonance imaging (MRI) (e.g., proton), spinal canal and contents; with contrast material(s)
72146	Magnetic resonance imaging (MRI) (e.g., proton), spinal canal and contents, thoracic; without contrast material
72147	Magnetic resonance imaging (MRI) (e.g., proton), spinal canal and contents, thoracic; with contrast material(s)
72148	Magnetic resonance imaging (MRI) (e.g., proton), spinal canal and contents, lumbar; without contrast material
72149	Magnetic resonance imaging (MRI) (e.g., proton), spinal canal and contents, lumbar; with contrast material(s)
72156	Magnetic resonance imaging (MRI) (e.g., proton), spinal canal and contents, without contrast material, followed

	by contrast material(s) and further sequences; cervical
72157	Magnetic resonance imaging (MRI) (e.g., proton), spinal canal and contents, without contrast material, followed by contrast material(s) and further sequences; thoracic
72158	Magnetic resonance imaging (MRI) (e.g., proton), spinal canal and contents, without contrast material, followed by contrast material(s) and further sequences; lumbar

Medical Evidence

Mathieu and Talbotts (2022) review using magnetic resonance imaging (MRI) to assess spinal emergencies. As an adjunct to CT, MRI proves to be most valuable in situations where additional imaging is necessary to assess spinal stability or compromise of neural elements. Vessel wall imaging techniques and MRA may also be utilized in cases of blunt traumatic cerebrovascular injury, mainly when findings from CTA are inconclusive. The American College of Radiology (ACR) has established guidelines outlining the appropriateness of MRI usage in various clinical scenarios related to spinal injuries and emergencies. MRI is considered 'usually appropriate' for patients with confirmed or suspected spinal cord or nerve root injuries.⁶

Suri et al. (2021) report on a randomized control trial (RCT) that investigated the impact of inserting epidemiological benchmarks into lumbar spine imaging reports as part of the Lumbar Imaging with Reporting of Epidemiology (LIRE) trial. The trial analyzed secondary outcomes, focusing on subsequent nonsurgical and surgical procedures involving the thoracolumbosacral spine and sacroiliac joints. The study included 238,886 adult patients from primary care clinics across four integrated healthcare systems in the United States. All participants underwent lumbar diagnostic imaging between 2013 and 2016. Results indicate that including epidemiological benchmarks (the 'LIRE intervention') did not significantly affect the utilization of non-surgical procedures (e.g., lumbosacral epidural steroid injections, facet joint injections, or facet joint radiofrequency ablation). In addition, the intervention did not impact surgical procedures such as decompression surgery, spinal fusion, or other spine surgeries involving the lumbar, sacral, or thoracic spine. The intervention also did not significantly affect any specific spine procedure.⁹

Ghaffari-Rafi et al. (2021) performed a systematic review and meta-analysis on the role of MRI in clinical decision-making in acute spinal cord injury. Obtaining MRI scans significantly influences the clinical management of patients experiencing acute spinal cord injury (SCI) across all presentations. Guidelines support MRI scans in adult patients with acute SCI before surgical intervention, when feasible, to enhance clinical decision-making. Additional

research is needed to establish the utility and efficacy of MRI in various types of SCI further.⁸

References

1. American College of Radiology (ACR), American Society of Neuroradiology (ASNR), Society of Advanced Body Imaging (SABI), Society of Skeletal Radiology (SSR). ACR–ASNR–SABI–SSR practice parameter for the performance of magnetic resonance imaging (MRI) of the adult spine – resolution 6. Updated 2023. Accessed August 28, 2024. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/mr-adult-spine.pdf>.
2. Expert Panel on Neurological Imaging, McDonald MA, Kirsch CFE, et al. ACR appropriateness criteria – cervical neck pain or cervical radiculopathy. *J Am Coll Radiol*. 2019 May;16(5S):S57–S76. doi: 10.1016/j.jacr.2019.02.023. PMID: 31054759.
3. Expert Panel on Neurological Imaging, Hutchins TA, Peckham M, et al. ACR appropriateness criteria – low back pain: 2021 update. *J Am Coll Radiol*. 2021 Nov;18(11S):S361–S379. doi: 10.1016/j.jacr.2021.08.002. PMID: 34794594.
4. Berbari EF, Kanj SS, Kowalski TJ, et al. 2015 Infectious Diseases Society of America (IDSA) clinical practice guidelines for the diagnosis and treatment of native vertebral osteomyelitis in adults. *Clin Infect Dis*. 2015 Sep 15;61(6):e26–46. doi: 10.1093/cid/civ482. PMID: 26229122.
5. Expert Panel on Neurological Imaging and Musculoskeletal Imaging, Beckmann NM, West OC, et al. ACR appropriateness criteria – suspected spine trauma. *J Am Coll Radiol*. 2019 May;16(5S):S264–S285. doi: 10.1016/j.jacr.2019.02.002. PMID: 31054754.
6. Mathieu J, Talbott JF. Magnetic resonance imaging for spine emergencies. *Magn Reson Imaging Clin N Am*. 2022 Aug;30(3):383–407. doi: 10.1016/j.mric.2022.04.004. PMID: 35995469; PMCID: PMC9926664.
7. Expert Panel on Neurological Imaging, Agarwal V, Shah LM, et al. ACR appropriateness criteria – myelopathy: 2021 update. *J Am Coll Radiol*. 2021 May;18(5S):S73–S82. doi: 10.1016/j.jacr.2021.01.020. PMID: 33958120.
8. Ghaffari-Rafi A, Peterson C, Leon-Rojas JE, et al. The role of magnetic resonance imaging to inform clinical decision-making in acute spinal cord injury: A Systematic review and meta-analysis. *J Clin Med*. 2021 Oct 26;10(21):4948. doi: 10.3390/jcm10214948. PMID: 34768468; PMCID: PMC8584859.

9. Suri P, Meier EN, Gold LS, et al. Providing epidemiological data in lumbar spine imaging reports did not affect subsequent utilization of spine procedures: Secondary outcomes from a stepped-wedge randomized controlled trial. *Pain Med*. 2021 Jun 4;22(6):1272-1280. doi: 10.1093/pm/pnab065. PMID: 33595635; PMCID: PMC8185556.
10. Holly LT, Batzdorf U. Chiari malformation and syringomyelia. *J Neurosurg Spine*. 2019 Nov 1;31(5):619-628. doi: 10.3171/2019.7.SPINE181139. PMID: 31675698.
11. Expert Panel on Pediatric Imaging, Jones JY, Saigal G, et al. ACR appropriateness criteria – scoliosis, child. *J Am Coll Radiol*. 2019 May;16(5S):S244-S251. doi: 10.1016/j.jacr.2019.02.018. PMID: 31054751.

Clinical Guideline Revision History/Information

Original Date: November 21, 2024		
Review History		