



## **Low Back Pain**

*Clinical Guidelines for Medical Necessity Review*

**Version:** V7.0  
**Effective Date:** December 29, 2022

## Important Notices

### Notices & Disclaimers:

**GUIDELINES SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION MAKING OF END USERS.**

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine the medical necessity of services (the "**Guidelines**") for informational purposes only and solely for use by Cohere's authorized "**End Users.**" These Guidelines (and any attachments or linked third party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision-making of any kind; their sole purpose and intended use are to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

©2022 Cohere Health, Inc. All Rights Reserved.

### Other Notices:

CPT copyright 2019 American Medical Association. All rights reserved.  
CPT is a registered trademark of the American Medical Association.

### Guideline Information:

**DiseaseSpecialty Area:** Diseases of the musculoskeletal system and connective tissue (M00-M99)

**CarePath Group:** Spine

**CarePath Name:** Low Back Pain (M54)

**Type:**  Adult (18+ yo) |  Pediatric (0-17yo)

**Physician author:** Mandy Armitage, MD (Sports Medicine)

**Peer reviewed by:** Adrian Thomas, MD (Orthopedic Spine Surgeon), Vijay Yanamadala, MD (Orthopedic Spine Surgeon), Traci Granston, MD (Orthopedic Hand Surgeon)

**Literature review current through:** December 29, 2022

**Document last updated:** December 29, 2022

## Table of Contents

<b>Important Notices</b>	<b>2</b>
<b>Care Path Overview</b>	<b>7</b>
Care Path Clinical Discussion	7
Key Information	8
Definitions	8
<b>Care Path Diagnostic Criteria</b>	<b>11</b>
Disease Classification	11
ICD-10 Codes Associated with Classification	11
Presentation and Etiology	13
Causes and Risk Factors	13
Clinical Presentation	13
Typical Physical Exam Findings	13
Typical Diagnostic Findings	13
<b>CarePath Services &amp; Medical Necessity Criteria</b>	<b>14</b>
<b>Conservative Therapy</b>	<b>14</b>
<hr/>	
<b>Service: Physical Therapy</b>	<b>14</b>
General Guidelines	14
Medical Necessity Criteria	14
Indications	14
Non-Indications	15
Site of Service Criteria	15
Procedure Codes (HCPCS/CPT)	15
<b>Service: Cognitive Behavioral Therapy (CBT)</b>	<b>22</b>
General Guidelines	22
Medical Necessity Criteria	22
Indications	22
Non-Indications	22
Site of Service Criteria	22
Procedure Codes (HCPCS/CPT)	22
<b>Advanced Imaging</b>	<b>24</b>
<hr/>	
<b>Service: Magnetic Resonance Imaging (MRI)</b>	<b>24</b>
General Guidelines	24
Medical Necessity Criteria	24
Indications	24
Non-Indications	25
Site of Service Criteria	25
Procedure Codes (HCPCS/CPT)	25

<b>Service: Computed Tomography (CT)</b>	<b>26</b>
General Guidelines	26
Medical Necessity Criteria	26
Indications	26
Non-Indications	27
Site of Service Criteria	27
Procedure Codes (HCPCS/CPT)	27
<b>Service: Single Photon Emission Computed Tomography/Computed Tomography (SPECT/CT)</b>	<b>28</b>
General Guidelines	28
Medical Necessity Criteria	28
Indications	28
Non-Indications	28
Site of Service Criteria	28
Procedure Codes (HCPCS/CPT)	28
<b>Diagnosics</b>	<b>36</b>
Service: Initial Facet Injection/Medial Branch Block	36
General Guidelines	36
Medical Necessity Criteria	36
Indications	36
Non-Indications	36
Site of Service Criteria	37
Procedure Codes (HCPCS/CPT)	37
Non-Surgical Management	39
<b>Service: Epidural Steroid Injections</b>	<b>39</b>
General Guidelines	39
Medical Necessity Criteria	39
Indications	39
Non-Indications	40
Site of Service Criteria	40
Procedure Codes (HCPCS/CPT)	40
<b>Service: Sacroiliac (SI) Joint Injections</b>	<b>42</b>
General Guidelines	42
Medical Necessity Criteria	42
Indications	42
Non-Indications	42
Site of Service Criteria	42
Procedure Codes (HCPCS/CPT)	43
Service: Subsequent Facet Injection/Medial Branch Block	44
General Guidelines	44

Medical Necessity Criteria	44
Indications	44
Non-Indications	44
Site of Service Criteria	45
Procedure Codes (HCPCS/CPT)	45
<b>Service: Initial/Subsequent Radiofrequency Ablation (RFA)</b>	<b>47</b>
General Guidelines	47
Medical Necessity Criteria	47
Indications	47
Site of Service Criteria	47
Procedure Codes (HCPCS/CPT)	47
Service: Facet Cyst Aspiration	49
General Guidelines	49
Medical Necessity Criteria	49
Indications	49
Site of Service Criteria	49
Procedure Codes (HCPCS/CPT)	49
<b>Service: Intradiscal Biacuplasty, Percutaneous Intradiscal Radiofrequency Thermocoagulation (PIRFT), or Intradiscal Electrothermal Therapy (IDET)</b>	<b>50</b>
General Guidelines	50
Medical Necessity Criteria	50
Indications	50
Non-Indications	50
Site of Service Criteria	50
Procedure Codes (HCPCS/CPT)	51
<b>Surgical Management</b>	<b>52</b>
<b>Service: Lumbar Fusion</b>	<b>52</b>
General Guidelines	52
Medical Necessity Criteria	52
Indications	52
Non-Indications	53
Site of Service Criteria	53
Procedure Codes (HCPCS/CPT)	53
<b>Service: Sacroiliac (SI) Joint Fusion</b>	<b>57</b>
General Guidelines	57
Medical Necessity Criteria	57
Indications	57
Non-Indications	57
Site of Service Criteria	57
Procedure Codes (HCPCS/CPT)	57

Surgical Risk Factors	59
Postoperative Care	63
<b>Service: Physical Therapy</b>	<b>63</b>
General Guidelines	63
Medical Necessity Criteria	63
Indications	63
Non-Indications	63
Site of Service Criteria	63
Procedure Codes (HCPCS/CPT)	63
<b>Service: Home Health Care</b>	<b>70</b>
General Guidelines	70
Medical Necessity Criteria	70
Indications	70
Non-Indications	70
Site of Service Criteria	70
Procedure Codes (HCPCS/CPT)	70
<b>Service: Inpatient Rehabilitation</b>	<b>71</b>
General Guidelines	71
Medical Necessity Criteria	71
Indications	71
Non-Indications	71
Site of Service Criteria	71
Procedure Codes (HCPCS/CPT)	71
<b>Service: Skilled Nursing Facility (SNF)</b>	<b>72</b>
General Guidelines	72
Medical Necessity Criteria	72
Indications	72
Non-Indications	72
Site of Service Criteria	72
Procedure Codes (HCPCS/CPT)	72
<b>References</b>	<b>75</b>
<b>Clinical Guideline Revision History/Information</b>	<b>79</b>

# Care Path Overview

## Care Path Clinical Discussion

The North American Spine Society defines **low back pain (LBP)** as “pain of musculoskeletal origin extending from the lowest rib to the gluteal fold that may at times extend as somatic referred pain into the thigh (above the knee).”<sup>1</sup> Nonspecific pain refers to pain in which no specific cause or structure accounts for the symptoms. Frequent signs and symptoms include pain in the low back that may extend into the gluteal area or thigh but typically not distal to the knee. While many patients experience resolution of LBP symptoms within one month, some will go on to experience subacute (6–12 weeks) or chronic low back pain (more than 12 weeks).<sup>1,2</sup> Anatomic drivers of low back pain include the facet joints, intervertebral disc, or the sacroiliac (SI) joint(s).<sup>4</sup> While the term ‘low back pain’ can encompass several diagnoses, here it explicitly excludes diagnoses or complaints involving leg pain below the knee (radiculopathy) and low back pain due to stenosis.

Routine imaging is not appropriate for patients with acute low back pain (fewer than 6 weeks) without red flags.<sup>3</sup> Advanced imaging (commonly magnetic resonance imaging or MRI) should be reserved for patients whose presentation is complicated, prolonged, or raises suspicion for severe pathology.<sup>3</sup> Electrodiagnostic testing may be helpful if imaging and clinical assessment are inconclusive in determining the etiology of nerve pathology or other comorbid conditions. Appropriate non-surgical interventions for subacute or chronic LBP include physical therapy, ‘back school’ (comprehensive, multidisciplinary rehabilitation), or cognitive behavioral therapy. Several non-surgical interventional procedures are available as alternatives to open surgical intervention. Surgical treatment options for chronic nonspecific LBP include artificial disc replacement or fusion of 1 or 2 lumbar levels, but the latter does not have more superior outcomes than intensive rehabilitation and cognitive therapy.<sup>4</sup>

*The information contained herein gives a general overview of the pathway of nonspecific LBP, including presentation, recommended assessments, and treatment options as supported by the medical literature and existing guidelines. It should be noted that data are commonly conflicting and incomplete, and as such, guidelines and recommendations from various work groups also conflict. The care of patients with spinal health problems is complex, and the information herein is meant to support clinical decision making in adult patients. It is not necessarily applicable to every case, as the entire clinical picture (including comorbidities, spinal health history, etc.) should be considered. Case-by-case treatment decisions are encouraged.*

## Key Information

- Acute low back pain is the most common musculoskeletal complaint seen by primary care providers associated with significant morbidity, increased healthcare cost, and missed work.
- 90% of people experience low back pain in their lifetime; 5%–10% will develop chronic back pain; approximately 9.8% will develop sciatic nerve or radicular back pain.
- Advanced imaging is generally not necessary on initial evaluation or in a primary care setting in the absence of red flags. Most patients will improve within one month without therapy of any kind.
- Initial conservative management includes nonopioid analgesia, muscle relaxants, and physical therapy.
- Alternative, evidence-based treatments include exercise therapy, yoga, chiropractic interventions, aquatic exercise therapy, behavioral therapy, and herbal medicine.
- There is no evidence supporting the use of corticosteroids, with an increased risk of adverse events.
- There is insufficient evidence to recommend for or against the use of epidural injections, facet injections, or spinal cord stimulators as second-line non-surgical therapies.

## Definitions


- **Distraction test:** an examination used to detect abnormalities and inflammation of the lumbar vertebrae and sacroiliac (SI) joint. The test begins with a supine patient. The examiner applies a posterolateral force to the anterior superior iliac spines of a supine patient.
- **Compression:** an SI joint provocative test in which the examiner applies a posterior force directed at the lateral side of the ilium of a supine patient.
- **Thigh thrust test:** an SI joint provocative test in which the supine patient flexes their hip and knee. The examiner applies a posterior force directed at the SIJ of the flexed lower extremity while the other hand stabilizes the sacrum.
- **Gaenslen's test:** an examination used to detect abnormalities and inflammation of the lumbar vertebrae and sacroiliac joint. The supine patient flexes their asymptomatic hip with their knee also flexed. The examiner drops the opposite lower extremity off the side of the table and applies a posterior force so that the symptomatic leg hyperextends at the hip. A positive test occurs when the patient's typical pain is reproduced.<sup>5,6</sup>
- **Sacral thrust test:** an SI joint provocative test in which the examiner applies a posterior force directed at the sacral base of the supine patient.

- **FABER (Flexion, Abduction, and External Rotation) Test:** may be used to assess a suspected labral tear. The examination begins with a supine patient. The examiner places the patient in a figure-4 position with their hip flexed and abducted with the lateral ankle resting on the contralateral thigh proximal to the knee. The examiner then applies gentle downward force against the knee of the abducted leg. A positive result occurs when the patient reports groin pain on the side of the lateral leg.<sup>7</sup>

# Low Back Pain

## What is a "Cohere Care Path"?

These Care Paths organize the services typically considered most clinically optimal and likely to be automatically approved. These service recommendations also include the suggested sequencing and quantity or frequency determined clinically appropriate and medically necessary for the management of most patient care scenarios in this Care Path's diagnostic cohort.



### Key

- <sup>PA</sup> = Service may require prior authorization
- ★ = Denotes preferred service
- AND = Services completed concurrently
- OR = Services generally mutually exclusive

- = Non-surgical management prior authorization group of services
- = Surgical management prior authorization group of services
- - - = Subsequent service
- - - = Management path moves to a different management path

# Care Path Diagnostic Criteria

## Disease Classification

Low Back Pain (LBP)

### ICD-10 Codes Associated with Classification

ICD-10 Code	Code Description/Definition
G95.9	Disease of spinal cord, unspecified
M40.205	Unspecified kyphosis, thoracolumbar region
M45.9	Ankylosing spondylitis of unspecified sites in spine
M46.00	Spinal enthesopathy, site unspecified
M46.05	Spinal enthesopathy, thoracolumbar region
M46.06	Spinal enthesopathy, lumbar region
M46.07	Spinal enthesopathy, lumbosacral region
M46.08	Spinal enthesopathy, sacral and sacrococcygeal region
M46.09	Spinal enthesopathy, multiple sites in spine
M46.1	Sacroiliitis, not elsewhere classified
M46.20	Osteomyelitis of vertebra, site unspecified
M46.25	Osteomyelitis of vertebra, thoracolumbar region
M46.26	Osteomyelitis of vertebra, lumbar region
M46.27	Osteomyelitis of vertebra, lumbosacral region
M46.28	Osteomyelitis of vertebra, sacral and sacrococcygeal region
M46.30	Infection of intervertebral disc (pyogenic), site unspecified
M46.35	Infection of intervertebral disc (pyogenic), thoracolumbar region
M46.36	Infection of intervertebral disc (pyogenic), lumbar region
M46.37	Infection of intervertebral disc (pyogenic), lumbosacral region
M46.38	Infection of intervertebral disc (pyogenic), sacral and sacrococcygeal region
M46.39	Infection of intervertebral disc (pyogenic), multiple sites in spine
M46.40	Discitis, unspecified, site unspecified
M46.45	Discitis, unspecified, thoracolumbar region
M46.46	Discitis, unspecified, lumbar region
M46.47	Discitis, unspecified, lumbosacral region

M46.48	Discitis, unspecified, sacral and sacrococcygeal region
M46.49	Discitis, unspecified, multiple sites in spine
M47.814	Spondylosis without myelopathy or radiculopathy, thoracic region
M47.816	Spondylosis without myelopathy or radiculopathy, lumbar region
M47.817	Spondylosis without myelopathy or radiculopathy, lumbosacral region
M48.54XA	Collapsed vertebra, not elsewhere classified, thoracic region, initial encounter for fracture
M51.2	Other thoracic, thoracolumbar and lumbosacral intervertebral disc displacement
M51.24	Other intervertebral disc displacement, thoracic region
M51.25	Other intervertebral disc displacement, thoracolumbar region
M51.26	Other intervertebral disc displacement, lumbar region
M51.27	Other intervertebral disc displacement, lumbosacral region
M53.3	Sacrococcygeal disorders, not elsewhere classified
M53.84	Other specified dorsopathies, thoracic region
M53.85	Other specified dorsopathies, thoracolumbar region
M53.86	Other specified dorsopathies, lumbar region
M54.14	Radiculopathy, thoracic region
M54.40	Lumbago with sciatica, unspecified side
M54.41	Lumbago with sciatica, right side
M54.42	Lumbago with sciatica, left side
M54.5	Low back pain
M54.6	Pain in thoracic spine
M54.89	Other dorsalgia
M54.9	Dorsalgia, unspecified
M80.08XA	Age-related osteoporosis with current pathological fracture, vertebra(e), initial encounter for fracture
M81.0	Age-related osteoporosis without current pathological fracture
M99.04	Segmental and somatic dysfunction of sacral region
S22.000A	Wedge compression fracture of unspecified thoracic vertebra, initial encounter for closed fracture
S22.060A	Wedge compression fracture of T7-T8 vertebra, initial encounter for closed fracture
S22.080A	Wedge compression fracture of T11-T12 vertebra, initial encounter

	for closed fracture
S22.080D	Wedge compression fracture of T11-T12 vertebra, subsequent encounter for fracture with routine healing
S32.000A	Wedge compression fracture of unspecified lumbar vertebra, initial encounter for closed fracture
S32.009K	Unspecified fracture of unspecified lumbar vertebra, subsequent encounter for fracture with nonunion
S32.010A	Wedge compression fracture of first lumbar vertebra, initial encounter for closed fracture
S32.010D	Wedge compression fracture of first lumbar vertebra, subsequent encounter for fracture with routine healing
S32.020A	Wedge compression fracture of second lumbar vertebra, initial encounter for closed fracture
S32.030A	Wedge compression fracture of third lumbar vertebra, initial encounter for closed fracture
S32.040A	Wedge compression fracture of fourth lumbar vertebra, initial encounter for closed fracture
S33.5XXD	Sprain of ligaments of lumbar spine, subsequent encounter
S39.012A	Strain of muscle, fascia and tendon of lower back, initial encounter

## **Presentation and Etiology**

### ***Causes and Risk Factors***

The following are risk factors for conversion from acute to chronic pain:

- History of LBP
- Psychosocial factors
- Workplace factors
- Pain severity
- Functional impairment

### ***Clinical Presentation***

The predominant symptom is low back pain without “red flag” symptoms (see advanced imaging section). Lower extremity pain or paresthesias may be present, typically does not travel distal to the knee, and does not predominate as a presenting symptom.

### ***Typical Physical Exam Findings***

The examination is typically nonspecific and without indication for pathophysiological abnormality (i.e., provocative testing such as the straight leg test should be negative).

There is insufficient evidence to recommend for or against using specific findings to determine which anatomic structure is causing pain.

The patient may have positive results to SI joint provocative tests such as distraction, compression, thigh thrust, Gaenslen’s, and sacral thrust.

### ***Typical Diagnostic Findings***

There is insufficient evidence to recommend for or against using assessment tools or questionnaires to help identify the cause of low back pain.

# CarePath Services & Medical Necessity Criteria

## Conservative Therapy

**Service: Physical Therapy**

### General Guidelines

- **Units, Frequency, & Duration:** Referrals commonly suggest 4-6 weeks for most programs.
- **Criteria for Subsequent Requests:** The medical necessity of subsequent physical therapy should be evaluated based on the individual's response to previous sessions (e.g., clinically relevant sustained reductions in pain, improvement in the individual's functional abilities).
- **Recommended Clinical Approach:** The following are suitable treatment options in addition to physical therapy<sup>8</sup>:
  - Mindfulness-based stress reduction (MBSR) programs
  - McKenzie exercise
  - Yoga
  - Spinal manipulative therapy (SMT)
  - Work hardening/conditioning.

The following are recommended to improve pain or functional outcomes for patients with subacute or chronic LBP:

- The combination of physical therapy (PT) and cognitive behavioral therapy (CBT)
- General exercise or stabilization in PT
- Back school (multidisciplinary rehabilitation with body mechanics education, exercise programs, and behavioral therapy)
- Acupuncture
- Aerobic exercise
- **Exclusions:** None.

### Medical Necessity Criteria

#### Indications

→ **Physical therapy** for education, spinal manipulation therapy, or acupuncture is considered appropriate if **ALL** of the following are **TRUE**<sup>8-9</sup>:

- Acute low back pain for less than 6 weeks
- Chronic low back pain for greater than 6 weeks

## Non-Indications

→ **Physical therapy** may not be considered appropriate if **ANY** of the following is **TRUE**:

- ◆ Progressive neurological deficits
- ◆ Severely unsteady gait related to myelopathy/balance or generalized lower extremity weakness
- ◆ Hyperreflexia
- ◆ Positive Babinski or clonus
- ◆ Bowel or bladder incontinence
- ◆ Saddle anesthesia

## Site of Service Criteria

Outpatient

## Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
97010	Application of hot or cold packs
97012	Application of mechanical traction
97014	Application of electrical stimulation
97016	Application of vasopneumatic devices
97018	Application of paraffin bath
97022	Application of whirlpool
97024	Application of diathermy
97026	Application of infrared modality
97028	Application of ultraviolet modality
97032	Application of manual electrical stimulation
97033	Application of iontophoresis
97034	Application of contrast baths
97035	Application of ultrasound modality
97036	Application of Hubbard tank

97039	Modality service
97110*	Therapeutic exercises to develop strength and endurance, range of motion and flexibility
97112	Neuromuscular reeducation of movement, balance, coordination, kinesthetic sense, posture, and proprioception for sitting and standing activities
97113	Aquatic therapy with therapeutic exercises
97116	Gait training including stair climbing
97124	Massage including effleurage and petrissage; Massage including effleurage and tapotement; Massage including effleurage, petrissage and tapotement; Massage including petrissage and tapotement
97139	Therapeutic procedure
97140	Manual therapy techniques
97150	Group therapeutic procedures
97164	Physical therapy re-evaluation of established plan of care, high complexity, typical time with patient 20 minutes; Physical therapy re-evaluation of established plan of care, high complexity, typical time with patient and family 20 minutes; Physical therapy re-evaluation of established plan of care, high complexity, typical time with patient's family 20 minutes
97530	Direct therapeutic activities with use of dynamic activities to improve functional performance, each 15 minutes
97535	Home management training, direct one-on-one contact, each 15 minutes; Self-care management training, direct one-on-one contact, each 15 minutes
97537	Community reintegration training, direct one-on-one contact, each 15 minutes; Work reintegration training, direct one-on-one contact, each 15 minutes
97542	Wheelchair management, each 15 minutes

97545	Work conditioning, initial 2 hours; Work hardening, initial 2 hours
97546	Work conditioning, each additional hour; Work hardening, each additional hour
97750	Physical performance measurement with written report, each 15 minutes; Physical performance test with written report, each 15 minutes
97755	Assistive technology assessment with written report, direct one-on-one contact, each 15 minutes
97760	Initial orthotic management and training with assessment and fitting of lower extremities and trunk, each 15 minutes; Initial orthotic management and training with assessment and fitting of lower extremities, each 15 minutes; Initial orthotic management and training with assessment and fitting of lower extremity and trunk, each 15 minutes; Initial orthotic management and training with assessment and fitting of lower extremity, each 15 minutes; Initial orthotic management and training with assessment and fitting of trunk, each 15 minutes; Initial orthotic management and training with assessment and fitting of upper and lower extremities and trunk, each 15 minutes
97761	Initial prosthetic training of lower extremities, each 15 minutes; Initial prosthetic training of lower extremity, each 15 minutes Initial prosthetic training of upper and lower extremities, each 15 minutes; Initial prosthetic training of upper extremities, each 15 minutes; Initial prosthetic training of upper extremity, each 15 minutes
97763	Subsequent orthotic management and training of lower extremities and trunk, each 15 minutes Subsequent orthotic management and training of lower extremity and trunk, each 15 minutes Subsequent orthotic management and training of lower extremity, each 15 minutes Subsequent orthotic management and training of upper

	<p>and lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management and training of upper extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management and training of upper extremities, each 15 minutes</p> <p>Subsequent orthotic management and training of upper extremity and trunk, each 15 minutes</p> <p>Subsequent orthotic management and training of upper extremity, each 15 minutes</p> <p>Subsequent orthotic management of lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management of lower extremity and trunk, each 15 minutes</p> <p>Subsequent orthotic management of lower extremity, each 15 minutes</p> <p>Subsequent orthotic management of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management of upper extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management of upper extremities, each 15 minutes</p> <p>Subsequent orthotic management of upper extremity and trunk, each 15 minutes</p> <p>Subsequent orthotic management of upper extremity, each 15 minutes</p> <p>Subsequent orthotic training of lower extremity, each 15 minutes</p> <p>Subsequent orthotic training of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic training of upper extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic training of upper extremities, each 15 minutes</p> <p>Subsequent orthotic training of upper extremity and trunk, each 15 minutes</p> <p>Subsequent orthotic training of upper extremity, each 15 minutes</p> <p>Subsequent prosthetic management and training of lower extremities and trunk, each 15 minutes</p>
--	--

	<p>Subsequent prosthetic management and training of lower extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of lower extremity, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper extremities, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper extremity, each 15 minutes</p> <p>Subsequent prosthetic management of lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of lower extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of lower extremity, each 15 minutes</p> <p>Subsequent prosthetic management of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of upper extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of upper extremities, each 15 minutes</p> <p>Subsequent prosthetic management of upper extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of upper extremity, each 15 minutes</p> <p>Subsequent prosthetic training of lower extremity, each 15 minutes</p> <p>Subsequent prosthetic training of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic training of upper extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic training of upper extremities, each 15 minutes</p> <p>Subsequent prosthetic training of upper extremity and</p>
--	---

	<p>trunk, each 15 minutes</p> <p>Subsequent prosthetic training of upper extremity, each 15 minutes</p> <p>Subsequent orthotic management and training of lower extremities, each 15 minutes</p> <p>Subsequent orthotic management of lower extremities, each 15 minutes</p> <p>Subsequent orthotic training of lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic training of lower extremities, each 15 minutes</p> <p>Subsequent orthotic training of lower extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of lower extremities, each 15 minutes</p> <p>Subsequent prosthetic management of lower extremities, each 15 minutes</p> <p>Subsequent prosthetic training of lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic training of lower extremities, each 15 minutes</p> <p>Subsequent prosthetic training of lower extremity and trunk, each 15 minutes</p>
97799	Unlisted physical medicine/rehabilitation service or procedure
420	Physical Therapy
421	Physical Therapy: Visit Charge
422	Physical Therapy: Hourly Charge
423	Physical Therapy: Group Rate
424	Physical Therapy: Evaluation/Re-evaluation
429	Physical Therapy: Other Physical Therapy
97163	Evaluation of physical therapy, typically 45 minutes
97161	Evaluation of physical therapy, typically 20 minutes
97162	Evaluation of physical therapy, typically 30 minutes
97168	Re-evaluation of occupational therapy established plan of care, typically 30 minutes

97165	Evaluation of occupational therapy, typically 30 minutes
97166	Evaluation of occupational therapy, typically 45 minutes
97167	Evaluation of occupational therapy established plan of care, typically 60 minutes
G0151	Hhcp-serv of pt,ea 15 min

\*Default codes for suggested services

## **Service: Cognitive Behavioral Therapy (CBT)**

### **General Guidelines**

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach:** Cognitive behavioral therapy (CBT) is required for lumbar fusion for axial back pain where at least 6 months of conservative treatment has failed, with no instability.
- **Exclusions:** None.

### **Medical Necessity Criteria**

#### **Indications**

→ **Cognitive behavioral therapy** is considered appropriate if **ANY** of the following is **TRUE**:

- ◆ The patient has **ANY** positive findings from the [clinical presentation](#) and [typical physical exam findings](#) lists.
- ◆ The patient requires a lumbar fusion.
- ◆ The patient completed more than 6 months of non-surgical treatment.

#### **Non-Indications**

None.

### **Site of Service Criteria**

Outpatient

### **Procedure Codes (HCPCS/CPT)**

<b>HCPCS Code</b>	<b>Code Description/Definition</b>
96125	Standardized cognitive performance testing, per hour by a qualified health care professional
97129	Therapeutic interventions that focus on cognitive function and compensatory strategies to manage the performance of an activity, direct (one-on-one) patient contact, initial 15 minutes
97130	Therapeutic interventions that focus on cognitive function and compensatory strategies to manage the performance of an activity, direct (one-on-one) patient

	contact, each additional 15 minutes
99483	Assessment of and care planning for outpatient with cognitive impairment, typical time 50 minutes
G0515	Cognitive skills development

## Advanced Imaging

### Service: Magnetic Resonance Imaging (MRI)

#### General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach<sup>10</sup>:**
  - MRI without contrast is the preferred method of advanced imaging if radiographs are inconclusive or do not provide enough detail for a definitive diagnosis.
  - MRI with contrast may be appropriate if MRI without contrast was indeterminate or if the patient previously had lumbar surgery.
- **Exclusions:** None.

#### Medical Necessity Criteria

#### Indications

- **MRI** is considered appropriate if **ANY** of the following are **TRUE<sup>10</sup>**:
- ◆ The patient has **ALL** of the following:
    - The patient has **ANY** positive findings from the [clinical presentation](#) and [typical physical exam findings](#) lists.
    - The patient failed to show significant improvement in pain or disability level due to symptoms, despite more than 6 weeks of conservative care (conservative care includes a combination of physical therapy, provider-directed home exercise program, and anti-inflammatory/pain management medications or oral steroids).
    - Radiographs are inconclusive or do not provide enough detail for a definitive diagnosis.
  - ◆ The patient presents with **ANY** of the following “red flags”:
    - Progressive neurological deficits
    - Unsteady gait related to myelopathy/balance or generalized lower extremity weakness
    - Hyperreflexia
    - Positive Babinski or clonus
    - Bowel or bladder incontinence
    - Saddle anesthesia

## Non-Indications

- **MRI** may not be appropriate if **ANY** of the following is **TRUE**<sup>11-12</sup>:
- ◆ Non-compatible implanted devices
  - ◆ Metallic intraocular foreign bodies
  - ◆ Claustrophobia

## Site of Service Criteria

None.

## Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
72148	MRI of lumbar spinal canal and contents; MRI of lumbar spinal canal and contents without contrast
72149	MRI of lumbar spinal canal and contents with contrast
72158	MRI of lumbar spinal canal and contents without contrast, followed by contrast and further sections
72146	MRI scan of middle spinal canal
72157	MRI scan of middle spinal canal before and after contrast
72195	MRI scan of pelvis
72197	MRI scan of pelvis before and after contrast
72196	MRI scan of pelvis with contrast
72130	CT scan of middle spine before and after contrast

## **Service: Computed Tomography (CT)**

### General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach<sup>10</sup>:** A CT recommended if there is a concern for fracture. In the absence of red flag signs/symptoms, advanced imaging may not be required at initial presentation; CTM may be utilized if MRI is contraindicated or indeterminate. CT may be utilized in this patient population if MRI contraindicated or indeterminate. CT may be ordered with an MRI to provide additional visibility to the bony anatomy or to assess previous surgery.
- **Exclusions:** None.

### Medical Necessity Criteria

#### Indications

- **CT/CTM** is considered appropriate if **ANY** of the following is **TRUE<sup>10</sup>**:
- ◆ The patient is being considered for a CTM and **ALL** of the following are **TRUE**:
    - The patient has **ANY** positive findings from the [clinical presentation](#) and [typical physical exam findings](#) lists.
    - The patient has failed to show significant improvement in pain or disability level due to symptoms, despite more than 6 weeks of conservative care (conservative care includes a combination of physical therapy, provider-directed home exercise program, and anti-inflammatory/pain management medications or oral steroids).
    - MRI is contraindicated or indeterminate for reasons such as an artifact from a previous surgery (e.g., anomalies in visual representation that impact imaging quality).
  - ◆ The patient is being considered for a CTM and presents with **ANY** of the following “red flags”:
    - Progressive neurological deficits
    - Unsteady gait related to myelopathy/balance or generalized lower extremity weakness
    - Hyperreflexia
    - Positive Babinski or clonus
    - Bowel or bladder incontinence
    - Saddle anesthesia
  - ◆ The patient is being considered for a CT and **ANY** of the following is **TRUE**:

- The patient meets the criteria for MRI or CTM but cannot receive either service due to contraindications.
- MRI or CTM studies are indeterminate.
- There is a need to obtain additional information that is not provided by an MRI (e.g., details of the bony anatomy or previous surgery).

### Non-Indications

→ **CT/CTM** may not be considered appropriate if **ANY** of the following is **TRUE**:

- ◆ If the patient is being considered for a **CTM** and **ANY** of the following are **TRUE**:
  - In patients with bleeding disorders
  - In patients with an allergy to iodinated contrast agents<sup>13</sup>
  - In patients who are pregnant<sup>14</sup>
- ◆ If the patient is being considered for a CT and **ANY** of the following is **TRUE**<sup>14</sup>:
  - The patient is pregnant.

### Site of Service Criteria

None.

### Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
72131	CT of lumbar spine; CT of lumbar spine without contrast
72132	CT of lumbar spine with contrast;
72133	CT of lumbar spine without contrast, followed by contrast and further sections
72128	CT scan of middle spine
72129	CT scan of middle spine with contrast
72130	CT scan of middle spine before and after contrast
72192	CT scan pelvis
72193	CT scan pelvis with contrast
72194	CT scan of pelvis before and after contrast

## ***Service: Single Photon Emission Computed Tomography/Computed Tomography (SPECT/CT)***

### **General Guidelines**

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach<sup>10</sup>:** SPECT may be appropriate to detect infection or occult fractures of the vertebrae if suspected. There is insufficient evidence to recommend for or against the use of SPECT imaging in the diagnosis of zygapophyseal joint pain.
- **Exclusions:** None.

### **Medical Necessity Criteria**

#### **Indications**

- **SPECT** is considered appropriate if **ALL** of the following are **TRUE<sup>10</sup>**:
- ◆ Zygapophyseal joint pain is suspected

#### **Non-Indications**

None.

### **Site of Service Criteria**

None.

### **Procedure Codes (HCPCS/CPT)**

<b>HCPCS Code</b>	<b>Code Description/Definition</b>
78803	<p>Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in limited area, single day;</p> <p>Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in limited area, with vascular flow and blood pool imaging, single day;</p> <p>Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in limited area, single day;</p> <p>Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in limited area, with vascular flow and blood pool imaging, single day;</p> <p>Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in</p>

	<p>limited area, single day;  Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in limited area, with vascular flow and blood pool imaging, single day;  Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in limited area, single day;  Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in limited area, with vascular flow and blood pool imaging, single day</p>
78830	<p>Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in limited area, with concurrently acquired computed tomography (CT) transmission, single day;  Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in limited area, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day;  Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in limited area, with concurrently acquired computed tomography (CT) transmission, single day;  Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in limited area, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day;  Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in limited area, with concurrently acquired computed tomography (CT) transmission, single day;  Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in limited area, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day;  Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in limited area, with concurrently acquired computed tomography (CT) transmission, single day;  Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in limited area,</p>

	with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day
78831	<p>Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in abdomen and pelvis, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in abdomen and pelvis, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in minimum of 2 areas, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in minimum of 2 areas, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in pelvis and knees, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in pelvis and knees, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in single area, multiple days; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in single area, with vascular flow and blood pool imaging, multiple days; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in abdomen and pelvis, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in abdomen and pelvis, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in minimum of 2 areas, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in minimum of 2 areas, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in pelvis and knees, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in pelvis and knees, with</p>

	<p>vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in single area, multiple days; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in single area, with vascular flow and blood pool imaging, multiple days; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in abdomen and pelvis, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in abdomen and pelvis, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in minimum of 2 areas, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in minimum of 2 areas, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in pelvis and knees, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in pelvis and knees, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in single area, multiple days; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in single area, with vascular flow and blood pool imaging, multiple days; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in abdomen and pelvis, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in abdomen and pelvis, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in minimum of 2 areas, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in minimum of 2 areas, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in pelvis and knees, single day; Single photon emission</p>
--	--

	<p>computed tomography (SPECT) radiopharmaceutical localization of tumor in pelvis and knees, with vascular flow and blood pool imaging, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in single area, multiple days; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in single area, with vascular flow and blood pool imaging, multiple days</p>
78832	<p>Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in abdomen and pelvis with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in abdomen and pelvis, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in minimum of 2 areas, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in minimum of 2 areas, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in pelvis and knees with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in pelvis and knees, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agent in single area, with concurrently acquired computed tomography (CT) transmission, multiple days; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in abdomen and pelvis with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT)</p>

	<p>imaging of distribution of radiopharmaceutical agents in abdomen and pelvis, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in minimum of 2 areas with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in minimum of 2 areas, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in pelvis and knees with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in pelvis and knees, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) imaging of distribution of radiopharmaceutical agents in single area, with concurrently acquired computed tomography (CT) transmission, multiple days; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in abdomen and pelvis with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in abdomen and pelvis, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in minimum of 2 areas with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in minimum of 2 areas, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in pelvis</p>
--	--

	<p>and knees with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in pelvis and knees, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of inflammation in single area, with concurrently acquired computed tomography (CT) transmission, multiple days; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in abdomen and pelvis with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in abdomen and pelvis, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in minimum of 2 areas with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in minimum of 2 areas, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in pelvis and knees in minimum of 2 areas, with vascular flow and blood pool imaging, with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in pelvis and knees with concurrently acquired computed tomography (CT) transmission, single day; Single photon emission computed tomography (SPECT) radiopharmaceutical localization of tumor in single area, with concurrently acquired computed tomography (CT) transmission, multiple days</p>
--	--

## Diagnostics

### ***Service: Initial Facet Injection/Medial Branch Block***

#### General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** For second injections, the previous injection must result in greater than 80% relief of symptoms. If the patient has had greater than or equal to 2 injections, the previous injection must result in 50% improvement of symptoms for 3 months.<sup>15-21</sup>
- **Recommended Clinical Approach:** Blocks to be performed with imaging (fluoroscopy or ultrasound with Doppler) guidance. Medial branch blocks with a double-injection technique and pain improvement threshold of 80% may be used to establish the diagnosis of lumbar facet-mediated pain and predict a favorable response to medial nerve ablation by thermocoagulation. There is no evidence to support the use of diagnostic facet blocks as a predictor of lumbar fusion outcome in patients with chronic low back pain. There is insufficient evidence to make a recommendation for or against the diagnostic utility of zygapophyseal/facet joint injections. Pregnancy is a relative contraindication.
- **Exclusions:** None.

#### Medical Necessity Criteria

##### Indications

- **Initial Facet Injection/Medial Branch Block** may be considered appropriate if **ALL** of the following are **TRUE**:
- ◆ The patient fails to show significant improvement in pain or disability level due to symptoms despite receiving more than 12 weeks of conservative management (conservative management includes a combination of physical therapy, provider-directed home exercise program, and anti-inflammatory/pain management medications or oral steroids).<sup>22</sup>
  - ◆ The patient had advanced imaging that revealed either facet arthritis or cervical/thoracic spondylosis (arthritis) at the level requested.

##### Non-Indications

- **Initial Facet Injection/Medial Branch Block** may not be appropriate if **ANY** of the following is **TRUE**:<sup>2</sup>
- ◆ Systemic infection
  - ◆ Local infection at the procedure site

- ◆ Coagulopathy or recent use of blood-thinning agents
- ◆ Previous fusion at the symptomatic level

### **Site of Service Criteria**

Outpatient

### **Procedure Codes (HCPCS/CPT)**

<b>HCPCS Code</b>	<b>Code Description/Definition</b>
64493	Injection of diagnostic agent into nerve of single lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into nerve of single lumbar paravertebral facet joint using imaging guidance; Injection of diagnostic agent into single lumbar paravertebral facet joint using imaging guidance; Injection of therapeutic agent into nerve of single lumbar paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into single lumbar paravertebral facet joint using computed tomography (CT) guidance
64494	Injection of diagnostic agent into nerve of second lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into nerve of second lumbar paravertebral facet joint using imaging guidance; Injection of diagnostic agent into second lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into second lumbar paravertebral facet joint using imaging guidance; Injection of therapeutic agent into nerve of second lumbar paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into second lumbar paravertebral facet joint using computed tomography (CT) guidance
64495	Injection of diagnostic agent into nerve of third and any additional lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into nerve of third and any additional lumbar paravertebral facet joint using imaging guidance; Injection of diagnostic agent into nerve of third and any additional sacral paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into nerve of third and any additional sacral paravertebral facet joint using imaging guidance; Injection of diagnostic agent into third and any

	additional lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into third and any additional lumbar paravertebral facet joint using imaging guidance; Injection of diagnostic agent into third and any additional sacral paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into third and any additional sacral paravertebral facet joint using imaging guidance; Injection of therapeutic agent into nerve of third and any additional lumbar paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into nerve of third and any additional sacral paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into third and any additional lumbar paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into third and any additional sacral paravertebral facet joint using computed tomography (CT) guidance
0216T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, lumbar or sacral; single level
64999	Unlisted procedure, nervous system
0213T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; single level
0214T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; second level (list separately in addition to code for primary procedure)
0215T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; third and any additional level(s) (list separately in addition to code for primary procedure)

## Non-Surgical Management

### ***Service: Epidural Steroid Injections***

#### General Guidelines

- **Units, Frequency, & Duration:** When the medical necessity criteria are met, a total of 3 epidural steroid injections per episode of pain per region may be performed in 6 months.
- **Criteria for Subsequent Requests:** A second injection may be considered if the patient has had one epidural steroid injection within the last six months. If the patient has had two or more epidural steroid injections in the past six months and the previous injection resulted in a 50% improvement of symptoms for three months, another injection may be considered.<sup>23-39</sup>
- **Recommended Clinical Approach:** Perform epidural steroid injections with radiographic image guidance. One interlaminar injection is recommended at a time. Up to two transforaminal injections may be appropriate at a time bilaterally (at the same nerve level) or unilaterally at adjacent levels.
- **Exclusions:** None.

#### Medical Necessity Criteria

#### Indications

- **Epidural steroid injections (ESI)** are considered appropriate if **ALL** of the following are **TRUE**:
- ◆ The patient has **ANY** positive findings from the [clinical presentation](#) and [typical physical exam findings](#) lists.
  - ◆ Advanced imaging corresponds to clinical presentation and shows nerve impingement.
  - ◆ Patient has failed to show significant improvement in pain or disability level due to symptoms, despite receiving conservative care for more than 6 weeks, or the patient is unable to complete conservative care due to the severity of symptoms. Conservative care is a combination of either physical therapy or provider-directed home exercise program **AND** medications, anti-inflammatories, or oral steroids.
  - ◆ The patient is outside the acute period (greater than 4 weeks).
  - ◆ The patient has had fewer than 3 epidural steroid injections per region within the past 6 months with at least 3 weeks in between injections, at which time the patient had 50% improvement of symptoms.<sup>34</sup>

## Non-Indications

→ **Epidural steroid injections** are non indicated if **ANY** of the following is **TRUE**<sup>37</sup>:

- ◆ Hypersensitivity (or allergy) to steroids.
- ◆ Local or systemic infection.
- ◆ Coagulopathy or recent use of blood-thinning agents.
- ◆ Uncontrolled diabetes.

## Site of Service Criteria

Outpatient

## Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
62322	Injection of substance into lumbar spinal canal; Insertion of catheter and injection of substance into lumbar interlaminar epidural space;
62323	Injection of substance into lumbar spinal canal using imaging guidance; Insertion of catheter and injection of substance into lumbar interlaminar epidural space using imaging guidance; Insertion of catheter and injection of substance into lumbar interlaminar subarachnoid space using imaging guidance; Insertion of needle and injection of substance into lumbar interlaminar epidural space using imaging guidance; Insertion of needle and injection of substance into lumbar interlaminar subarachnoid space using imaging guidance
64483	Transforaminal injection of anesthetic agent and steroid into epidural space of lumbar spine using CT guidance; Transforaminal injection of anesthetic agent and steroid into epidural space of lumbar spine using fluoroscopic guidance; Transforaminal injection of anesthetic agent and steroid into epidural space of lumbar spine using imaging guidance; Transforaminal injection of anesthetic agent into epidural space of lumbar spine using CT guidance; Transforaminal injection of anesthetic agent into epidural space of lumbar spine using fluoroscopic guidance; Transforaminal injection of anesthetic agent into epidural space of lumbar spine using imaging guidance; Transforaminal injection of steroid into epidural space of lumbar spine using CT guidance; Transforaminal

	injection of steroid into epidural space of lumbar spine using fluoroscopic guidance; Transforaminal injection of steroid into epidural space of lumbar spine using imaging guidance
64484	Transforaminal injection of anesthetic agent and steroid into epidural space of lumbar spine using CT guidance; Transforaminal injection of anesthetic agent and steroid into epidural space of lumbar spine using fluoroscopic guidance; Transforaminal injection of anesthetic agent and steroid into epidural space of lumbar spine using imaging guidance; Transforaminal injection of anesthetic agent into epidural space of lumbar spine using CT guidance; Transforaminal injection of anesthetic agent into epidural space of lumbar spine using fluoroscopic guidance; Transforaminal injection of anesthetic agent into epidural space of lumbar spine using imaging guidance; Transforaminal injection of steroid into epidural space of lumbar spine using CT guidance; Transforaminal injection of steroid into epidural space of lumbar spine using fluoroscopic guidance; Transforaminal injection of steroid into epidural space of lumbar spine using imaging guidance
0230T	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with ultrasound guidance, lumbar or sacral; single level
0231T	Injections of anesthetic agent and/or steroid into lower or sacral spinal canal using ultrasound guidance
64999	Nervous system procedure
76000	Imaging guidance for procedure, up to 1 hour

## **Service: Sacroiliac (SI) Joint Injections**

### **General Guidelines**

- **Units, Frequency, & Duration: Units, Frequency, & Duration:** When the medical necessity criteria are met, a total of 4 sacroiliac joint injections per episode of pain per region may be performed in 6 months.
- **Criteria for Subsequent Requests:** If the first injection resulted in a 50% improvement of symptoms for three months, a second injection might be appropriate. If the first injection was not beneficial, a second injection is not recommended.<sup>39</sup>
- **Recommended Clinical Approach:** None.
- **Exclusions:** None.

### **Medical Necessity Criteria**

#### **Indications**

- **Sacroiliac (SI) joint injections** are considered appropriate if **ALL** of the following are **TRUE**:
- ◆ The patient had a positive response to 3 or more SI joint provocative tests.<sup>16</sup> SI joint provocative tests include distraction, compression, thigh thrust, Gaenslen's, FABER maneuver, and sacral thrust.
  - ◆ The patient failed to show significant improvement in pain or disability level due to symptoms, despite receiving more than 6 weeks of conservative management (conservative management includes a combination of physical therapy, provider-directed home exercise program, and anti-inflammatory/pain management medications or oral steroids).

#### **Non-Indications**

- **Sacroiliac (SI) joint injections** may not be appropriate if **ANY** of the following is **TRUE**<sup>16</sup>:
- ◆ Allergy to cortisone injections
  - ◆ Coagulopathy or recent use of blood-thinning agents

### **Site of Service Criteria**

Outpatient

**Procedure Codes (HCPCS/CPT)**

HCPCS Code	Code Description/Definition
27096	Injection of anesthetic into sacroiliac joint using imaging guidance; Injection of anesthetic into sacroiliac joint with arthrography using imaging guidance; Injection of steroid into sacroiliac joint using imaging guidance; Injection of steroid into sacroiliac joint with arthrography using imaging guidance; Injection of anesthetic into sacroiliac joint using fluoroscopic guidance; Injection of anesthetic into sacroiliac joint with arthrography using fluoroscopic guidance; Injection of steroid into sacroiliac joint using fluoroscopic guidance; Injection of anesthetic into sacroiliac joint using computed tomography (CT) guidance; Injection of anesthetic into sacroiliac joint with arthrography using computed tomography (CT) guidance; Injection of steroid into sacroiliac joint using computed tomography (CT) guidance; Injection of steroid into sacroiliac joint with arthrography using computed tomography (CT) guidance; Injection of steroid into sacroiliac joint with arthrography using fluoroscopic guidance
76000	Imaging guidance for procedure, up to 1 hour

## **Service: Subsequent Facet Injection/Medial Branch Block**

### **General Guidelines**

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** For second injections, the previous injection must result in greater than 80% relief of symptoms. If the patient has had greater than or equal to 2 injections, the previous injection must result in 50% improvement of symptoms for 3 months.<sup>15-21</sup>
- **Recommended Clinical Approach:** Blocks to be performed with imaging (fluoroscopy or ultrasound with Doppler) guidance. Medial branch blocks with a double-injection technique and pain improvement threshold of 80% may be used to establish the diagnosis of lumbar facet-mediated pain and predict a favorable response to medial nerve ablation by thermocoagulation. There is no evidence to support the use of diagnostic facet blocks as a predictor of lumbar fusion outcome in patients with chronic low back pain. Pregnancy is a relative contraindication.
- **Exclusions:** None.

### **Medical Necessity Criteria**

#### **Indications**

- **Subsequent Facet Injection/Medial Branch Block** may be considered appropriate if **ALL** of the following are **TRUE**:
- ◆ The patient fails to show significant improvement in pain or disability level due to symptoms despite receiving more than 12 weeks of conservative management (conservative management includes a combination of physical therapy, provider-directed home exercise program, and anti-inflammatory/pain management medications or oral steroids).<sup>22</sup>
  - ◆ The patient had advanced imaging that revealed either facet arthritis or cervical/thoracic spondylosis (arthritis) at the level requested.

#### **Non-Indications**

- **Subsequent Facet Injection/Medial Branch Block** may not be appropriate if **ANY** of the following is **TRUE**:<sup>7</sup>
- ◆ Systemic infection
  - ◆ Local infection at the procedure site
  - ◆ Coagulopathy or recent use of blood-thinning agents
  - ◆ Previous fusion at the symptomatic level

**Site of Service Criteria**

Outpatient

**Procedure Codes (HCPCS/CPT)**

HCPCS Code	Code Description/Definition
64493	Injection of diagnostic agent into nerve of single lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into nerve of single lumbar paravertebral facet joint using imaging guidance; Injection of diagnostic agent into single lumbar paravertebral facet joint using imaging guidance; Injection of therapeutic agent into nerve of single lumbar paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into single lumbar paravertebral facet joint using computed tomography (CT) guidance
64494	Injection of diagnostic agent into nerve of second lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into nerve of second lumbar paravertebral facet joint using imaging guidance; Injection of diagnostic agent into second lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into second lumbar paravertebral facet joint using imaging guidance; Injection of therapeutic agent into nerve of second lumbar paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into second lumbar paravertebral facet joint using computed tomography (CT) guidance
64495	Injection of diagnostic agent into nerve of third and any additional lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into nerve of third and any additional lumbar paravertebral facet joint using imaging guidance; Injection of diagnostic agent into nerve of third and any additional sacral paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into nerve of third and any additional sacral paravertebral facet joint using imaging guidance; Injection of diagnostic agent into third and any additional lumbar paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into

	third and any additional lumbar paravertebral facet joint using imaging guidance; Injection of diagnostic agent into third and any additional sacral paravertebral facet joint using fluoroscopic guidance; Injection of diagnostic agent into third and any additional sacral paravertebral facet joint using imaging guidance; Injection of therapeutic agent into nerve of third and any additional lumbar paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into nerve of third and any additional sacral paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into third and any additional lumbar paravertebral facet joint using computed tomography (CT) guidance; Injection of therapeutic agent into third and any additional sacral paravertebral facet joint using computed tomography (CT) guidance
0216T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, lumbar or sacral; single level
64999	Unlisted procedure, nervous system
0213T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; single level
0214T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; second level (list separately in addition to code for primary procedure)
0215T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; third and any additional level(s) (list separately in addition to code for primary procedure)

## **Service: Initial/Subsequent Radiofrequency Ablation (RFA)**

### General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** Patients may receive a second radiofrequency ablation after 6 months if there was 50% improvement of symptoms.<sup>15-21</sup>
- **Recommended Clinical Approach:** Radiofrequency ablation for the facet joint may provide pain relief in 45% -60% of patients.<sup>23,40</sup>
- **Exclusions:** None.

### Medical Necessity Criteria

#### Indications

- **Initial/subsequent Radiofrequency Ablation (RFA)** is considered appropriate if **ALL** the following are **TRUE**<sup>1,41</sup>:
- ◆ The patient has had 2 or more medial branch diagnostic nerve blocks to locate the target level and they have provided 50% or greater relief of symptoms.
  - ◆ The patient failed to show significant improvement in pain or disability level due to symptoms, despite receiving more than 6 weeks of conservative management (conservative management includes a combination of physical therapy, provider-directed home exercise program, and anti-inflammatory/pain management medications or oral steroids).

#### Non-Indications

- **Initial/subsequent Radiofrequency Ablation (RFA)** may not be appropriate if **ANY** of the following is **TRUE**<sup>24-25</sup>:
- ◆ Infection at the injection site
  - ◆ The patient is allergic to local anesthetic.

### Site of Service Criteria

Outpatient

### Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
63435	Destruction of single lumbar paravertebral facet joint nerve using neurolytic agent and using fluoroscopic guidance; Destruction of single lumbar paravertebral

	facet joint nerve using neurolytic agent and using imaging guidance; Destruction of single sacral paravertebral facet joint nerve using neurolytic agent and using fluoroscopic guidance; Destruction of single sacral paravertebral facet joint nerve using neurolytic agent and using imaging guidance
64636	Destruction of each additional lumbar paravertebral facet joint nerve using neurolytic agent and using computed tomography (CT) guidance; Destruction of each additional lumbar paravertebral facet joint nerve using neurolytic agent and using fluoroscopic guidance; Destruction of each additional lumbar paravertebral facet joint nerve using neurolytic agent and using imaging guidance; Destruction of each additional sacral paravertebral facet joint nerve using neurolytic agent and using computed tomography (CT) guidance; Destruction of each additional sacral paravertebral facet joint nerve using neurolytic agent and using fluoroscopic guidance; Destruction of each additional sacral paravertebral facet joint nerve using neurolytic agent and using imaging guidance
64999	Unlisted procedure, nervous system

## **Service: Facet Cyst Aspiration**

### General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests<sup>42</sup>:** Patients may receive a second Cyst Aspiration/Rupture after 3 months if there was a 50% or more improvement of symptoms.
- **Recommended Clinical Approach:** Blocks to be performed with imaging (fluoroscopy or ultrasound with Doppler) guidance. Pregnancy is a relative contraindication.
- **Exclusions:** None.

### Medical Necessity Criteria

#### Indications

→ **Facet Cyst Aspiration** is considered appropriate if **ALL** the following are **TRUE<sup>42</sup>**:

- ◆ If this is the first request for a Cyst Aspiration and **ALL** of the following are true:
  - Advanced diagnostic imaging confirms compression or displacement of the nerve root.
  - Clinical and physical symptoms related to the synovial cyst are documented.
- ◆ If this is a subsequent request for a Cyst Aspiration and **ALL** of the following are true:
  - Experience greater than or equal to 50% or more consistent improvement in pain for at least 3 months.

#### Non-Indications

→ **Facet Cyst Aspiration** may not be appropriate if **ANY** of the following is **TRUE<sup>24-25</sup>**:

- ◆ Infection at the injection site
- ◆ The patient is allergic to local anesthetic.

### Site of Service Criteria

Outpatient

### Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
64999	Unlisted procedure, nervous system

**Service: Intradiscal Biacuplasty, Percutaneous Intradiscal Radiofrequency Thermocoagulation (PIRFT), or Intradiscal Electrothermal Therapy (IDET)**

### General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach:** Intradiscal electrothermal therapy may be appropriate in a select subgroup of people who have chronic low back pain for at least 3–6 months that is attributed to a spinal disc (discogenic pain).<sup>26</sup>
- **Exclusions:** None.

### Medical Necessity Criteria

#### Indications

→ **Intradiscal biacuplasty, PIRFT, or IDET** are considered appropriate if **ALL** of the following are **TRUE**!

- ◆ The patient fails to show significant improvement in pain or disability level due to symptoms despite receiving **3 or more months** of conservative management (conservative management includes a combination of physical therapy, provider-directed home exercise program, and anti-inflammatory/pain management medications or oral steroids).
- ◆ At least one of the following is **TRUE**:
  - Advanced imaging found an annular tear
  - Disc disease is limited to one or two levels
  - Disc height is greater than or equal to 50% of normal

#### Non-Indications

→ **Intradiscal biacuplasty, PIRFT, or IDET** may not be appropriate if **ANY** of the following is **TRUE**!

- ◆ Severe disc degeneration.
- ◆ Spinal stenosis.
- ◆ Spinal instability such as spondylolisthesis.
- ◆ The patient previously had back surgery or intradiscal electrothermal therapy.

### Site of Service Criteria

Outpatient

### Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
22526	Bilateral percutaneous intradiscal electrothermal annuloplasty of a single level of spine using fluoroscopic guidance; Unilateral percutaneous intradiscal electrothermal annuloplasty of a single level of spine using fluoroscopic guidance
22527	Bilateral percutaneous intradiscal electrothermal annuloplasty of a single additional level using fluoroscopic guidance; Bilateral percutaneous intradiscal electrothermal annuloplasty of multiple additional levels using fluoroscopic guidance; Unilateral percutaneous intradiscal electrothermal annuloplasty of multiple additional levels using fluoroscopic guidance; Unilateral percutaneous intradiscal electrothermal annuloplasty of single additional level using fluoroscopic guidance

## Surgical Management

### **Service: Lumbar Fusion**

#### General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach:** Surgery for low back pain (LBP) may include 1 or 2 levels of lumbar fusion. There is no established consensus or set of criteria regarding timing of surgical intervention. The literature shows no significant clinical difference in outcomes between comprehensive inpatient rehabilitation and fusion for chronic LBP. Advanced imaging is recommended before surgical intervention
- **Exclusions:** Lumbar fusion is not recommended as a routine treatment after discectomy.

#### Medical Necessity Criteria

#### **Indications**

- **Lumbar fusion for back pain** is considered appropriate if **ALL** of the following are **TRUE**<sup>43</sup>:
  - ◆ The patient has **ANY** positive findings from the [clinical presentation](#) and [typical physical exam findings](#) lists.
  - ◆ The patient has failed to show significant improvement in pain or disability level due to symptoms despite receiving more than 6-12 months of non-surgical management (i.e., a combination of physical therapy, provider-directed home exercise programs, facet injections/MBBBs/ESIs, and anti-inflammatory/pain management medication or oral steroids).
  - ◆ The patient has severe pain or disability due to symptoms or has progressive signs and symptoms.
  - ◆ There is radiographic evidence of moderate to severe discogenic disease or spondylosis.
  - ◆ The patient has completed cognitive behavioral therapy.
- **Revision surgery** for pseudarthrosis following an initial spine surgery may be appropriate if **ALL** the following conditions are **TRUE**<sup>44</sup>:
  - It has been at least one year since the previous surgery
  - There is clear radiographic evidence of pseudoarthrosis
  - The patient exhausted available conservative treatment measures.

## Non-Indications

None.

## Site of Service Criteria

Inpatient

## Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
22533	Arthrodesis of lumbar vertebra with minimal discectomy; Arthrodesis of lumbar vertebra by lateral extracavitary technique with minimal discectomy
+22534	Arthrodesis of lumbar vertebra with minimal discectomy; Arthrodesis of lumbar vertebra with minimal discectomy
22558	Arthrodesis by anterior interbody technique of lumbar region with discectomy; Arthrodesis by anterior interbody technique of lumbar region, with minimal discectomy
+22585	Arthrodesis by anterior interbody technique of vertebral region with discectomy; Arthrodesis by anterior interbody technique of each additional interspace, with minimal discectomy
22612	Arthrodesis of lumbar vertebral segment by posterior and lateral transverse technique; Arthrodesis of lumbar vertebral segment by posterior technique; Arthrodesis of lumbar vertebral segment by posterolateral and lateral transverse technique; Arthrodesis of lumbar vertebral segment by posterolateral technique
+22614	Arthrodesis of each additional vertebral segment of single level by posterior technique; Arthrodesis of each additional vertebral segment of single level by posterolateral technique; Arthrodesis of vertebral segment by posterior technique; Arthrodesis of vertebral segment by posterolateral technique
22630	Arthrodesis of lumbar vertebral segment by posterior interbody technique with discectomy; Arthrodesis of lumbar vertebral segment by posterior interbody technique with laminectomy; Arthrodesis of lumbar vertebral segment by posterior interbody technique with laminectomy and discectomy; Arthrodesis of single

	lumbar vertebral interspace by posterior interbody technique with discectomy; Arthrodesis of single lumbar vertebral interspace by posterior interbody technique with laminectomy; Arthrodesis of single lumbar vertebral interspace by posterior interbody technique with laminectomy and discectomy
+22632	Arthrodesis of each additional vertebral interspace by posterior interbody technique with discectomy; Arthrodesis of each additional vertebral interspace by posterior interbody technique with laminectomy; Arthrodesis of each additional vertebral interspace by posterior interbody technique with laminectomy and discectomy; Arthrodesis of lumbar vertebral segment by posterior interbody technique with discectomy; Arthrodesis of lumbar vertebral segment by posterior interbody technique with laminectomy; Arthrodesis of lumbar vertebral segment by posterior interbody technique with laminectomy and discectomy
22633	Arthrodesis of single lumbar vertebral interspace by combined posterior technique and posterior interbody technique, with discectomy; Arthrodesis of single lumbar vertebral interspace by combined posterior technique and posterior interbody technique, with laminectomy; Arthrodesis of single lumbar vertebral interspace by combined posterior technique and posterior interbody technique, with laminectomy and discectomy; Arthrodesis of single lumbar vertebral interspace by combined posterolateral technique and posterior interbody technique, with discectomy; Arthrodesis of single lumbar vertebral interspace by combined posterolateral technique and posterior interbody technique, with laminectomy; Arthrodesis of single lumbar vertebral interspace by combined posterolateral technique and posterior interbody technique, with laminectomy and discectomy; Arthrodesis of single lumbar vertebral segment by combined posterior technique and posterior interbody technique, with laminectomy; Arthrodesis of single lumbar vertebral segment by combined posterior technique and posterior interbody technique, with laminectomy and discectomy; Arthrodesis of single lumbar vertebral segment by combined posterolateral technique and posterior interbody technique, with discectomy

+22634	Arthrodesis by combined technique of lumbar vertebral segment with discectomy; Arthrodesis by combined technique of lumbar vertebral segment with laminectomy; Arthrodesis by combined technique of lumbar vertebral segment with laminectomy and discectomy; Arthrodesis of each additional lumbar vertebral interspace by combined posterolateral technique and posterior interbody technique, with discectomy; Arthrodesis of each additional lumbar vertebral interspace by combined posterolateral technique and posterior interbody technique, with laminectomy; Arthrodesis of each additional lumbar vertebral interspace by combined posterolateral technique and posterior interbody technique, with laminectomy and discectomy; Arthrodesis of each additional lumbar vertebral segment by combined posterior technique and posterior interbody technique, with discectomy; Arthrodesis of each additional lumbar vertebral segment by combined posterior technique and posterior interbody technique, with laminectomy; Arthrodesis of each additional lumbar vertebral segment by combined posterior technique and posterior interbody technique, with laminectomy and discectomy; Arthrodesis of each additional lumbar vertebral segment by combined posterolateral technique and posterior interbody technique, with discectomy; Arthrodesis of each additional lumbar vertebral segment by combined posterolateral technique and posterior interbody technique, with laminectomy; Arthrodesis of each additional lumbar vertebral segment by combined posterolateral technique and posterior interbody technique, with laminectomy and discectomy
22865	Removal of artificial disc from total disc arthroplasty of lumbar vertebral interspace by anterior approach; Removal of artificial disc from total disc arthroplasty of single lumbar vertebral interspace by anterior approach
+63052	Laminectomy, facetectomy, or foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s] [eg, spinal or lateral recess stenosis]), during posterior interbody arthrodesis, lumbar; single vertebral segment (List separately in addition to code for primary procedure)
+63053	Laminectomy, facetectomy, or foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda

	equina and/or nerve root[s] [eg, spinal or lateral recess stenosis]), during posterior interbody arthrodesis, lumbar; each additional vertebral segment (List separately in addition to code for primary procedure)
63087	Vertebral corpectomy lumbar (vertebral body resection), partial or complete, anterior approach with decompression of spinal cord and/or nerve root(s)
63088	Vertebral corpectomy lumbar (vertebral body resection), partial or complete, anterior approach with decompression of spinal cord and/or nerve root(s)
22840	Posterior non segmental instrumentation
22842	Posterior segmental instrumentation 3-6 vertebral segments
22843	Posterior segmental instrumentation 7-12 vertebral segments
22844	Posterior segmental instrumentation >/13 vertebral segments
22845	Anterior Instrumentation 2-3 vertebral segments
22846	Anterior Instrumentation 4-7 vertebral segments
22847	Anterior Instrumentation >/8 vertebral segments
22853	Insertion of biomechanical interbody device with arthrodesis
22854	Insertion of biomechanical device with corpectomy defect w/ arthrodesis

## Service: Sacroiliac (SI) Joint Fusion

### General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach:** SI joint fusion may be appropriate for patients with low back pain originating from the SI joint and that does not improve with nonoperative treatment.<sup>22</sup>
- **Exclusions:** None.

### Medical Necessity Criteria

#### Indications

- **SI joint fusion** is considered appropriate if **ALL** of the following are **TRUE**:
- ◆ The patient has **ANY** positive findings from the [clinical presentation](#) and [typical physical exam findings](#) lists.
  - ◆ The patient fails to show significant improvement in pain or disability level despite receiving non-surgical management for more than 6 months (i.e., a combination of physical therapy, provider-directed home exercise programs, facet injections/MBBB/ESIs, and anti-inflammatory/pain management medications or oral steroids).
  - ◆ The patient has severe pain or disability due to symptoms or has progressive signs and symptoms.
  - ◆ The patient has a positive response to 3 or more provocative SI joint tests.<sup>45</sup> Provocative SI joint tests include distraction, compression, thigh thrust, Gaenslen's, FABER maneuver, and sacral thrust.
  - ◆ There was a positive response from SI joint injections.

#### Non-Indications

None.

#### Site of Service Criteria

Inpatient

#### Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
27279	Percutaneous arthrodesis of sacroiliac joint with insertion

	of fixation device using imaging guidance
27280	Open arthrodesis of sacroiliac joint

## Surgical Risk Factors

### Patient Medical Risk Stratification

Patient Risk Score	Patient Characteristic	Min Range	Max Range	Guidance
<b>1- Very Low Risk</b>	No known medical problems			
<b>2- Low Risk</b>	Hypertension		180/110 mm Hg	
<b>2- Low Risk</b>	Asthma	peak flow >80% of predicted or personal best value		
<b>2- Low Risk</b>	Prior history of alcohol abuse			Screen for liver disease and malnutrition
<b>2- Low Risk</b>	Prior history of tobacco use			
<b>3- Intermediate Risk</b>	Asthma	peak flow <80% of predicted or personal best value		
<b>3- Intermediate Risk</b>	Active alcohol abuse			
<b>3- Intermediate Risk</b>	Age	65	75	
<b>3- Intermediate Risk</b>	History of treated, stable coronary artery disease (CAD)			
<b>3- Intermediate Risk</b>	Stable atrial fibrillation			
<b>3- Intermediate Risk</b>	Diabetes mellitus	HbA1C >7%		
<b>3- Intermediate Risk</b>	Morbid obesity	BMI 30	BMI 40	
<b>3- Intermediate Risk</b>	Anemia	hemoglobin <11 (females), <12 (males)		Workup to identify etiology
<b>3- Intermediate Risk</b>	HIV	CD4 <200 cells/mm <sup>3</sup>		Get clearance from HIV specialist

<b>3- Intermediate Risk</b>	Rheumatologic disease			Preoperative consultation with rheumatologist re: perioperative medication management
<b>3- Intermediate Risk</b>	Peripheral vascular disease or history of peripheral vascular bypass	ankle-brachial pressure index (ABPI) <0.9		Preoperative consultation with vascular surgeon
<b>3- Intermediate Risk</b>	History of venous thromboembolism (VTE)			
<b>3- Intermediate Risk</b>	Well-controlled obstructive sleep apnea			
<b>3- Intermediate Risk</b>	Malnutrition	transferrin <200 mg/dL albumin <3.5 g/dL prealbumin <22.5 mg/dL total lymphocyte count <1200-1500 cell/mm <sup>3</sup> BMI <18		Preoperative consultation with nutritionist
<b>3- Intermediate Risk</b>	Active tobacco Use			Enroll patient in smoking cessation program
<b>4- High Risk</b>	Diabetes mellitus with complications	HbA1c >8%		
<b>4- High Risk</b>	Age	76	85	
<b>4- High Risk</b>	Oxygen dependent pulmonary disease			
<b>4- High Risk</b>	Sickle cell anemia			
<b>4- High Risk</b>	Obesity	BMI 40		
<b>4- High Risk</b>	Cirrhosis, history of hepatic decompensation or variceal bleeding			
<b>4- High Risk</b>	Impaired cognition; dementia			

<b>4- High Risk</b>	Compensated CHF			
<b>4- High Risk</b>	Cerebrovascular disease			
<b>4- High Risk</b>	Uncontrolled or suspected obstructive sleep apnea (OSA)			
<b>4- High Risk</b>	Renal insufficiency	serum creatinine >1.5 mg/dL or creatinine clearance <100 mL/min		
<b>4- High Risk</b>	Opioid dependence			
<b>4- High Risk</b>	End Stage Liver Disease			
<b>4- High Risk</b>	Uncontrolled Seizure Disorder			
<b>4- High Risk</b>	History of Malignant Hyperthermia			
<b>5- Very High Risk</b>	Cardiovascular: unstable angina, recent myocardial infarction (60 days), uncontrolled atrial fibrillation or other high-grade abnormal rhythm, severe valvular disease, decompensated heart failure			
<b>5- Very High Risk</b>	Primary pulmonary hypertension			Preoperative consultation with pulmonologist warranted
<b>5- Very High Risk</b>	Cirrhosis or severe liver disease, history of hepatic decompensation or variceal bleeding			
<b>5- Very High Risk</b>	Severe frailty, dependence for ADLs, or history of 3 or more falls in last 6 mos			
<b>5- Very High Risk</b>	Obesity		BMI >50	
<b>5- Very High Risk</b>	Age		>85	

<b>5- Very High Risk</b>	History of VTE with CI to anticoagulation, failure of anticoagulation, cessation of anticoagulation therapy secondary to bleeding			Preoperative consultation with hematologist or internist
<b>5- Very High Risk</b>	Renal failure requiring dialysis			
<b>5- Very High Risk</b>	Immunosuppression			
<b>5- Very High Risk</b>	Chronic Pain			

## Postoperative Care

**Service: Physical Therapy**

### General Guidelines

- **Units, Frequency, & Duration:** Physical therapy may be prescribed 4–6 weeks after lumbar fusion surgery. Education and activity supervision may begin immediately. Formal spine rehabilitation may begin later to allow fusion healing (approximately 8–12 weeks). Physical therapy may be appropriate for up to 6 months post fusion.<sup>46</sup>
- **Criteria for Subsequent Requests:** The patient should be progressing towards goals in the physical therapy plan without fully obtaining all goals.
- **Recommended Clinical Approach:** None.
- **Exclusions:** None.

### Medical Necessity Criteria

#### Indications

- **Post-acute physical therapy** is considered appropriate if **ANY** of the following is **TRUE**<sup>46</sup>:
- ◆ The patient underwent lumbar fusion.

#### Non-Indications

None.

### Site of Service Criteria

Outpatient

### Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
97010	Application of hot or cold packs
97012	Application of mechanical traction
97014	Application of electrical stimulation
97016	Application of vasopneumatic devices
97018	Application of paraffin bath

97022	Application of whirlpool
97024	Application of diathermy
97026	Application of infrared modality
97028	Application of ultraviolet modality
97032	Application of manual electrical stimulation
97033	Application of iontophoresis
97034	Application of contrast baths
97035	Application of ultrasound modality
97036	Application of Hubbard tank
97039	Modality service
97110*	Therapeutic exercises to develop strength and endurance, range of motion and flexibility
97112	Neuromuscular reeducation of movement, balance, coordination, kinesthetic sense, posture, and proprioception for sitting and standing activities
97113	Aquatic therapy with therapeutic exercises
97116	Gait training including stair climbing
97124	Massage including effleurage and petrissage; Massage including effleurage and tapotement; Massage including effleurage, petrissage and tapotement; Massage including petrissage and tapotement
97139	Therapeutic procedure
97140	Manual therapy techniques
97150	Group therapeutic procedures
97164	Physical therapy re-evaluation of established plan of care, high complexity, typical time with patient 20 minutes; Physical therapy re-evaluation of established plan of care, high complexity, typical time with patient and family 20 minutes; Physical therapy re-evaluation of established plan of care, high complexity, typical time with patient's family 20

	minutes
97530	Direct therapeutic activities with use of dynamic activities to improve functional performance, each 15 minutes
97535	Home management training, direct one-on-one contact, each 15 minutes; Self-care management training, direct one-on-one contact, each 15 minutes
97537	Community reintegration training, direct one-on-one contact, each 15 minutes; Work reintegration training, direct one-on-one contact, each 15 minutes
97542	Wheelchair management, each 15 minutes
97545	Work conditioning, initial 2 hours; Work hardening, initial 2 hours
97546	Work conditioning, each additional hour; Work hardening, each additional hour
97750	Physical performance measurement with written report, each 15 minutes; Physical performance test with written report, each 15 minutes
97755	Assistive technology assessment with written report, direct one-on-one contact, each 15 minutes
97760	Initial orthotic management and training with assessment and fitting of lower extremities and trunk, each 15 minutes; Initial orthotic management and training with assessment and fitting of lower extremities, each 15 minutes; Initial orthotic management and training with assessment and fitting of lower extremity and trunk, each 15 minutes; Initial orthotic management and training with assessment and fitting of lower extremity, each 15 minutes; Initial orthotic management and training with assessment and fitting of trunk, each 15 minutes; Initial orthotic management and training with assessment and fitting of upper and lower extremities and trunk, each 15 minutes
97761	Initial prosthetic training of lower extremities, each 15 minutes; Initial prosthetic training of lower extremity, each 15 minutes

	<p>Initial prosthetic training of upper and lower extremities, each 15 minutes;</p> <p>Initial prosthetic training of upper extremities, each 15 minutes;</p> <p>Initial prosthetic training of upper extremity, each 15 minutes</p>
97763	<p>Subsequent orthotic management and training of lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management and training of lower extremity and trunk, each 15 minutes</p> <p>Subsequent orthotic management and training of lower extremity, each 15 minutes</p> <p>Subsequent orthotic management and training of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management and training of upper extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management and training of upper extremities, each 15 minutes</p> <p>Subsequent orthotic management and training of upper extremity and trunk, each 15 minutes</p> <p>Subsequent orthotic management and training of upper extremity, each 15 minutes</p> <p>Subsequent orthotic management of lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management of lower extremity and trunk, each 15 minutes</p> <p>Subsequent orthotic management of lower extremity, each 15 minutes</p> <p>Subsequent orthotic management of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management of upper extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic management of upper extremities, each 15 minutes</p> <p>Subsequent orthotic management of upper extremity and trunk, each 15 minutes</p> <p>Subsequent orthotic management of upper extremity, each 15 minutes</p> <p>Subsequent orthotic training of lower extremity, each 15</p>

	<p>minutes</p> <p>Subsequent orthotic training of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic training of upper extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic training of upper extremities, each 15 minutes</p> <p>Subsequent orthotic training of upper extremity and trunk, each 15 minutes</p> <p>Subsequent orthotic training of upper extremity, each 15 minutes</p> <p>Subsequent prosthetic management and training of lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of lower extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of lower extremity, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper extremities, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of upper extremity, each 15 minutes</p> <p>Subsequent prosthetic management of lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of lower extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of lower extremity, each 15 minutes</p> <p>Subsequent prosthetic management of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of upper extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of upper extremities, each 15 minutes</p>
--	--

	<p>Subsequent prosthetic management of upper extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management of upper extremity, each 15 minutes</p> <p>Subsequent prosthetic training of lower extremity, each 15 minutes</p> <p>Subsequent prosthetic training of upper and lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic training of upper extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic training of upper extremities, each 15 minutes</p> <p>Subsequent prosthetic training of upper extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic training of upper extremity, each 15 minutes</p> <p>Subsequent orthotic management and training of lower extremities, each 15 minutes</p> <p>Subsequent orthotic management of lower extremities, each 15 minutes</p> <p>Subsequent orthotic training of lower extremities and trunk, each 15 minutes</p> <p>Subsequent orthotic training of lower extremities, each 15 minutes</p> <p>Subsequent orthotic training of lower extremity and trunk, each 15 minutes</p> <p>Subsequent prosthetic management and training of lower extremities, each 15 minutes</p> <p>Subsequent prosthetic management of lower extremities, each 15 minutes</p> <p>Subsequent prosthetic training of lower extremities and trunk, each 15 minutes</p> <p>Subsequent prosthetic training of lower extremities, each 15 minutes</p> <p>Subsequent prosthetic training of lower extremity and trunk, each 15 minutes</p>
97799	Unlisted physical medicine/rehabilitation service or procedure
420	Physical Therapy

421	Physical Therapy: Visit Charge
422	Physical Therapy: Hourly Charge
423	Physical Therapy: Group Rate
424	Physical Therapy: Evaluation/Re-evaluation
429	Physical Therapy: Other Physical Therapy
97163	Evaluation of physical therapy, typically 45 minutes
97161	Evaluation of physical therapy, typically 20 minutes
97162	Evaluation of physical therapy, typically 30 minutes
97168	Re-evaluation of occupational therapy established plan of care, typically 30 minutes
97165	Evaluation of occupational therapy, typically 30 minutes
97166	Evaluation of occupational therapy, typically 45 minutes
97167	Evaluation of occupational therapy established plan of care, typically 60 minutes
G0151	Hhcp-serv of pt,ea 15 min

\*Default codes for suggested services

## **Service: Home Health Care**

### **General Guidelines**

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach:** Home health care may be recommended for postoperative care if outpatient treatment is not indicated based on hospital case management recommendations.
- **Exclusions:** None.

### **Medical Necessity Criteria**

#### **Indications**

- **Home health care** may be appropriate if **ALL** of the following are **TRUE**:
  - The patient lives with those who are unable to care for the patient postoperatively.
  - The patient underwent lumbar fusion.

#### **Non-Indications**

None.

### **Site of Service Criteria**

Home

### **Procedure Codes (HCPCS/CPT)**

<b>HCPCS Code</b>	<b>Code Description/Definition</b>
99509	Home visit for assistance with activities of daily living and personal care
99600	Unlisted home visit procedure; Unlisted home visit service
99334	Level 1 rest home visit for evaluation and management of established patient with minor and/or self-limited problem, including problem-focused interval history and physical examination, and straightforward medical decision-making, typical time with patient, family, and/or caregiver 15 minutes
G0129	Partial hosp prog service
G0283	Elec stim other than wound

## **Service: Inpatient Rehabilitation**

### **General Guidelines**

- **Units, Frequency, & Duration:** Postoperative rehabilitation is recommended to begin as soon as possible for all patients. No guidelines are available for the specific duration, timing, or frequency.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach:** There are no firmly established criteria for discharge appropriateness. Discharge depends upon medical stability, pain control, home situation, and if PT/OT goals were met. Some patients may require non-home discharge after surgery depending upon their age, comorbidities, and functional needs. Rehabilitation guidelines are not firmly established. There are no data available on outcomes nor firm recommendations on appropriate discharge destination. Post-acute care may be warranted. Cognitive behavioral therapy (CBT) may be included in postoperative rehabilitation programs.
- **Exclusions:** None.

### **Medical Necessity Criteria**

#### **Indications**

→ **Post-acute inpatient rehabilitation** is considered appropriate if **ALL** of the following is **TRUE**:

- ◆ **ANY** of the following are **TRUE**:
  - Neurologic deficit occurs postoperatively.
  - Postoperative complications.
  - Multiple medical comorbidities.
  - The patient requires maximum assistance for mobility.
  - The patient is a potentially unsafe discharge to home.
- ◆ The patient underwent lumbar fusion.

#### **Non-Indications**

None.

### **Site of Service Criteria**

Inpatient

### **Procedure Codes (HCPCS/CPT)**

<b>HCPCS Code</b>	<b>Code Description/Definition</b>
97799	Physical medicine service

## **Service: Skilled Nursing Facility (SNF)**

### General Guidelines

- **Units, Frequency, & Duration:** None.
- **Criteria for Subsequent Requests:** None.
- **Recommended Clinical Approach:** May be indicated for postoperative care in cases where the surgery occurred at an inpatient hospital, and outpatient physical therapy or home health care are not indicated.
- **Exclusions:** None.

### Medical Necessity Criteria

#### Indications

→ **Skilled nursing facilities (SNF)** are considered appropriate if **ALL** of the following are **TRUE**:

- ◆ The patient underwent lumbar fusion and **ANY** of the following is **TRUE**:
  - Neurologic deficit occurs postoperatively.
  - Postoperative complications.
  - Multiple medical comorbidities.
  - The patient requires maximum assistance for mobility.
  - The patient is a potentially unsafe discharge to home.

#### Non-Indications

None.

### Site of Service Criteria

Nursing facility

### Procedure Codes (HCPCS/CPT)

HCPCS Code	Code Description/Definition
99304	Level 1 initial nursing facility care for evaluation and management of patient with problem of low severity, including comprehensive history and physical examination, and medical decision-making of low complexity, typical time 25 minutes; Level 1 initial nursing facility care for evaluation and management of patient with problem of low severity, including detailed history and physical examination, and straightforward medical

	decision-making, typical time 25 minutes
99305	Level 2 initial nursing facility care for evaluation and management of patient with problem of moderate severity, including comprehensive history and physical examination, and medical decision-making of moderate complexity, typical time 35 minutes
99306	Level 3 initial nursing facility care for evaluation and management of patient with problem of high severity, including comprehensive history and physical examination, and medical decision-making of high complexity typical time 45 minutes
99307	Level 1 subsequent nursing facility care for evaluation and management of patient, including problem-focused interval history and physical examination, and straightforward medical decision-making, typical time 10 minutes; Level 1 subsequent nursing facility care for evaluation and management of patient, including problem-focused interval history and physical examination, typical time 10 minutes; Level 1 subsequent nursing facility care for evaluation and management of patient, including problem-focused interval history and straightforward medical decision-making, typical time 10 minutes; Level 1 subsequent nursing facility care for evaluation and management of patient, including problem-focused physical examination and straightforward medical decision-making, typical time 10 minutes
99308	Level 2 subsequent nursing facility care for evaluation and management of patient, including expanded problem-focused interval history and medical decision-making of low complexity, typical time 15 minutes; Level 2 subsequent nursing facility care for evaluation and management of patient, including expanded problem-focused interval history and physical examination, and medical decision-making of low complexity, typical time 15 minutes; Level 2 subsequent nursing facility care for evaluation and management of patient, including expanded problem-focused interval history and physical examination, typical time 15 minutes; Level 2 subsequent nursing facility care for evaluation and management of patient, including expanded problem-focused physical examination and medical

	decision-making of low complexity, typical time 15 minutes
99309	Level 3 subsequent nursing facility care for evaluation and management of patient, including detailed interval history and medical decision-making of moderate complexity, typical time 25 minutes; Level 3 subsequent nursing facility care for evaluation and management of patient, including detailed interval history and physical examination, and medical decision-making of moderate complexity. typical time 25 minutes; Level 3 subsequent nursing facility care for evaluation and management of patient, including detailed interval history and physical examination, typical time 25 minutes; Level 3 subsequent nursing facility care for evaluation and management of patient, including detailed physical examination and medical decision-making of moderate complexity, typical time 25 minutes
99310	Level 4 subsequent nursing facility care for evaluation and management of patient, including comprehensive interval history and medical decision-making of high complexity, typical time 35 minutes; Level 4 subsequent nursing facility care for evaluation and management of patient, including comprehensive interval history and physical examination, and medical decision-making of high complexity, typical time 35 minutes; Level 4 subsequent nursing facility care for evaluation and management of patient, including comprehensive interval history and physical examination, typical time 35 minutes; Level 4 subsequent nursing facility care for evaluation and management of patient, including comprehensive physical examination and medical decision-making of high complexity, typical time 35 minutes
99315	Nursing facility discharge day management, 30 minutes or less
99316	Nursing facility day management, more than 30 minutes
G0128	Corf skilled nursing service

## References

1. Kreiner DS, Matz P, Cho CH, et al. Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis & Treatment of Low Back Pain. 2020. <https://www.spine.org/Portals/0/assets/downloads/ResearchClinicalCare/Guidelines/LowBackPain.pdf>.
2. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. *Spine*. 2009;34(10):1066-1077.
3. Patel ND, Broderick DF, Burns J, et al. ACR Appropriateness Criteria Low Back Pain. *J Am Coll Radiol*. 2016;13(9):1069-1078. doi:10.1016/j.jacr.2016.06.008
4. Eck JC, Sharan A, Ghogawala Z, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 7: lumbar fusion for intractable low-back pain without stenosis or spondylolisthesis. *J Neurosurg Spine*. 2014;21(1):42-47.
5. Gaenslen FJ. Sacro-iliac arthrodesis: indications, author's technic and end-results. *JAMA*. 1927;10;89(24)2031-2035.
6. Dreyfuss P, Michaelsen M, Pauza K, McLarty J, Bogduk N. The value of medical history and physical examination in diagnosing sacroiliac joint pain. *Spine*. 1996;21(22)2594-2602.
7. Bartleson J, Maus T. Diagnostic and therapeutic spinal interventions: Facet joint interventions. *Neurology: Clinical Practice*. 2014;4(4):342-346. doi:10.1212/cpj.0000000000000044
8. Delitto A, George SZ, Van Dillen L, et al. Low back pain. *J Orthop Sports Phys Ther*. 2012;42(4):A1-A57. doi:10.2519/jospt.2012.42.4.A1
9. Kreiner DS, Matz P, Bono CM, et al. Guideline summary review: an evidence-based clinical guideline for the diagnosis and treatment of low back pain [published correction appears in *Spine J*. 2021 Feb 24;]. *Spine J*. 2020;20(7):998-1024. doi:10.1016/j.spinee.2020.04.006
10. American College of Radiology. ACR Appropriateness Criteria® Low Back Pain. ACR.org. Revised 2021.
11. American College of Radiology. ACR PRACTICE PARAMETER FOR PERFORMING AND INTERPRETING MAGNETIC RESONANCE IMAGING (MRI). ACR.org. Revised 2022 (Resolution 8).
12. Manchikanti L, Singh V, Cash KA, Pampati V, Datta S. Management of pain of post lumbar surgery syndrome: one-year results of a randomized, double-blind, active controlled trial of fluoroscopic caudal epidural injections. *Pain Physician*. 2010;13(6):509-21.
13. ACR Committee on Drugs and Contrast Media. ACR Manual on Contrast Media: 2022 Updates. ACR.org. 2022.

14. American College of Radiology. ACR–SPR PRACTICE PARAMETER FOR IMAGING PREGNANT OR POTENTIALLY PREGNANT ADOLESCENTS AND WOMEN WITH IONIZING RADIATION. ACR.org. Revised 2018 (Resolution 39).
15. Bogduk N, Dreyfuss P, Govind J. A Narrative Review of Lumbar Medial Branch Neurotomy for the Treatment of Back Pain. *Pain Med.* 2009;10:1035–1045
16. Lord SM, Barnsley L, Bogduk N. The utility of comparative local anesthetic blocks versus placebo-controlled blocks for the diagnosis of cervical zygapophysial joint pain. *Clin J Pain.* 1995 Sep;11(3): 208–213.
17. MacVicar J, Borowczyk J, MacVicar A, et al. Lumbar medial branch radiofrequency neurotomy in New Zealand. *Pain Medicine* 2013; 14: 639–645
18. Nath S, Nath CA, Pettersson K. Percutaneous lumbar zygapophysial (facet) joint neurotomy using radiofrequency current, in the management of chronic low back pain. A randomized double-blind trial. *Spine* 2008;33:1291–1297.
19. Schofferman J, Kine G. Effectiveness of repeated radiofrequency neurotomy for lumbar facet pain. *Spine* 2004;29: 2471–2473.
20. Tekin I, Mirzai H, Ok G, Erbuyun K, Vatansever D. A comparison of conventional and pulsed radiofrequency denervation in the treatment of chronic facet joint pain. *Clin J Pain* 2007;23: 524–529.
21. Van Kleef M, Barendse GA, Kessels A, et al. Randomized trial of radiofrequency lumbar facet denervation for chronic low back pain. *Spine* 1999; 24: 1937–1942.
22. Peh W. Image-guided facet joint injection. *Biomed Imaging Interv J.* 2011;7(1):e4. doi:10.2349/bij.7.1.e4.
23. Perolat R, Kastler A, Nicot B, et al. Facet joint syndrome: from diagnosis to interventional management. *Insights Imaging.* 2018;9(5):773–789.
24. Senthelal S, Dydyk AM, Mesfin FB. Ablative Nerve Block. [Updated 2021 Apr 19]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK499975/>.
25. Wray JK, Dixon B, Przkora R. Radiofrequency Ablation. [Updated 2020 Oct 21]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK482387/>.
26. Ackerman WE III, Ahmad M. The efficacy of lumbar epidural steroid injections in patients with lumbar disc herniations. *Anesth Analg.* 2007;104(5):1217–22.
27. Arden NK, Price C, Reading I, et al. A multicentre randomized controlled trial of epidural corticosteroid injections for sciatica: the WEST study. *Rheumatology (Oxford).* 2005;44(11):1399–1406.
28. Briggs VG, Li W, Kaplan MS, Eskander MS, Franklin PD. Injection treatment and back pain associated with degenerative lumbar spinal stenosis in older adults. *Pain Physician.* 2010;13(6):E347–55.

29. Buttermann GR. The effect of spinal steroid injections for degenerative disc disease. *Spine J*. 2004;4(5):495–505.
30. Candido KD, Raghavendra MS, Chinthagada M, Badiee S, Trepashko DW. A prospective evaluation of iodinated contrast flow patterns with fluoroscopically guided lumbar epidural steroid injections: the lateral parasagittal interlaminar epidural approach versus the transforaminal epidural approach. *Anesth Analg*. 2008;106(2):638–44.
31. Furman MB, Kothari G, Parikh T, Anderson JG, Khawaja A. Efficacy of fluoroscopically guided, contrast-enhanced lumbosacral interlaminar epidural steroid injections: a pilot study. *Pain Med*. 2010;11(9):1328–34.
32. Kim D, Brown J. Efficacy and safety of lumbar epidural dexamethasone versus methylprednisolone in the treatment of lumbar radiculopathy: a comparison of soluble versus particulate steroids. *Clin J Pain*. 2011;27(6):518–522.
33. Lee JH, Moon J, Lee SH. Comparison of effectiveness according to different approaches of epidural steroid injection in lumbosacral herniated disk and spinal stenosis. *J Back Musculoskelet Rehabil*. 2009;22(2):83–89.
34. Manchikanti L, Cash KA, McManus CD, Pampati V, Benyamin RM. Fluoroscopic lumbar interlaminar epidural injections in managing chronic lumbar axial or discogenic pain. *J Pain Res*. 2012;5:301–11.
35. Manchikanti L, Cash KA, McManus CD, Pampati V, Benyamin RM. Preliminary results of a randomized, double-blind, controlled trial of fluoroscopic lumbar interlaminar epidural injections in managing chronic lumbar discogenic pain without disc herniation or radiculitis. *Pain Physician*. 2010;13(4):E279–92.
36. Manchikanti L, Cash KA, McManus CD, Pampati V, Fellows B. Fluoroscopic caudal epidural injections with or without steroids in managing pain of lumbar spinal stenosis: one year results of randomized, double-blind, active-controlled trial. *J Spinal Disord Tech*. 2012;25(4):226–34.
37. Manchikanti L, Pakanati RR, Pampati V. Comparison of three routes of epidural steroid injections in low back pain. *Pain Digest*. 1999;9:277–85.
38. Park CH, Lee SH, Kim BI. Comparison of the effectiveness of lumbar transforaminal epidural injection with particulate and nonparticulate corticosteroids in lumbar radiating pain. *Pain Med*. 2010;11(11):1654–1658.
39. Rados I, Sakic K, Fingler M, Kapural L. Efficacy of interlaminar vs transforaminal epidural steroid injection for the treatment of chronic unilateral radicular pain: prospective, randomized study. *Pain Med*. 2011;12(9):1316–21.
40. McCormick ZL, Marshall B, Walker J, McCarthy R, Walega DR. Long-Term Function, Pain and Medication Use Outcomes of Radiofrequency Ablation for Lumbar Facet Syndrome. *Int J Anesth Anesth*. 2015;2(2):028.
41. Cohen SP, Bhaskar A, Bhatia A, et al. Consensus practice guidelines on interventions for lumbar facet joint pain from a multispecialty, international working group. *Reg Anesth Pain Med*. 2020;45(6):424–467. doi:10.1136/rapm-2019-101243

42. Cohen SP, Hurley RW, Buckenmaier CC 3rd, Kurihara C, Morlando B, Dragovich A. Randomized placebo-controlled study evaluating lateral branch radiofrequency denervation for sacroiliac joint pain. *Anesthesiology*. 2008;109(2):279-288. doi:10.1097/ALN.0b013e31817f4c7c
43. Eck JC, Sharan A, Ghogawala Z, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 7: lumbar fusion for intractable low-back pain without stenosis or spondylolisthesis. *J Neurosurg Spine*. 2014;21(1):42-47. doi:10.3171/2014.4.SPINE14270
44. Montenegro TS, Gonzalez GA, Saiegh FA, et al. Clinical outcomes in revision lumbar spine fusions: an observational cohort study. *Journal of Neurosurgery: Spine*. 2021;35(4):437-445. doi:10.3171/2020.12.SPINE201908
45. Wu L, Tafti D, Varacallo M. Sacroiliac Joint Injection. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021. <https://www.ncbi.nlm.nih.gov/books/NBK513245/>.
46. Debono B, Wainwright TW, Wang MY, et al. Consensus statement for perioperative care in lumbar spinal fusion: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *Spine J*. 2021;21(5):729-752. doi:10.1016/j.spinee.2021.01.001

## Clinical Guideline Revision History/Information

Original Date: September 1, 2020	
<b>Review History</b>	
April 21, 2021 (V1 - V5)	<b>Approving Physician:</b> Dr. Brian Covino
October 14, 2021 (V.6)	<b>Reviewing Physician:</b> Dr. Vijay Yanamadala <b>Approving Physician:</b> Dr. Brian Covino
December 29, 2022 (V.7)	<b>Reviewing Physician:</b> Dr. Vijay Yanamadala <b>Approving Physician:</b> Dr. Traci Granston