



Cohere Medicare Advantage Policy – Sleep Study/Polysomnography (PSG)

Clinical Guidelines for Medical Necessity Review

Version: 3.1
Revision Date: April 21, 2025

Important Notices

Notices & Disclaimers:

GUIDELINES ARE SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION-MAKING OF END USERS.

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine the medical necessity of services (the "**Guidelines**") for informational purposes only, and solely for use by Cohere's authorized "**End Users**". These Guidelines (and any attachments or linked third-party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision-making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice. This policy may be superseded by existing and applicable Centers for Medicare & Medicaid Services (CMS) statutes.

© 2025 Cohere Health, Inc. All Rights Reserved.

Other Notices:

HCPCS® and CPT® copyright 2025 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

Guideline Information:

Specialty Area: Sleep Medicine

Guideline Name: Cohere Medicare Advantage Policy - Sleep Study/Polysomnography (PSG)

Date of last literature review: 04/16/2025

Document last updated: 04/21/2025

Type: ☒ Adult (18+ yo) | ☒ Pediatric (0-17 yo)

Table of Contents

Important Notices	2
Medical Necessity Criteria	4
Service: Sleep Study/Polysomnography (PSG)	4
Benefit Category	4
Related CMS Documents	4
Recommended Clinical Approach	4
Evaluation of Clinical Harms and Benefits	5
Medical Necessity Criteria	6
Indications	6
Non-Indications	22
Level of Care Criteria	23
Procedure Codes (CPT/HCPCS)	23
Medical Evidence	26
References	28
Clinical Guideline Revision History/Information	32

Medical Necessity Criteria

Service: Sleep Study/Polysomnography (PSG)

Benefit Category

Diagnostic Tests (other)

Please Note: This may not be an exhaustive list of all applicable Medicare benefit categories for this item or service.¹⁻⁶

Related CMS Documents

Please refer to the [CMS Medicare Coverage Database](#) for the most current applicable CMS National Coverage.¹⁻⁶

- [National Coverage Determination. Continuous positive airway pressure \(CPAP\) therapy for obstructive sleep apnea \(OSA\) \(240.4\)](#)
- [National Coverage Determination. Sleep testing for obstructive sleep apnea \(OSA\) \(240.4.1\)](#)
- [Local Coverage Determination \(LCD\). Polysomnography \(L36593\)](#)
- [Local Coverage Determination \(LCD\). Polysomnography and other sleep studies \(L36902\)](#)
- [Billing and Coding Article. Polysomnography \(A56995\)](#)
- [Billing and Coding: Polysomnography and other sleep studies \(A57049\)](#)

Recommended Clinical Approach

Sleep study/polysomnography (PSG) is the continuous and simultaneous monitoring and recording of various physiological and pathophysiological parameters during sleep (e.g., brain waves, blood oxygen level, heart rate, breathing rate, and eye and leg movements) for 6 hours or more, together with the review and interpretation of the records by a physician. Results are reviewed and interpreted by a physician. A comprehensive sleep evaluation must include a sleep history (e.g., snoring, apneas, daytime sleepiness), body mass index (BMI), neck circumference, cardiopulmonary examination, and identification of co-morbid sleep disorders and medical conditions.

Sleep study/PSG is performed to diagnose various sleep disorders. The gold standard for diagnosing obstructive sleep apnea (OSA) is overnight or split-night technologist-attended facility-based PSG. Split-night studies are performed to diagnose strongly suspected OSA and evaluate response to continuous positive airway pressure (CPAP) treatment over a one-night period¹. Completing the CPAP titration study on a second night may be appropriate. Selected patients without severe co-morbid conditions may be appropriate candidates for a portable or home sleep study. Two tests may be used in addition to PSG testing. Multiple Sleep Latency Tests (MSLT) measure daytime sleepiness and aid in diagnosing types 1 and 2 narcolepsy and idiopathic hypersomnia. Maintenance of Wakefulness Tests (MWTs) measures the ability to stay awake.²

Evaluation of Clinical Harms and Benefits

Cohere Health uses the criteria below to ensure consistency in reviewing the conditions to be met for coverage of sleep study or polysomnography. This process helps prevent incorrect denials and inappropriate approvals of medically necessary services. Specifically, limiting incorrect approvals reduces the risks associated with unnecessary procedures, such as complications from surgery, infections, and prolonged recovery times.

The potential clinical harms of using these criteria may include:

- Generally, no major risks or clinical harms are associated with sleep study/ polysomnography, which is non-invasive and painless.
- The most common side effect is skin irritation caused by the adhesive used to attach test sensors to the skin.
- Some patients may find it uncomfortable or difficult to sleep in the laboratory setting with sensors attached to the skin.
- Some patients (e.g., trauma survivors, or those hypersensitive to sound) may experience anxiety due to the unfamiliar settings of the sleep lab and the noise from the recording medical instruments.
- Although most sleep studies are performed at night, night-shift workers who generally sleep during the daytime might find it difficult to sleep at night. This problem can be circumvented by adjusting the time of the procedure based on individualized sleep schedules.
- Increased healthcare costs and complications from the inappropriate use of emergency services and additional treatments.

The clinical benefits of using these criteria include:

- A comprehensive sleep study may help diagnose various sleep disorders (e.g., sleep-related breathing disorders, periodic limb movement disorders, narcolepsy, sleepwalking, insomnia, etc.), including potentially life-threatening conditions such as obstructive sleep apnea (OSA), or adjust treatment plans for known sleep disorders.⁸⁻⁹
- Enhanced overall patient satisfaction and healthcare experience.

This policy includes provisions for expedited reviews and flexibility in urgent cases to mitigate risks of delayed access. Evidence-based criteria are employed to prevent inappropriate denials and ensure that patients receive medically necessary care. The criteria aim to balance the need for effective treatment with minimizing potential harms, providing numerous clinical benefits, and helping avoid unnecessary complications from inappropriate care. In addition, using these criteria is likely to create a consistent set of review criteria, thereby supporting optimal patient outcomes and efficient healthcare utilization.

Medical Necessity Criteria

Indications

→ A **sleep study** is considered appropriate if **ANY** of the following are **TRUE**¹⁰:

- ◆ The sleep study/polysomnography (PSG) includes **ANY** of the following³:
 - A 1-4 lead electroencephalogram (EEG) to measure global neural encephalographic activity using electrodes placed on the scalp; **OR**
 - Electrooculogram (EOG) to measure eye movements using electrodes placed near the outer canthus of each eye; **OR**
 - A submental electromyogram (EMG) to measure submental electromyographic activity using electrodes placed over the mentalis, submental muscle, and/or masseter regions; **OR**
 - Rhythm electrocardiogram (ECG) with 2 or 3 chest leads; **OR**

- Nasal and/or oral airflow; **OR**
 - Ventilation and respiratory effort by chest-wall and abdominal movement are measured using strain gauges, piezoelectric belts, inductive plethysmography, impedance or inductance pneumography, endo esophageal pressure, or by intercostal EMG; **OR**
 - Gas exchange (oxygen saturation [SpO₂]) by oximetry, transcutaneous monitoring, or end-tidal gas analysis; **OR**
 - Extremity muscle activity, motor activity-movement using EMG; **OR**
 - Body positions via mercury switches or by direct observation; **OR**
 - Recordings of vibration (frequency and/or volume) may be recorded; **OR**
 - Transcutaneous CO₂, esophageal pH, penile tumescence, and bipolar EEG; **OR**
 - **ALL** of the following for PSG:
 - Sleep is recorded; **AND**
 - Sleep is staged; **OR**
- ◆ Home-based, non-attended sleep study with **ANY** of the following:
- **ALL** of the following:
 - For an initial home study when the patient is an adult with suspected sleep apnea, including excessive daytime sleepiness with **AT LEAST TWO** of the following^{[10,12-13](#)}:
 - ◆ Snoring loudly (e.g., enough to be heard through a closed door, waking others); **OR**
 - ◆ Daytime sleepiness (e.g., falling asleep while driving); **OR**
 - ◆ Observed sleep apnea, gasping, choking, or breathing that has stopped; **OR**
 - ◆ Diagnosed hypertension; **OR**
 - ◆ Body mass index (BMI) greater than or equal to 35 kg/m²; **OR**
 - ◆ Age 50 years or older; **OR**

- ◆ Large neck circumference (greater than 17 inches in men; greater than 16 inches in women); **AND**
- The patient is an adult with a high pretest probability of moderate to severe obstructive sleep apnea (OSA) with **ALL** of the following^{4,14}:
 - ◆ Sleep history and symptoms including, but not limited to, snoring, daytime sleepiness, observed apneas, choking or gasping during sleep, morning headaches; **AND**
 - ◆ Epworth Sleepiness Scale; **AND**
 - ◆ Physical examination that documents body mass index, neck circumference, and a focused cardiopulmonary and upper airway system evaluation; **AND**
 - ◆ The sleep study is recorded and staged³; **AND**
 - ◆ The sleep study is performed in conjunction with a comprehensive sleep evaluation⁴; **OR**
- A study is required that demonstrates less than 25% of central events are performed within 24 months of the first consultation for hypoglossal nerve stimulation (HGNS) implant¹⁵⁻¹⁶; **OR**
- Repeat home study with **ANY** of the following^{10-11,14}:
 - ◆ The first study was inconclusive due to technical or equipment failure; **OR**
 - ◆ The patient is unable to sleep or complete enough hours of sleep to allow a clinical diagnosis; **OR**
 - ◆ The results were inconclusive or ambiguous; **OR**
 - ◆ For re-evaluation due to weight change; **OR**
- ◆ Facility-based, technologist-attended PSG (type I) with **ALL** of the following:
 - **ANY** of the following:
 - Adult with suspected sleep apnea with **ALL** of the following:
 - ◆ Home sleep study is contraindicated; **AND**
 - ◆ For initial facility-based testing when the patient is an adult with suspected sleep apnea,

including excessive daytime sleepiness with **AT LEAST TWO** of the following¹⁰⁻¹²:

- Snoring loudly (e.g., enough to be heard through a closed door, waking others); **OR**
- Daytime sleepiness (e.g., falling asleep while driving); **OR**
- Observed sleep apnea, gasping, choking, or breathing that has stopped; **OR**
- Diagnosed hypertension; **OR**
- Body mass index (BMI) greater than or equal to 35 kg/m²; **OR**
- Age 50 years or older; **OR**
- Large neck circumference (greater than 17 inches in men; greater than 16 inches in women); **AND**

◆ **ANY** of the following co-morbid medical conditions¹⁰:

- Significant cardiopulmonary disease (forced expiratory volume in one second [FEV₁] % predicted of less than 70) or heart failure with left ventricular ejection fraction (LVEF) less than 45%^{8,17-18}; **OR**
- Potential respiratory muscle weakness due to neuromuscular conditions; **OR**
- History of stroke; **OR**
- Chronic opiate medication use¹¹; **OR**
- Concern for a significant non-respiratory sleep disorder(s) that require evaluation (e.g., disorders of central hypersomnolence⁹, sleep-related movement disorders) or that interfere with the accuracy of home sleep apnea test (HSAT) (e.g., severe insomnia); **OR**
- Environmental or personal factors that preclude the adequate acquisition and interpretation of data from HSAT; **OR**
- The patient or caregiver is unable to safely use the equipment for home sleep

- study due to dexterity, mobility, or cognitive function; **OR**
- A home sleep study was negative, inconclusive, or technically inadequate^{10,19}; **OR**
- Periodic leg movement disorder (PLMD) is considered because of complaints by the patient or an observer of repetitive limb movements during sleep and frequent awakenings, fragmented sleep, difficulty maintaining sleep, or excessive daytime sleepiness³; **OR**
- **ANY** of the following are suspected with planned Multiple Sleep Latency Testing (MSLT)⁷:
 - ◆ Narcolepsy with **ANY** of the following:
 - Cataplexy; **OR**
 - Excessive daytime sleepiness; **OR**
 - Hallucinations with the onset of sleep or awakening; **OR**
 - Disrupted nighttime sleep; **OR**
 - Sleep paralysis; **OR**
 - ◆ Central disorders of hypersomnia with **ANY** of the following:
 - Sleep inertia; **OR**
 - Unrefreshed sleep with adequate or long sleep time; **OR**
- An in-lab sleep study is needed when a home sleep study cannot be performed with **ANY** of the following²⁰:
 - ◆ Obesity hypoventilation syndrome (defined as a body mass index [BMI] greater than 30, daytime hypercapnia [partial pressure of carbon dioxide, PaCO₂, greater than 45 mm Hg without other causes such as kyphosis, myopathy, hypothyroidism, or lung disease]); **OR**
 - ◆ Awake daytime hypercapnia (partial pressure of carbon dioxide, PaCO₂, greater than or equal

to 45 mm Hg without other causes such as kyphosis, myopathy, hypothyroidism, or lung disease – serum bicarbonate greater than 28 is considered an alternative in the absence of PaCO₂ from arterial blood gases); **OR**

- Suspected central sleep apnea (CSA); **OR**
- Evaluation of parasomnias (e.g., undesirable or unpleasant occurrences during sleep, sleepwalking, sleep terrors, rapid eye movement, sleep behavior disorder²¹, history of repeated violent or injurious episodes during sleep) with **ANY** of the following^{3,14}:
 - ◆ The patient has a history of parasomniac episodes during sleep that result in harm to the patient or others; **OR**
 - ◆ To assist with the diagnosis of paroxysmal arousals or other sleep disruptions that are thought to be seizure-related when the initial clinical evaluation and results of a standard EEG are inconclusive; **OR**
 - ◆ To evaluate sleep-related behaviors that are violent or otherwise potentially injurious to the patient; **OR**
 - ◆ The patient has a sleep behavior suggestive of parasomnias that are unusual or atypical due to **ALL** of the following:
 - Age at onset; **AND**
 - Time, duration, or frequency of occurrence of the behavior; **AND**
 - Specifics of the particular motor patterns are in question (e.g. stereotypical, repetitive, or focal); **OR**
 - ◆ In situations with forensic considerations (e.g., if onset follows trauma or if the events themselves have been associated with personal injury); **OR**
 - ◆ When the presumed parasomnia or sleep-related epilepsy does not respond to conventional therapy; **OR**

- ◆ In cases of typical, uncomplicated, and non-injurious parasomnias when the diagnosis is not clearly delineated; **OR**
- The pediatric patient has suspected sleep apnea with an initial PSG test indicated by **ANY** of the following²²⁻²⁴:
 - ◆ Evaluation for obstructive sleep apnea (OSA) pre- or post-removal of enlarged tonsils or adenoids; **OR**
 - ◆ Down syndrome; **OR**
 - ◆ Chiari malformation; **OR**
 - ◆ Craniofacial malformation; **OR**
 - ◆ Neuromuscular disorder (e.g., Parkinson's disease, stroke with persistent neurological sequelae, Duchenne muscular dystrophy, multiple sclerosis with associated pulmonary disease, amyotrophic lateral sclerosis, myotonic dystrophy); **OR**
 - ◆ Skeletal dysplasia (e.g., achondroplasia); **OR**
 - ◆ Suspected periodic limb movement disorder²⁵; **OR**
 - ◆ Signs and symptoms of OSA with **ANY** of the following:
 - Snoring; **OR**
 - Daytime sleepiness; **OR**
 - Mouth breathing; **OR**
 - Nocturnal apnea; **OR**
 - Enuresis; **OR**
 - Pulmonary hypertension; **OR**
 - Nasal flaring or other signs of breathing difficulty; **OR**
 - Failure to thrive (weight less than 5th percentile for age); **OR**
 - Hyponasal speech; **OR**
 - Behavioral problems (e.g., hyperactivity, developmental delay, difficulties in school); **OR**

- An attended, full-night titration study including **ALL** of the following:
 - ◆ Unattended auto-titration with APAP or auto bi-level PAP is contraindicated; **AND**
 - ◆ OSA with **ANY** of the following:
 - AHI, RDI, or REI greater than or equal to 15 events per hour; **OR**
 - AHI, RDI, or REI greater than or equal to 5-14 events per hour with **ANY** of the following:
 - Excessive daytime sleepiness; **OR**
 - Insomnia; **OR**
 - Mood disorders (e.g., anxiety, depression); **OR**
 - Impaired cognition; **OR**
 - History of stroke; **OR**
 - Hypertension; **OR**
 - Ischemic heart disease; **AND**
 - ◆ **ANY** of the following:
 - The patient has **ANY** of the following co-morbid conditions:
 - Heart failure with New York Heart Association (NYHA) Classification III or IV or reduced ejection fraction less than or equal to 40%; **OR**
 - Cardiac arrhythmia(s) (acute, uncontrolled, or refractory) with documented symptoms; **OR**
 - Severe asthma with daily use of oral corticosteroids and/or immunomodulator biologics; **OR**
 - Pulmonary disease (e.g., moderate to severe COPD or interstitial lung disease) as diagnosed on pulmonary function studies (PFTs) and the patient requires chronic oxygen use; **OR**

- Obesity hypoventilation syndrome (OHS); **OR**
- Moderate to severe pulmonary hypertension; **OR**
- Neuromuscular/neurodegenerative disorders with restrictive disease or hypoventilation (e.g., amyotrophic lateral sclerosis [ALS], post-polio syndrome, myasthenia gravis, Guillian-Barré syndrome, polymyositis, kyphoscoliosis); **OR**
- Chronic opioid medication use; **OR**
- The patient has a second or associated sleep disorder other than OSA including **ANY** of the following:
 - Central nervous system disorders that increase the risk of CSA (e.g., Arnold Chiari malformation); **OR**
 - CSA or treatment-emergent sleep apnea; **OR**
 - Acute nocturnal seizures; **OR**
 - Narcolepsy or related symptoms following the evaluation and treatment of OSA in accordance with the patient's documented adherence to therapy; **OR**
 - Complex parasomnias that may include injurious, disruptive, or violent behavior (e.g., sleepwalking, REM behavior disorder); **OR**
 - Periodic limb movement disorder (PLMD); **OR**
- The patient failed a recent home APAP trial due to **ANY** of the following:
 - Auto bi-level therapy is contraindicated or was ineffective; **OR**

- PAP therapy was not tolerated after a trial of at least 1 month with no previous attended titration; **OR**
- Adequate objective adherence to therapy (greater than or equal to 4 hours per night for at least 70% of nights in a 30-day consecutive period as documented by APAP download) with **ANY** of the following:
 - ◆ Symptoms of residual excessive daytime sleepiness; **OR**
 - ◆ Residual AHI greater than or equal to 5 as evidenced by APAP download; **OR**
- Split-night protocol for strong pre-test suspicion of OSA with the initiation of treatment with positive pressure device including **ANY** of the following^{2,4,10}:
 - ◆ For initial testing with **ALL** of the following:
 - Sleep-disordered breathing as indicated by **ANY** of the following²⁶:
 - Loud snoring; **OR**
 - Gasping or choking during sleep; **OR**
 - Excessive daytime sleepiness; **OR**
 - Cognitive deficits (e.g., concentration, memory); **OR**
 - Morning headaches; **OR**
 - BMI greater than or equal to 30; **OR**
 - Large neck circumference (greater than 17 inches in men; greater than 16 inches in women); **OR**
 - Unexplained nocturnal reflux; **OR**
 - Sleep-related bruxism; **OR**
 - Erectile dysfunction; **OR**

- Apneas or hypoxemia during procedures requiring anesthesia;
AND
- **ANY** of the following:
 - A non-diagnostic HSAT performed no more than 1 year prior with **ANY** of the following:
 - ◆ An OSA diagnosis was not determined when there was a high pretest probability of OSA; **OR**
 - ◆ An effort was made to perform the test again if the original test was determined to be technically inadequate; **OR**
 - The patient has **ANY** of the following co-morbid conditions:
 - ◆ Heart failure with New York Heart Association (NYHA) Classification III or IV or reduced ejection fraction less than or equal to 40%; **OR**
 - ◆ Cardiac arrhythmia(s) (acute, uncontrolled, or refractory) with documented symptoms; **OR**
 - ◆ Severe asthma with daily use of oral corticosteroids and/or immunomodulator biologics; **OR**
 - ◆ Pulmonary disease (e.g., moderate to severe COPD or interstitial lung disease) as diagnosed on pulmonary function studies (PFTs) and the patient requires chronic oxygen use; **OR**

- ◆ Obesity hypoventilation syndrome (OHS); **OR**
- ◆ Moderate to severe pulmonary hypertension; **OR**
- ◆ Neuromuscular/neurodegenerative disorders with restrictive disease or hypoventilation (e.g., amyotrophic lateral sclerosis [ALS], post-polio syndrome, myasthenia gravis, Guillian-Barré syndrome, polymyositis, kyphoscoliosis); **OR**
- ◆ Chronic opioid medication use; **OR**
- The patient has a second or associated sleep disorder other than OSA including **ANY** of the following:
 - ◆ Central nervous system disorders that increase the risk of CSA (e.g., Arnold Chiari malformation); **OR**
 - ◆ CSA or treatment-emergent sleep apnea; **OR**
 - ◆ Acute nocturnal seizures; **OR**
 - ◆ Narcolepsy or related symptoms following the evaluation and treatment of OSA in accordance with the patient's documented adherence to therapy; **OR**
 - ◆ Complex parasomnias that may include injurious, disruptive, or violent behavior (e.g., sleepwalking, REM behavior disorder); **OR**

- ◆ Periodic limb movement disorder (PLMD); **OR**
- ◆ For repeat split-night testing with **ANY** of the following:
 - The patient has **ANY** of the following:
 - Symptoms of OSA that persist or recur despite PAP therapy; **OR**
 - A change in weight (gain or loss greater than or equal to 10% of total body weight) when OSA symptoms have worsened or improved and re-evaluation is required to determine needed therapy modifications; **OR**
 - A change in cardiovascular status (e.g., stroke, arrhythmia, uncontrolled hypertension, hospitalization for heart failure; **OR**
 - To re-evaluate the indicators of OSA after **ANY** of the following:
 - Adenoidectomy; **OR**
 - Tonsillectomy; **OR**
 - Maxillomandibular advancement surgery (MMA); **OR**
 - Uvulopalatoplasty (UPPP); **OR**
 - Other surgery related to the upper airway; **OR**
 - A non-diagnostic HSAT performed no more than 1 year prior with **ANY** of the following:
 - An OSA diagnosis was not determined when there was a high pretest probability of OSA; **OR**
 - An effort was made to perform the test again if the original test was determined to be technically inadequate; **OR**

- The patient requires evaluation to begin using a fabricated oral mandibular advancement appliance (OAT); **OR**
- Repeat PSG test, as indicated by **ANY** of the following:
 - ◆ Confirmation of the efficacy of prescribed therapy is needed (e.g., oral appliance, postoperative assessment of response to intervention)⁴; **OR**
 - ◆ To ascertain whether CPAP is still needed at the previously titrated pressure if the patient experiences substantial weight gain or weight loss on CPAP for the treatment of sleep-related breathing disorders⁴; **OR**
 - ◆ The patient shows insufficient clinical response or symptoms return despite a good initial response to CPAP⁴; **OR**
 - ◆ Assessing treatment response after upper airway surgical procedures, or initial treatment with oral appliances for **ANY** of the following^{4,27-28}:
 - Initial treatment with oral appliances (pre-implantation or re-evaluation of known OSA) with **ANY** of the following:
 - PAP failure or PAP intolerance with BMI less than or equal to 35 with no recent sleep study; **OR**
 - A significant change in weight and/or symptoms; **OR**
 - After upper airway surgical procedures (post-implantation) with **ANY** of the following:
 - Initial PSG titration; **OR**
 - PSG titration previously performed with insufficient clinical response, weight gain and/or return of symptoms; **OR**
- Multiple Sleep Latency Test (MSLT) performed in a sleep laboratory for **ANY** of the following⁷:

- ◆ For initial MSLT with **ALL** of the following:
 - Evaluation of presence or treatment response for features of **ANY** of the following:
 - Central disorders of hypersomnia;
 - OR**
 - Narcolepsy (including, cataplexy, EDS, sleep paralysis, hypersomnia);
 - AND**
 - Testing consists of 5 episodes of 20-minute nap trials at 2-hour intervals, measuring the onset of sleep and rapid eye movement (REM) sleep, immediately following a negative PSG when narcolepsy is suspected; **OR**
- ◆ Repeat MSLT may be required if initial results are indeterminate or negative when narcolepsy is suspected; **OR**
- Maintenance of Wakefulness Tests (MWT) performed in a sleep laboratory are considered appropriate with **ANY** of the following:
 - ◆ For initial MWT with **ALL** of the following²⁹:
 - To assess an individual's ability to remain awake when his or her inability to remain awake constitutes a public or personal safety issue; **AND**
 - To assess response to treatment and the patient has excessive daytime sleepiness;
 - OR**
 - ◆ Repeat MWT may be required if initial results are indeterminate or negative when symptoms persist; **AND**
- **ALL** of the following:
 - Sleep studies are performed in **ANY** of the following settings:
 - ◆ Hospital³; **OR**
 - ◆ Sleep laboratory (hospital and freestanding facilities including sleep clinics that are part of

a physician's office, and all other non-hospital-based facilities where sleep studies are performed) that is accredited by **ANY** of the following^{3,14,24}:

- American Academy of Sleep Medicine (AASM); **OR**
- Accreditation Commission for Health Care (ACHC); **OR**
- The Joint Commission; **OR**
- ◆ Independent Diagnostic Treatment Facility (IDTF) that is supervised by a physician (MD/DO) trained in analyzing and interpreting the recordings, and should be attended by a trained technologist³; **AND**
- Sleep studies are performed by **ANY** of the following providers:
 - ◆ The physician performing the service must meet **ANY** of the following⁴:
 - Diplomate of the American Board of Sleep Medicine (ABSM); **OR**
 - Sleep certification issued by **ANY** of the following Boards:
 - American Board of Internal Medicine (ABIM); **OR**
 - American Board of Family Medicine (ABFM); **OR**
 - American Board of Pediatrics (ABP); **OR**
 - American Board of Psychiatry and Neurology (ABPN); **OR**
 - American Board of Otolaryngology (ABOto); **OR**
 - American Osteopathic Board of Neurology and Psychiatry (AOBNP); **OR**
 - American Osteopathic Board of Family Medicine, (AOBFP); **OR**

- American Osteopathic Board of Internal Medicine, (AOBIM); **OR**
 - American Osteopathic Board of Ophthalmology and Otorhinolaryngology (AOBOO); **OR**
- Be an active physician staff member of a credentialed sleep center or laboratory that has active physician staff members meeting the physician credential criteria above; **OR**
- ◆ The technician performing the service must meet the requirements of **ANY** of the following⁴:
 - American Board of Sleep Medicine (ABSM) (e.g., Registered Sleep Technologist [RST]); **OR**
 - Board of Registered Polysomnographic Technologists (BRPT) (e.g., Registered Polysomnographic Technologist [RPSGT]); **OR**
 - National Board for Respiratory Care (NBRC) (e.g., Certified Pulmonary Function Technologist [CPFT], Registered Pulmonary Function Technologist [RPFT], Certified Respiratory Therapist [CRT], Registered Respiratory Therapist [RRT]).

Non-Indications

→ A **home sleep apnea study** is not considered appropriate if **ANY** of the following is **TRUE**¹⁰:

- ◆ The request is for a pediatric home-based sleep study²⁴; **OR**
- ◆ The patient is an adult with testing being performed for **ANY** of the following:
 - For the diagnosis of **ANY** of the following:
 - Circadian rhythm sleep disorders; **OR**
 - Chronic lung disease; **OR**
 - Preoperative evaluation for laser-assisted uvulopalatopharyngoplasty without clinical evidence of suspicion of OSA; **OR**

- **ANY** of the following:
 - Seizure disorder⁴; **OR**
 - Significant cardiopulmonary disease (forced expiratory volume in one second [FEV₁] % predicted of less than 70 or heart failure with left ventricular ejection fraction (LVEF) less than 45%)^{8,17-18}; **OR**
 - Potential respiratory muscle weakness due to neuromuscular conditions³⁰; **OR**
 - Periodic limb movement disorder or restless leg syndrome without suspicion of OSA³⁰; **OR**
 - Hypoventilation syndrome (awake hypoventilation or suspected sleep-related hypoventilation); **OR**
 - Acute opioid use (medication not normally taken by the patient)³¹; **OR**
 - Chronic opioid medication use³¹; **OR**
 - History of stroke²⁶; **OR**
 - Suspected sleep disorder other than OSA (e.g., CSA, parasomnia, narcolepsy)^{26,30}; **OR**
 - General screening of an asymptomatic patient⁴; **OR**
 - To confirm typical, uncomplicated, and non-injurious parasomnias when the diagnosis is clearly delineated⁴; **OR**
 - Actigraphy used for the diagnosis of sleep disorders; **OR**
 - The patient or caregiver is unable to manage equipment; **OR**
 - The patient is asymptomatic and testing is for general screening.

Level of Care Criteria

Inpatient or Outpatient

Procedure Codes (CPT/HCPCS)

CPT/HCPCS Code	Code Description
95782	Polysomnography; younger than 6 years, sleep staging with 4 or more additional parameters of sleep, attended by a technologist

95783	Polysomnography; younger than 6 years, sleep staging with 4 or more additional parameters of sleep, with initiation of continuous positive airway pressure therapy or bi-level ventilation, attended by a technologist
95800	Sleep study, unattended, simultaneous recording; heart rate, oxygen saturation, respiratory analysis (eg, by airflow or peripheral arterial tone), and sleep time
95801	Sleep study, unattended, simultaneous recording; minimum of heart rate, oxygen saturation, and respiratory analysis (eg, by airflow or peripheral arterial tone)
95803	Actigraphy testing, recording, analysis, interpretation, and report (minimum of 72 hours to 14 consecutive days of recording)
95805	Multiple sleep latency or maintenance of wakefulness testing, recording, analysis and interpretation of physiological measurements of sleep during multiple trials to assess sleepiness
95806	Sleep study, unattended, simultaneous recording of, heart rate, oxygen saturation, respiratory airflow, and respiratory effort (eg, thoracoabdominal movement)
95807	Sleep study, simultaneous recording of ventilation, respiratory effort, ECG or heart rate, and oxygen saturation, attended by a technologist
95808	Polysomnography; any age, sleep staging with 1-3 additional parameters of sleep, attended by a technologist
95810	Polysomnography; age 6 years or older, sleep staging with 4 or more additional parameters of

	sleep, attended by a technologist
95811	Polysomnography; age 6 years or older, sleep staging with 4 or more additional parameters of sleep, with initiation of continuous positive airway pressure therapy or bi-level ventilation, attended by a technologist

Disclaimer: S Codes are non-covered per CMS guidelines due to their experimental or investigational nature.

Medical Evidence

Queisi et al. (2024) performed a retrospective review to compare polysomnographic (PSG) sleep parameters for people with multiple sclerosis (PwMS). The study evaluates a large single cohort from a single center versus existing published standards. A total of 299 PwMS were evaluated at a facility for polysomnography (PSG). Data included total sleep time (TST), sleep efficiency (SE), sleep onset latency (SOL), relative REM latency, total apnea-hypopnea indices (AHI), spontaneous arousal indices (AI), total periodic leg movements indices (PLMI), and sleep architecture metrics (e.g., the percentage spent in stages N1/N2, N3, and REM). Compared to normative data, PwMS had an average of 85.9 minutes shorter TST, 27.3 minutes longer SOL, 62.1 minutes longer REM latency, 10.7% lower SE, 16.4% more N1/N2, and 11.4% less N3. The population demonstrated a high prevalence of obstructive sleep apnea (OSA; 60.7%); the mean AHI was also higher by 11.1 events per hour. Fatigue is a primary symptom among PwMS. The study highlighted the need for established parameters for PwMS.³²

The American Heart Association (AHA) (2021) issued a scientific statement regarding obstructive sleep apnea (OSA) and cardiovascular disease. Testing is recommended for cardiovascular conditions, including resistant hypertension, pulmonary hypertension, recurrent atrial fibrillation, heart failure, stroke, and for survivors of sudden cardiac death. Follow-up testing is recommended to determine the effectiveness of treatment.⁸

The American Academy of Sleep Medicine (AASM) has published a guideline and position statements related to testing for OSA and other sleep disorders, including the following:

- Das et al. (2022) developed a position statement for AASM focusing on enhancing public health and safety by diagnosing and treating OSA in those in the transportation industry. Recommendations have included mandatory testing and treatment for OSA for rail and highway personnel in safety-sensitive positions.³³
- Kapur et al. (2017) published the *Clinical Practice Guideline for Diagnostic Testing for Adult Sleep Apnea*, with clinical recommendations using the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) system. A strong

recommendation was made for facility-based testing rather than home testing for patients with significant cardiorespiratory disease, neuromuscular conditions with respiratory muscle weakness, a history of stroke, severe insomnia, or chronic opioid use.¹⁰

- Kirk et al. (2017) published a position statement regarding home sleep apnea testing for diagnosing OSA in children. The authors concluded that home testing is not recommended in children less than 18 years of age. Limited evidence exists comparing attended PSG to home testing.²⁴

Khan et al. (2015) systematically reviewed the peer-reviewed literature regarding central disorders of hypersomnolence. They state that there have been significant advances in recent years, particularly in the diagnosis and management of narcolepsy type 1. A 24-hour PSG is important in the diagnosis of central disorders of hypersomnia.⁹

References

1. Centers for Medicare and Medicaid Services (CMS). National coverage determination (NCD): Continuous positive airway pressure (CPAP) therapy for obstructive sleep apnea (OSA) (240.4). Effective Date March 13, 2008.
<https://www.cms.gov/medicare-coverage-database/search.aspx>
2. Centers for Medicare and Medicaid Services (CMS). National coverage determination (NCD): Sleep testing for obstructive sleep apnea (OSA) (240.4.1). Effective Date March 3, 2009.
<https://www.cms.gov/medicare-coverage-database/search.aspx>
3. Centers for Medicare and Medicaid Services (CMS). Local coverage determination (LCD): Polysomnography (L36593). Revision Effective Date March 20, 2025. <https://www.cms.gov/medicare-coverage-database/view/lcd.aspx>
4. Centers for Medicare and Medicaid Services (CMS). Local coverage determination (LCD): Polysomnography and other sleep studies (L36902). Revision Effective Date February 27, 2025.
<https://www.cms.gov/medicare-coverage-database/search.aspx>
5. Centers for Medicare and Medicaid Services (CMS). Billing and coding article: Polysomnography (A56995). Revision Effective Date March 7, 2024. <https://www.cms.gov/medicare-coverage-database/search.aspx>
6. Centers for Medicare and Medicaid Services (CMS). Billing and coding article: Polysomnography and other sleep studies (A57049). Revision Effective Date March 7, 2024. <https://www.cms.gov/medicare-coverage-database/search.aspx>
7. Krahn LE, Arand DL, Avidan AY, et al. Recommended protocols for the Multiple Sleep Latency Test and Maintenance of Wakefulness Test in adults: Guidance from the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2021 Dec 1;17(12):2489–2498. doi:10.5664/jcsm.9620
8. Yeghiazarians Y, Jneid H, Tietjens JR, et al. Obstructive sleep apnea and cardiovascular disease: A scientific statement from the American Heart Association. *Circulation*. 2021 Jul 20;144(3):e56–e67. doi:10.1161/CIR.0000000000000988
9. Khan Z, Trotti LM. Central disorders of hypersomnolence: Focus on the narcolepsies and idiopathic hypersomnia. *Chest*. 2015 Jul;148(1):262–273. doi:10.1378/chest.14-1304

10. Kapur VK, Auckley DH, Chowdhuri S, et al. Clinical practice guideline for diagnostic testing for adult obstructive sleep apnea: An American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2017 Mar 15;13(3):479–504. doi:10.5664/jcsm.6506
11. Rosen IM, Kirsch DB, Carden KA, et al. Clinical use of a home sleep apnea test: An updated American Academy of Sleep Medicine position statement. *J Clin Sleep Med*. 2018 Dec 15;14(12):2075–2077. doi:10.5664/jcsm.7540
12. Chung F, Abdullah HR, Liao P. STOP-Bang questionnaire: A practical approach to screen for obstructive sleep apnea. *Chest*. 2016 Mar;149(3):631–8. doi:10.1378/chest.15-0903
13. Pivetta B, Chen L, Nagappa M, et al. Use and performance of the STOP-Bang questionnaire for obstructive sleep apnea screening across geographic regions: A systematic review and meta-analysis. *JAMA Netw Open*. 2021 Mar 1;4(3):e211009. doi:10.1001/jamanetworkopen.2021.1009
14. Centers for Medicare and Medicaid Services (CMS). Local coverage determination (LCD): Polysomnography and sleep testing (L33405). Revision Effective Date July 1, 2020. <https://www.cms.gov/medicare-coverage-database/search.aspx>
15. Corral J, Sánchez-Quiroga MÁ, Carmona-Bernal C, et al. Conventional polysomnography is not necessary for the management of most patients with suspected obstructive sleep apnea – noninferiority, randomized controlled trial. *Am J Respir Crit Care Med*. 2017 Nov 1;196(9):1181–1190. doi:10.1164/rccm.201612-2497OC
16. Masa JF, Corral J, Pereira R, et al. Effectiveness of home respiratory polygraphy for the diagnosis of sleep apnoea and hypopnoea syndrome. *Thorax*. 2011 Jul;66(7):567–73. doi:10.1136/thx.2010.152272
17. Pellegrino R, Viegi G, Brusasco V, et al. Interpretative strategies for lung function tests. *Eur Respir J*. 2005 Nov;26(5):948–68. doi:10.1183/09031936.05.00035205
18. Zhao YY, Blackwell T, Ensrud KE, et al. Sleep apnea and obstructive airway disease in older men: Outcomes of sleep disorders in older men study. *Sleep*. 2016;39(7):1343–1351. Published 2016 Jul 1. doi:10.5665/sleep.5960
19. United States Department of Veterans Affairs (VA). VA/DoD clinical practice guideline for the management of chronic insomnia disorder

- and obstructive sleep apnea. Updated 2019.
<https://www.healthquality.va.gov/guidelines/CD/insomnia/index.asp>
20. Mokhlesi B, Masa JF, Brozek JL, et al. Evaluation and management of obesity hypoventilation syndrome: An official American Thoracic Society clinical practice guideline. *Am J Respir Crit Care Med*. 2019 Aug 1;200(3):e6–e24. doi:10.1164/rccm.201905-1071ST
 21. Aurora RN, Zak RS, Maganti RK, et al. Best practice guide for the treatment of REM sleep behavior disorder (RBD). *J Clin Sleep Med*. 2010 Feb 15;6(1):85–95. PMID: 20191945.
 22. Aurora RN, Zak RS, Karippot A, et al. Practice parameters for the respiratory indications for polysomnography in children. *Sleep*. 2011 Mar 1;34(3):379–88. doi:10.1093/sleep/34.3.379
 23. Aurora RN, Lamm CI, Zak RS, et al. Practice parameters for the non-respiratory indications for polysomnography and multiple sleep latency testing for children. *Sleep*. 2012 Nov 1;35(11):1467–73. doi:10.5665/sleep.2190
 24. Kirk V, Baughn J, D’Andrea L, et al. American Academy of Sleep Medicine position paper for the use of a home sleep apnea test for the diagnosis of OSA in children. *J Clin Sleep Med*. 2017 Oct 15;13(10):1199–1203. doi:10.5664/jcsm.6772
 25. Beck SE, Marcus CL. Pediatric polysomnography. *Sleep Med Clin*. 2009 Sep;4(3):393–406. doi:10.1016/j.jsmc.2009.04.007
 26. American Association of Sleep Technologists (AASST). Technical guideline: Home sleep apnea testing (HSAT). Published December 2020. https://www.aastweb.org/Portals/0/Docs/Resources/Guidelines/HSAT%20Technical%20Guideline%202020_FINAL_New%20Template-1.pdf
 27. Baptista P, Costantino A, Moffa A, et al. Hypoglossal nerve stimulation in the treatment of obstructive sleep apnea: Patient selection and new perspectives. *Nat Sci Sleep*. 2020;12:151–159. doi:10.2147/nss.S221542
 28. Steffen A, König I, Baptista P, et al. Home sleep testing to direct upper airway stimulation therapy optimization for sleep apnea. *Laryngoscope*. Apr 2021; 131: E1375–e1379. doi:10.1002/lary.29043
 29. Littner MR, Kushida C, Wise M, et al. Practice parameters for clinical use of the multiple sleep latency test and the maintenance of wakefulness test. *Sleep*. 2005 Jan;28(1):113–21. doi:10.1093/sleep/28.1.113
 30. American Board of Internal Medicine (ABIM), American Academy of Sleep Medicine (AASM). Five things physicians and patients should question. Published December 2, 2014.

<https://aasm.org/wp-content/uploads/2017/11/choosingwisely-sleepmedicine.pdf>

31. Rosen IM, Aurora RN, Kirsch DB, et al. Chronic opioid therapy and sleep: An American Academy of Sleep Medicine position statement. *J Clin Sleep Med*. 2019 Nov 15;15(11):1671-1673. doi:10.5664/jcsm.8062
32. Queisi M, Cipriani V, Golan D, et al. Polysomnography parameters in a large cohort of people with multiple sclerosis. *Sleep Med*. 2024 Sep;121:236-240. doi:10.1016/j.sleep.2024.07.009
33. Das AM, Chang JL, Berneking M, et al. Enhancing public health and safety by diagnosing and treating obstructive sleep apnea in the transportation industry: An American Academy of Sleep Medicine position statement. *J Clin Sleep Med*. 2022 Oct 1;18(10):2467-2470. doi:10.5664/jcsm.9670

Clinical Guideline Revision

History/Information

Original Date: February 6, 2025

Review History

Version 2	03/13/2025	<ul style="list-style-type: none"> Added criteria from CMS LCD L36593 for sleep study/PSG. Updated the home-based, non-attended sleep study indication with STOP-BANG criteria. An indication was also added – “test is required within 24 months of the first consultation for hypoglossal nerve stimulation (HGNS) implant” (Corral et al, 2017 and Masa et al, 2011). For the suspicion indications and use of Multiple Sleep Latency Testing (MSLT), added “central disorders of hypersomnia with ANY of the following: <ul style="list-style-type: none"> Sleep inertia; OR Unrefreshed sleep with adequate or long sleep time; OR” Separated the indications for MSLT performed in a sleep laboratory and Maintenance of Wakefulness Tests (MWT). Added indication for full-night titration study. Rewrote the indication for the split-night protocol for strong pre-test suspicion of OSA. Per CMS, included requirements for place of service and accreditation of the provider.
-----------	------------	--

Version 3	04/17/2025	<p>Boolean logic updates made:</p> <ul style="list-style-type: none"> • On p. 6, previously, the first diamond bullet was “ANY of the following” – deleted and moved the bullets over one level to the left. • On p. 7 – bullet for “Transcutaneous CO2, esophageal pH, penile tumescence, and bipolar EEG” (<i>changed from an AND to an OR</i>) • On p. 7 – the main bullet for “home-based testing” was changed from an ANY to an ALL • On p. 8, under the bullet for “The patient is an adult with a high pretest probability of moderate to severe obstructive sleep apnea (OSA) with ALL of the following” – indented 2 bullets (“sleep study is recorded” and “sleep study is staged”) to the right to fix logic. • On p. 15, the indication for “split-night protocol” was updated from ALL to ANY (previously, the patient was required to meet both the initial and repeat criteria together). • On p. 20, changed the OR to AND – see the indication for “Maintenance of Wakefulness Tests (MWT)”, under the first sub-bullet for initial MWT (“to assess an individual’s ability to remain awake when his or her inability to remain awake constitutes a public or personal safety issue”).
Version 3.1	04/21/2025	<ul style="list-style-type: none"> • Updated policy per CMS revisions for 3/20/2025 • Updated Effective Date

		<ul style="list-style-type: none">• Updated Links and Bookmarks
--	--	---