



Cohere Medical Policy – Knee Arthroscopy

Clinical Policy for Medical Necessity Review

Version: 8

Cohere Health UMC Approval Date: September 11, 2025

Last Annual Review: April 17, 2025

Revision: September 11, 2025

Next Annual Review: April 17, 2026

This policy was reviewed by the American Association of Orthopaedic Surgeons (AAOS) prior to publication.

Important Notices

Notices & Disclaimers:

GUIDELINES ARE SOLELY FOR COHERE'S USE IN PERFORMING MEDICAL NECESSITY REVIEWS AND ARE NOT INTENDED TO INFORM OR ALTER CLINICAL DECISION-MAKING OF END USERS.

Cohere Health, Inc. ("**Cohere**") has published these clinical guidelines to determine the medical necessity of services (the "**Guidelines**") for informational purposes only, and solely for use by Cohere's authorized "**End Users**". These Guidelines (and any attachments or linked third-party content) are not intended to be a substitute for medical advice, diagnosis, or treatment directed by an appropriately licensed healthcare professional. These Guidelines are not in any way intended to support clinical decision-making of any kind; their sole purpose and intended use is to summarize certain criteria Cohere may use when reviewing the medical necessity of any service requests submitted to Cohere by End Users. Always seek the advice of a qualified healthcare professional regarding any medical questions, treatment decisions, or other clinical guidance. The Guidelines, including any attachments or linked content, are subject to change at any time without notice.

© 2025 Cohere Health, Inc. All Rights Reserved.

Other Notices:

HCPCS® and CPT® copyright 2025 American Medical Association. All rights reserved.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

HCPCS and CPT are registered trademarks of the American Medical Association.

Policy Information:

Specialty Area: Musculoskeletal Care

Policy Name: Knee Arthroscopy

Type: ☒ Adult (18+ yo) | ☒ Pediatric (0-17 yo)

Table of Contents

Important Notices	2
Medical Necessity Criteria	4
Service: Knee Arthroscopy	4
Description	5
Medical Necessity Criteria	5
Indications	5
Non-Indications	10
Definitions	10
Level of Care Criteria	10
Procedure Codes (CPT/HCPCS)	10
Medical Evidence	14
References	17
Policy Revision History/Information	23

Medical Necessity Criteria

Service: Knee Arthroscopy

Cohere Health takes an evidence-based approach to reviewing imaging and procedure requests, meaning that sufficient clinical information must be provided at the time of submission to determine medical necessity.

Documentation must include a recent and detailed history, physical examination related to the onset or change in symptoms, relevant lab results, prior imaging, and details of previous treatments. Advanced imaging or procedures should be requested after a clinical evaluation by the treating provider, which may include referral to a specialist.

- When a specific clinical indication is not explicitly addressed in the Cohere Health medical policy, medical necessity will be determined based on established clinical best practices, as supported by evidence-based literature, peer-reviewed sources, professional society guidelines, and state or national recommendations, unless otherwise directed by the health plan.
- Requests submitted without clinical documentation, or those that do not align with the provided clinical information—such as mismatched procedure, laterality, body part, or CPT code—may be denied for lack of medical necessity due to insufficient or inconsistent clinical information.
- When there are multiple diagnostic or therapeutic procedures requested simultaneously or within the past three months, each will be reviewed independently. Clinical documentation must clearly justify all of the following:
 - The medical necessity of each individual request
 - Why prior imaging or procedures were inconclusive, or why additional/follow-up studies are needed
 - How the results will impact patient management or treatment decisions
- Requests involving adjacent or contiguous body parts may be considered not medically necessary if the documentation demonstrates that the patient's primary symptoms can be adequately assessed with a single study or procedure.

Description

Knee arthroscopy is a minimally invasive method of visualizing and treating the knee joint. It is the standard treatment for the removal of inflamed synovial tissue, loose fragments of bone or cartilage, and resecting the damaged meniscus. The procedure may also be performed to trim or reconstruct damaged articular cartilage, as well as to reconstruct a torn anterior cruciate ligament (ACL) or posterior cruciate ligament (PCL). Knee arthroscopy is additionally performed to treat knee sepsis and issues related to the patella, such as chondromalacia patella. Osteochondritis dissecans, a condition in which a portion of bone and the cartilage covering it begins to separate from the joint, may be treated and evaluated arthroscopically. As an alternative to more invasive open procedures, knee arthroscopy is well-tolerated, safe, and leads to improved quality of life and function.¹⁻⁷

Medical Necessity Criteria

Indications

Knee arthroscopy is considered appropriate if **ANY** of the following is **TRUE**:

- **Intraarticular joint pathology evaluation, lateral retinacular release, or synovectomy/plica resection** when **ALL** of the following are **TRUE**⁸⁻¹²:
 - **ANY** of the following
 - Chronic knee pain and **ALL** of the following are **TRUE**:
 - Unknown etiology of symptoms; **AND**
 - Imaging does not yield definitive results; **AND**
 - Diagnostic arthrocentesis with synovial fluid analysis when **ANY** of the following is **TRUE**:
 - Nondiagnostic; **OR**
 - Not indicated; **OR**
 - Symptoms of locking, catching, and giving way; **OR**
 - A loose body noted on imaging; **OR**
 - Patellofemoral instability; **OR**
 - Abnormal patellar tracking; **OR**
 - Patellar compression syndrome; **OR**
 - Inflammatory (e.g., rheumatoid arthritis, psoriatic arthritis, Lyme arthritis) arthritis; **OR**
 - Benign tumor/neoplastic disorders (osteochondromatosis, tenosynovial giant cell tumor [pigmented villonodular synovitis – PVNS], synovial hemangioma, and recurrent hemarthrosis)¹³; **OR**

- Recurrent effusion¹⁴; **OR**
- Plica syndrome¹⁴; **OR**
- Hemophilic joint disease¹⁵; **OR**
- Lipoma arborescens (synovial lipomatosis)¹⁴; **AND**
- **ANY** of the following:
 - Conservative management does not apply – tenosynovial giant cell tumor (pigmented villonodular synovitis [PVNS]) or other pathologic synovial processes; **OR**
 - Failure of conservative management for greater than 3 months, including **ALL** of the following:
 - Anti-inflammatory medications, non-opioid analgesics, or prescription medications (e.g., oral steroids, neuropathic pain medications) if not contraindicated; **AND**
 - Physical therapy or physician-directed exercise program; **AND**
 - Bracing if medically appropriate; **AND**
 - **ANY** of the following:
 - Corticosteroid injection if medically appropriate; **OR**
 - Documentation that corticosteroid injection is contraindicated;
- OR**
- **Debridement, lysis, drainage, or lavage** when **ALL** of the following are **TRUE**⁸:
 - **ANY** of the following:
 - Infected joint (septic arthritis)^{9,10}; **OR**
 - A foreign body on imaging (e.g., hardware); **AND**
 - The patient has **ANY** of the following conditions:
 - Nonoperative care does not apply (infected joint; symptoms of locking, catching, giving way); **OR**
 - Focal articular cartilage lesion (less than 4 cm²) with **ALL** of the following^{2,16,17}:
 - Symptoms that are related to chondral injury; **AND**
 - Absence of advanced osteoarthritis; **AND**
 - Nonoperative care (e.g., anti-inflammatory medications, analgesics, physical therapy, bracing) has been attempted and failed; **OR**
 - Arthrofibrosis (e.g., after prior surgical procedure or trauma), as indicated by **ALL** of the following⁷:
 - Loss of range of motion; **AND**

- Nonoperative care (e.g., physical therapy, manipulation under anesthesia) has been attempted and failed for at least 6 weeks; **OR**
- **Ligament injury (ACL, PCL, LCL)** when **ALL** of the following are **TRUE**^{8,18-22}:
 - Absence of advanced arthritis; **AND**
 - Limited activities of daily living (ADLs) due to pain and instability; **AND**
 - Presence of **ANY** of the following:
 - Anterior cruciate ligament (ACL) tear; **OR**
 - Posterior cruciate ligament (PCL) tear; **OR**
 - Lateral collateral (fibular) ligament (LCL) tear; **OR**
 - Posterolateral corner (PLC) of the knee injury; **AND**
 - **ANY** of the following:
 - Decrease in swelling with improvement in range of motion since initial injury; **OR**
 - Clinically urgent injury for which ROM restoration would not be feasible prior to surgery; **AND**
 - Treatment is indicated for **ANY** of the following reasons:
 - Acute tear or injury; **OR**
 - Chronic injury associated with meniscal tear; **OR**
 - Concurrent with injury of **ANY** of the following:
 - Concomitant MCL/PCL/ACL/PLC/LCL injury; **OR**
 - Concomitant avulsion fracture; **OR**
 - Posterolateral corner of the knee is injured; **OR**
 - Tibial displacement of more than 8mm is demonstrated on stress radiographs; **OR**
 - Anterolateral ligament injury in conjunction with ACL injury; **OR**
 - Chronic injury with **ANY** of the following:
 - Failure of conservative management for greater than 3 months, including **ALL** of the following:
 - Anti-inflammatory medications, non-opioid analgesics, or prescription medications (e.g., oral steroids, neuropathic pain medications) if not contraindicated; **AND**
 - Physical therapy or physician-directed exercise program; **AND**
 - **ANY** of the following:
 - Corticosteroid injection if medically appropriate; **OR**
 - Documentation that corticosteroid injection is contraindicated; **OR**

- Persistent instability such that surgical treatment recommended; **OR**
- **Treatment of osteochondral defect (e.g., osteochondritis dissecans)** when **ALL** of the following are **TRUE**^{23,24}:
 - Plain radiograph or advanced imaging demonstrates an osteochondral defect; **AND**
 - Patient has **ANY** of the following:
 - Symptoms attributable to the osteochondral defect; **OR**
 - Acute traumatic injury resulting in osteochondral defect; **OR**
 - Unstable lesion/unstable defect on imaging; **OR**
- **Autologous chondrocyte implantation (ACI)** when **ALL** of the following are **TRUE**^{3,8}:
 - The patient has knee symptoms (e.g. pain, swelling, mechanical) or functional compromise; **AND**
 - Chondral lesion between 2 cm and 4 cm; **AND**
 - 12 years or older; **AND**
 - **ANY** of the following:
 - Underlying malalignment will be addressed with osteotomy in the same procedure; **OR**
 - Patient does not have concurrent malalignment; **OR**
- **Treatment of torn meniscus** when **ANY** of the following is **TRUE**^{1,25-33}:
 - **Partial meniscectomy** when **ANY** of the following is **TRUE**:
 - The tear is an acute tear due to injury or trauma, or a recurrent or failed prior repair, and **ALL** of the following are **TRUE**:
 - Advanced imaging demonstrates a meniscal tear; **AND**
 - Acute, painful knee with exam consistent with MRI finding of tear; **AND**
 - Persistent mechanical symptoms; **OR**
 - The tear is a chronic (degenerative) tear and **ALL** of the following are **TRUE**:
 - Advanced imaging demonstrates a meniscal tear; **AND**
 - Persistent mechanical symptoms; **AND**
 - Knee pain with exam consistent with MRI finding of meniscal tear; **AND**
 - Failure of conservative management for greater than 3 months with degenerative tears and minimal osteoarthritis, including **ALL** of the following:

- Anti-inflammatory medications, non-opioid analgesics, or prescription medications (e.g., oral steroids, neuropathic pain medications) if not contraindicated; **AND**
- Physical therapy or physician-directed exercise program; **AND**
- Activity modification; **AND**
- **ANY** of the following:
 - Corticosteroid injection if medically appropriate; **OR**
 - Documentation that corticosteroid injection is contraindicated; **OR**
- **Meniscus repair** when **ALL** of the following are **TRUE**:
 - The patient has mechanical symptoms following an acute injury; **AND**
 - The patient has a meniscal tear on advanced imaging findings or a failed prior repair; **OR**
- **Meniscus allograft transplantation** is considered appropriate if **ALL** of the following are **TRUE**⁸:
 - Significant portion of the meniscus is absent due to either prior meniscectomy or prior injury; **AND**
 - Persistent pain or impaired function with activity; **AND**
 - Age 50 years or younger; **AND**
 - Symptoms are localized to the tibiofemoral compartment; **AND**
 - No evidence of advanced arthrosis; **AND**
 - Over 50% joint space remaining; **AND**
 - **ANY** of the following:
 - Underlying malalignment will be addressed with osteotomy in the same procedure; **OR**
 - Patient does not have concurrent malalignment; **OR**
- **Arthroscopic management of fractures** when **ANY** of the following are **TRUE**⁸:
 - Fracture of the anterior intercondylar eminence of the tibia; **OR**
 - Other fracture pattern that would benefit from arthroscopically-assisted fracture reduction.

Non-Indications

Knee Arthroscopy is not considered appropriate if **ANY** of the following is **TRUE**:

- Osteoarthritis of the knee that is moderate or severe (or KL [Kellgren-Lawrence] grade III or grade IV)^{25,27,30}; **OR**
- Active infection at the surgical site (unless arthroscopy is indicated for surgical management of infection); **OR**
- The procedure is a meniscus allograft transplantation for **ANY** of the following:
 - Inflammatory arthropathy; **OR**
 - Malalignment or instability that is not being corrected at the time of the meniscal allograft surgery; **OR**
 - Irreparable/untreatable chondral damage.

Definitions

Advanced osteoarthritis is defined as complete or near-complete loss of joint space or moderate/severe KL grade arthritis^{1,25}:

- **Grade 3 (moderate)**: Moderate multiple osteophytes, definite narrowing of joint space, some sclerosis, and possible deformity of bone ends.
- **Grade 4 (severe)**: Large osteophytes, marked narrowing of joint space, severe sclerosis, and definite deformity of bone ends.

Level of Care Criteria

Inpatient or Outpatient

Procedure Codes (CPT/HCPCS)

CPT/HCPCS Code	Code Description
27405	Repair, primary, torn ligament and/or capsule, knee; collateral
27407	Repair, primary, torn ligament and/or capsule, knee; cruciate
27409	Repair, primary, torn ligament and/or capsule, knee; collateral and cruciate ligaments
27412	Autologous chondrocyte implantation, knee
27427	Ligamentous reconstruction (augmentation), knee; extra-articular
27599	Unlisted procedure, femur or knee
29850	Arthroscopically aided treatment of intercondylar spine(s) and/or tuberosity fracture(s) of the knee, with or without manipulation; without internal or external fixation (includes arthroscopy)
29851	Arthroscopically aided treatment of intercondylar spine(s) and/or tuberosity fracture(s) of the knee, with or without manipulation; with internal or external fixation (includes arthroscopy)
29855	Arthroscopically aided treatment of tibial fracture, proximal (plateau); unicondylar, includes internal fixation, when performed (includes arthroscopy)
29856	Arthroscopically aided treatment of tibial fracture, proximal (plateau); bicondylar, includes internal fixation, when performed (includes arthroscopy)
29866	Arthroscopy, knee, surgical; osteochondral autograft(s) (e.g., mosaicplasty) (includes harvesting of autograft[s])
29867	Arthroscopy, knee, surgical; osteochondral allograft

	(e.g., mosaicplasty)
29868	Arthroscopy, knee, surgical; meniscal transplantation (includes arthrotomy for meniscal insertion), medial or lateral
29870	Arthroscopy, knee, diagnostic, with or without synovial biopsy (separate procedure)
29871	Arthroscopy, knee, surgical; for infection, lavage and drainage
29873	Arthroscopy, knee, surgical; with lateral release
29874	Arthroscopy, knee, surgical; for removal of loose body or foreign body (e.g., osteochondritis dissecans fragmentation, chondral fragmentation)
29875	Arthroscopy, knee, surgical; synovectomy, limited (e.g., plica or shelf resection) (separate procedure)
29876	Arthroscopy, knee, surgical; synovectomy, major, 2 or more compartments (e.g., medial or lateral)
29877	Arthroscopy, knee, surgical; debridement/shaving of articular cartilage (chondroplasty)
29879	Arthroscopy, knee, surgical; abrasion arthroplasty (includes chondroplasty where necessary) or multiple drilling or microfracture
29880	Arthroscopy, knee, surgical; with meniscectomy (medial AND lateral, including any meniscal shaving) including debridement/shaving of articular cartilage (chondroplasty), same or separate compartment(s), when performed
29881	Arthroscopy, knee, surgical; with meniscectomy (medial OR lateral, including any meniscal shaving) including debridement/shaving of articular cartilage (chondroplasty), same or separate compartment(s), when performed
29882	Arthroscopy, knee, surgical; with meniscus repair (medial OR lateral)

29883	Arthroscopy, knee, surgical; with meniscus repair (medial AND lateral)
29884	Arthroscopy, knee, surgical; with lysis of adhesions, with or without manipulation (separate procedure)
29885	Arthroscopy, knee, surgical; drilling for osteochondritis dissecans with bone grafting, with or without internal fixation (including debridement of base of lesion)
29886	Arthroscopy, knee, surgical; drilling for intact osteochondritis dissecans lesion
29887	Arthroscopy, knee, surgical; drilling for intact osteochondritis dissecans lesion with internal fixation
29888	Arthroscopically aided anterior cruciate ligament repair/augmentation or reconstruction
29889	Arthroscopically aided posterior cruciate ligament repair/augmentation or reconstruction
29999	Unlisted procedure, arthroscopy
J7330	Cultured chondrocytes implant
S2112	Knee arthroscopy harvest

Medical Evidence

In a 2022 Cochrane review, O'Connor et al. reviewed the literature to determine the efficacy of arthroscopic knee surgery for the treatment of symptomatic knee osteoarthritis, including for degenerative meniscal tears. While current guidelines discourage the use of the procedure for these indications, it is still performed with some regularity. The review analyzed benefits and harms (e.g., debridement, partial meniscectomy, or both) when compared to non-surgical interventions (e.g., physical therapy, exercise, intra-articular glucocorticoid injections, non-arthroscopic lavage, non-steroidal anti-inflammatory drugs, hyaluronic acid injections) or placebo surgery. Sixteen trials were included, representing randomized controlled trials (RCTs) and quasi-randomized trials. A total of 2105 patients (age range 46 to 65 years; 56% women) met the inclusion criteria. The authors affirmed that arthroscopic surgery is not recommended for patients with symptomatic degenerative knee disease as the evidence does not demonstrate clinically significant improvements in pain, function, or quality of life. In addition, arthroscopy may advance the progression of knee osteoarthritis and may result in earlier or additional surgery (e.g., knee arthroplasty, osteotomy).¹

A 2024 publication by Sonesson et al. examined the 10-year outcomes among 150 middle-aged patients with meniscal symptoms who were randomized to either nonsurgical or surgical management. At the time of follow-up, the 142 surviving patients were evaluated for radiographic presence of osteoarthritis, as well as symptoms of osteoarthritis. The authors found that patient-reported outcomes were similar between cohorts and, given the short-term benefit and lack of long-term harm of knee arthroscopy, endorsed it as a first-line treatment for meniscal pathology after a failed trial of conservative care for a minimum of three months.³⁴

The Academy of Orthopaedic Surgeons (AAOS) has published four clinical guidelines to direct the management of patients with disease and damage of the knee that may be treated with arthroscopy. *Diagnosis and Treatment of Osteochondritis Dissecans* was published in 2023.²⁴ Therein, two recommendations, classified as limited in strength, were made regarding the surgical management of osteochondritis dissecans. The authors noted that the current body of evidence is insufficient to make a recommendation for or

against surgical intervention for both adult and pediatric patients. They cite the associated benefits and risks for nonoperative and operative treatment and ultimately opine, as a workgroup, that patients should be given the option of undergoing surgery. *Management of Acute Meniscal Pathology* was adopted in 2024, although the workgroup notes that current evidence is insufficient to establish strong recommendations.³³ Importantly, a moderate-strength recommendation is made regarding the preservation of meniscal tissue during surgical intervention in order to reduce the future risk of osteoarthritis. The authors opine that patients with a displaced acute meniscal tear or a symptomatic acute meniscal tear would benefit from early surgical intervention. Elsewhere in the guideline, the authors recommend surgical treatment of acute meniscal tears within six months of injury after failed conservative treatment, although this guidance was limited in strength and based on low-quality evidence. The workgroup advocates for future research in this area to allow for improved recommendations to be made in the future. *Management of Anterior Cruciate Ligament Injuries* was published in 2022.¹⁹ The AAOS recommends operative treatment of ACL injuries in select patients. Early reconstruction for patients with acute isolated ACL tears is recommended due to the increased risk profile (potential chondral injury, meniscal injury) beyond 3 months. In general, ACL reconstruction was endorsed over ACL repair due to the higher risk of revision surgery after ACL repair. ACL reconstruction was characterized as the standard of care for primary ACL injury. Both recommendations were categorized as strong and were based on evidence that was rated as high-quality.¹⁹ *Management of Osteoarthritis of the Knee (Non-Arthroplasty)* was adopted in 2021. In this guideline, the AAOS does not recommend arthroscopy with lavage and/or debridement for primary knee osteoarthritis. Among patients with meniscal tears and concomitant mild to moderate osteoarthritis, arthroscopic partial meniscectomy may be necessary when conservative treatment (e.g., physical therapy) has been unsuccessful. Both recommendations were made with moderate strength.²⁵

The AAOS has also issued position statements pertaining to knee arthroscopy. Information statement 1047, published in 2016, acknowledges the increased patient safety risks conferred by tobacco use – including increased ventilatory support, myocardial infarction, cardiac arrest, cerebrovascular accident, sepsis, and death.³⁵ The AAOS states that patients who are active smokers may reduce these risks through cessation of smoking prior to

surgery; they also note the special role orthopaedic surgeons play in counseling patients on the benefits of reduced or eliminated tobacco use before surgery. Importantly, unconfirmed cessation is not endorsed as a hard stop to surgery; rather, the surgeon's unique role as an advocate for preoperative smoking cessation is emphasized. Statements 1040 and 1184 discuss the impact of obesity on musculoskeletal conditions.^{36,37} Patients with morbid obesity (BMI of 40 or above) are encouraged to participate in a weight loss program, obtain weight reduction resources through their physician, rectify nutritional deficiencies, and consider a delay in surgical treatment if it would facilitate participation in weight loss interventions that may improve surgical outcomes. Statement 1040 notes that individuals with obesity face an increased risk for sports injuries and that when such injuries are treated arthroscopically, the procedure may be more technically difficult because of the loss of superficial landmarks.³⁶ Questions remain as to whether functional results are affected by obesity. Further, the authors note the risks associated with general anesthesia for patients with obesity and emphasize the importance of adequate patient positioning and padding to avoid pressure ulcers, nerve palsies, and compartment syndromes, which are more common among obese patients. In general, obesity is associated with a greater risk of premature complications and mortality during the perioperative period. The AAOS endorses compassionate, risk-informed patient counseling for obese patients who are considering surgery. Careful screening and appropriate referral to nutrition or endocrine care is also endorsed. Statement 1184 reinforces the risks associated with obesity in the setting of orthopaedic care and similarly encourages adequate patient counseling prior to surgery.³⁷

Social determinants of health remain an important area of ongoing orthopaedic surgery research, with recent literature raising questions regarding the healthcare disparities that may be potentiated by care limitations based on obesity and smoking status/nicotine dependence.³⁸⁻⁴⁴ Other ongoing research interrogates the impacts that biological sex, race, and socioeconomic status have on knee arthroscopy utilization and outcomes.⁴⁵⁻⁵³

References

1. O'Connor D, Johnston RV, Brignardello-Petersen R, et al. Arthroscopic surgery for degenerative knee disease (osteoarthritis including degenerative meniscal tears). *Cochrane Database Syst Rev*. 2022 Mar 3;3(3):CD014328. doi: 10.1002/14651858.CD014328. PMID: 35238404
2. Howell M, Liao Q, Gee CW. Surgical management of osteochondral defects of the knee: An educational review. *Curr Rev Musculoskelet Med*. 2021 Feb;14(1):60–66. doi: 10.1007/s12178-020-09685-1. PMID: 33587261; PMCID: PMC7930143
3. Mall NA, Harris JD, Cole BJ. Clinical evaluation and preoperative planning of articular cartilage lesions of the knee. *J Am Acad Orthop Surg*. 2015 Oct;23(10):633–40. doi: 10.5435/JAAOS-D-14-00241. PMID: 26377673
4. Harris JD, Siston RA, Pan X, et al. Autologous chondrocyte implantation: A systematic review. *J Bone Joint Surg Am*. 2010 Sep 15;92(12):2220–33. doi: 10.2106/JBJS.J.00049. PMID: 20844166; PMCID: PMC7373451
5. Lim HC, Bae JH, Song SH, et al. Current treatments of isolated articular cartilage lesions of the knee achieve similar outcomes. *Clin Orthop Relat Res*. 2012 Aug;470(8):2261–7. doi: 10.1007/s11999-012-2304-9. PMID: 22422593; PMCID: PMC3392395
6. Ulstein S, Årøen A, Røtterud JH, et al. Microfracture technique versus osteochondral autologous transplantation mosaicplasty in patients with articular chondral lesions of the knee: A prospective randomized trial with long-term follow-up. *Knee Surg Sports Traumatol Arthrosc*. 2014 Jun;22(6):1207–15. doi: 10.1007/s00167-014-2843-6. PMID: 24441734; PMCID: PMC4028546
7. Thompson R, Novikov D, Cizmic Z, et al. Arthrofibrosis after total knee arthroplasty: Pathophysiology, diagnosis, and management. *Orthop Clin North Am*. 2019 Jul;50(3):269–279. doi: 10.1016/j.ocl.2019.02.005. PMID: 31084828

8. Phillips BB, Mihalko MJ. Arthroscopy of the lower extremity. In: Azar FM, Beaty JH, editors. *Campbell's Operative Orthopaedics*. 14th ed. Philadelphia, PA: Elsevier; 2021:2576–2662.e8
9. Mascioli AA, Park AL. Infectious arthritis. In: Azar FM, Beaty JH, editors. *Campbell's Operative Orthopaedics*. 14th ed. Philadelphia, PA: Elsevier; 2021:842–867.e2
10. Donders CM, Spaans AJ, Bessems JHJM, et al. Arthrocentesis, arthroscopy or arthrotomy for septic knee arthritis in children: A systematic review. *J Child Orthop*. 2021 Feb 1;15(1):48–54. doi: 10.1302/1863-2548.15.200129. PMID: 33643458; PMCID: PMC7907762
11. Dabov GD. Osteomyelitis. In: Azar FM, Beaty JH, editors. *Campbell's Operative Orthopaedics*. 14th ed. Philadelphia, PA: Elsevier; 2021:817–841.e6
12. Miller RH III, Azar FM. Knee injuries. In: Azar FM, Beaty JH, editors. *Campbell's Operative Orthopaedics*. 14th ed. Philadelphia, PA: Elsevier; 2021:2198–2373.e18
13. Healey JH, Bernthal NM, van de Sande M. Management of tenosynovial giant cell tumor: A neoplastic and inflammatory disease. *J Am Acad Orthop Surg Glob Res Rev*. 2020 Nov;4(11):e20.00028. doi: 10.5435/JAAOSGlobal-D-20-00028. PMID: 33156160; PMCID: PMC7643913
14. Kelly MP, Bush-Joseph CA. Arthroscopic Synovectomy of the Knee. In: DeLee, Drez, Miller, editors. *Orthopaedic Sports Medicine*. 5th ed. Philadelphia, PA: Elsevier; 2020:1127–1131.e1
15. van Vulpen LFD, Thomas S, Keny SA, et al. Synovitis and synovectomy in haemophilia. *Haemophilia*. 2021 Feb;27 Suppl 3(Suppl 3):96–102. doi: 10.1111/hae.14025. PMID: 32490595; PMCID: PMC7984224.
16. Totlis T, Fermín TM, Kalifis G, et al. Arthroscopic debridement for focal articular cartilage lesions of the knee: A systematic review. *Surgeon*. 2021 Dec 1;19(6):356–64
17. Falah M, Nierenberg G, Soudry M, Hayden M, Volpin G. Treatment of articular cartilage lesions of the knee. *Int Ortho*. 2010 Jun;34:621–30
18. Diermeier TA, Rothrauff BB, Engebretsen L, et al. Treatment after ACL injury: Panther Symposium ACL Treatment Consensus Group. *Br J Sports*

Med. 2021 Jan;55(1):14–22. doi: 10.1136/bjsports-2020-102200. PMID: 32661128

19. American Academy of Orthopaedic Surgeons (AAOS). Management of anterior cruciate ligament injuries: Evidence-based clinical practice guideline. Published August 22, 2022. Accessed January 21, 2025. <https://www.aaos.org/globalassets/quality-and-practice-resources/anterior-cruciate-ligament-injuries/aclcp.pdf>
20. LaBella CR, Hennrikus W, Hewett TE, et al. Anterior cruciate ligament injuries: Diagnosis, treatment, and prevention. *Peds.* 2014 May;133(5):e1437–50. doi: 10.1542/peds.2014-0623. PMID: 24777218
21. Weiss S, Krause M, Frosch KH. Posterolateral corner of the knee: A systematic literature review of current concepts of arthroscopic reconstruction. *Arch Orthop Trauma Surg.* 2020 Dec;140(12):2003–2012. doi: 10.1007/s00402-020-03607-z. PMID: 32955608; PMCID: PMC7674327
22. Song JG, Nha KW, Lee SW. Open posterior approach versus arthroscopic suture fixation for displaced posterior cruciate ligament avulsion fractures: Systematic review. *Knee Surg & Rel Res.* 2018 Dec;30(4):275
23. Chau MM, Klimstra MA, Wise KL, et al. Osteochondritis dissecans: Current understanding of epidemiology, etiology, management, and outcomes. *J Bone Joint Surg Am.* 2021 Jun 16;103(12):1132–1151. doi: 10.2106/JBJS.20.01399. PMID: 34109940; PMCID: PMC8272630
24. American Academy of Orthopaedic Surgeons (AAOS) Diagnosis and Treatment of Osteochondritis Dissecans Evidence-Based Clinical Practice Guideline. AAOS.org/ocdcp.org Published December 1, 2023. Accessed January 21, 2025
25. Brophy RH, Fillingham YA. AAOS clinical practice guideline summary: Management of osteoarthritis of the knee (non-arthroplasty), third edition. *J Am Acad Orthop Surg.* 2022 May 1;30(9):e721–e729. doi: 10.5435/JAAOS-D-21-01233. PMID: 35383651
26. Katz JN, Arant KR, Loeser RF. Diagnosis and treatment of hip and knee osteoarthritis: A review. *JAMA.* 2021 Feb 9;325(6):568–578. doi: 10.1001/jama.2020.22171. PMID: 33560326; PMCID: PMC8225295

27. Abram SGF, Beard DJ, Price AJ, et al. Arthroscopic meniscal surgery: A national society treatment guideline and consensus statement. *Bone Joint J.* 2019 Jun;101-B(6):652–659. doi: 10.1302/0301-620X.101B6.BJJ-2019-0126.R1. PMID: 31154847
28. Beaufils P, Becker R, Kopf S, et al. Surgical management of degenerative meniscus lesions: The 2016 ESSKA meniscus consensus. *Knee Surg Sports Traumatol Arthrosc.* 2017 Feb;25(2):335–346. PMID: 28210788
29. Kopf S, Beaufils P, Hirschmann MT, et al. Management of traumatic meniscus tears: The 2019 ESSKA meniscus consensus. *Knee Surg Sports Traumatol Arthrosc.* 2020 Apr;28(4):1177–1194. doi: 10.1007/s00167-020-05847-3. PMID: 32052121; PMCID: PMC7148286
30. Chahla J, Hinckel BB, Yanke AB, et al. An expert consensus statement on the management of large chondral and osteochondral defects in the patellofemoral joint. *Orthop J Sports Med.* 2020 Mar 26;8(3):2325967120907343. doi: 10.1177/2325967120907343. PMID: 32258181; PMCID: PMC7099674
31. Khan M, Evaniew N, Bedi A, et al. Arthroscopic surgery for degenerative tears of the meniscus: A systematic review and meta-analysis. *CMAJ.* 2014 Oct 7;186(14):1057–64. doi: 10.1503/cmaj.140433. PMID: 25157057; PMCID: PMC4188648
32. Brignardello-Petersen R, Guyatt GH, Buchbinder R, et al. Knee arthroscopy versus conservative management in patients with degenerative knee disease: A systematic review. *BMJ Open.* 2017 May 11;7(5):e016114. doi: 10.1136/bmjopen-2017-016114. PMID: 28495819
33. American Academy of Orthopaedic Surgeons (AAOS). Management of Acute Meniscal Pathology Evidence-Based Clinical Practice Guideline. [aaos.org/ampcpg](https://www.aaos.org/ampcpg) Published June 10, 2024. Accessed January 21, 2025
34. Sonesson S, Springer I, Yakob J, Hedevis H, Gauffin H, Kvist J. Knee arthroscopic surgery in middle-aged patients with meniscal symptoms: a 10-year follow-up of a prospective, randomized controlled trial. *Amer J Sports Med.* 2024 Jul;52(9):2250–9
35. American Academy of Orthopaedic Surgeons (AAOS). Information Statement 1047: Tobacco Use and Orthopaedic Surgery. Published February 2016. Accessed January 3, 2025. <https://www.aaos.org/>

globalassets/about/bylaws-library/information-statements/1047-tobacco-use-and-orthopaedic-surgery-3.pdf

36. American Academy of Orthopaedic Surgeons (AAOS). Information Statement 1040: Obesity and Musculoskeletal Care. Published June 2022. Accessed January 3, 2025. <https://www.aaos.org/globalassets/about/bylaws-library/information-statements/1040-obesity-and-musculoskeletal-care.pdf>
37. American Academy of Orthopaedic Surgeons (AAOS). Information Statement 1184: The Impact of Obesity on Bone and Joint Health. Published March 2015. Accessed January 3, 2025. <https://www.aaos.org/contentassets/1cd7f41417ec4dd4b5c4c48532183b96/1184-the-impact-of-obesity-on-bone-and-joint-health1.pdf>
38. Sing DC, Luan TF, Feeley BT, Zhang AL. Is obesity a risk factor for adverse events after knee arthroscopy?. *Arthrosc*. 2016 Jul 1;32(7):1346–53
39. Zhang T, Jauregui JJ, Foster M, et al. Outcomes of partial meniscectomy in obese patients: A systematic review and meta-analysis. *Cartilage*. 2021 Dec;13(1_suppl):216S–27S
40. Ford BT, Kong R, Wellington IJ, et al. Impact of obesity, smoking, and age on 30-day postoperative outcomes of patients undergoing arthroscopic meniscus surgery. *Ortho*. 2024 Nov 1;47(6):332–6
41. Bi AS, Mojica ES, Markus DH, et al. Risk of postoperative stiffness following multiligamentous knee injury surgery is not affected by obesity: A multicenter study. *Arthrosc*. 2022 Dec 1;38(12):3175–81
42. Holle AM, Tummala SV, Pejic J, et al. Association of smokeless tobacco use with perioperative complications and revision surgery after anterior cruciate ligament reconstruction. *Amer J Sports Med*. 2025 Jan 4:03635465241303487.
43. Abdel Khalik H, Shanmugaraj A, Ekhtiari S, et al. Self-defined former smokers consume the highest opioid quantities following knee and shoulder arthroscopy. *Knee Surg, Sports Traumatol, Arthrosc*. 2024 Aug 6;33(3):856.

44. Everhart JS, Vajapey S, Kirven JC, et al. Symptom chronicity and tobacco use: Differences in athletic and nonathletic candidates for cartilage surgery. *Cartilage*. 2021 Oct;12(4):448–55.
45. Mok AC, Fancher AJ, Vopat ML, et al. Sex-specific outcomes after anterior cruciate ligament reconstruction: A systematic review and meta-analysis. *Ortho J Sports Med*. 2022 Feb 23;10(2):23259671221076883
46. Rohde MS, Shea KG, Dawson T, et al. Age, sex, and BMI differences related to repairable meniscal tears in pediatric and adolescent patients. *Amer J Sports Med*. 2023 Feb;51(2):389–97
47. Amen TB, Chatterjee A, Rudisill SS, et al. National patterns in utilization of knee and hip arthroscopy: An analysis of racial, ethnic, and geographic disparities in the United States. *Ortho J Sports Med*. 2023 Aug 16;11(8):23259671231187447
48. Bloise C, Fong B, Jeffers K, et al. Predictors of disparities in patient-reported outcomes before and after arthroscopic meniscectomy. *TJ Knee Surg*. 2023 Jun;36(07):792–800
49. Wentt CL, Farrow LD, Everhart JS, Spindler KP, Jones MH. Are there racial disparities in knee symptoms and articular cartilage damage in patients presenting for arthroscopic partial meniscectomy?. *JBJS Open Acc*. 2022 Jul 1;7(3):e21
50. Fang CJ, Miller JA, Yergensen CJ, et al. Racial/ethnic disparities exist among patients who undergo anterior cruciate ligament reconstruction in socioeconomic status, perception of health status and literacy. *Arthrosc, Sports Med, and Rehab*. 2024 Sep 11:101000
51. Prinz NW, Thompson XD, Leicht AS, Kuenze C, Hart JM. Associations between race and socioeconomic status, lower extremity strength, and patient-reported outcomes following anterior cruciate ligament reconstruction. *J Athl Training*. 2024 Dec 1;59(12):1171–7
52. Testa EJ, Modest JM, Brodeur P, et al. Do patient demographic and socioeconomic factors influence surgical treatment rates after ACL injury?. *J Racial Ethn Health Disp*. 2023 Feb;10(1):319–24

53. England P, Patel NM. Disparities in evaluation, treatment, and outcomes of pediatric knee injuries. *Sports Med and Arthrosc Rev*. 2024 Dec 1;32(4):182–8
54. Ro KH, Kim JH, Heo JW, Lee DH. Clinical and radiological outcomes of meniscal repair versus partial meniscectomy for medial meniscus root tears: a systematic review and meta-analysis. *Ortho J Sports Med*. 2020 Nov 10;8(11):2325967120962078
55. Saltzman BM, Meyer MA, Leroux TS, et al. The influence of full-thickness chondral defects on outcomes following meniscal allograft transplantation: A comparative study. *Arthrosc*. 2018 Feb 1;34(2):519–29
56. Kempshall PJ, Parkinson B, Thomas M, et al. Outcome of meniscal allograft transplantation related to articular cartilage status: Advanced chondral damage should not be a contraindication. *Knee Surg, Sports Traumatol, Arthrosc*. 2015 Jan;23(1):280–9
57. Frank R, Gilat R, Haunschild ED, et al. Do outcomes of meniscal allograft transplantation differ based on age and sex? A comparative group analysis. *Arthrosc*. 2022 Feb 1;38(2):452–65

Policy Revision History/Information

Original Date: March 20, 2023		
Review History		
Version 2	10/13/2023	Policy criteria reviewed and updated per medical literature
Version 3	12/1/2023	Policy criteria reviewed and updated per medical literature
Version 4	05/30/2024	Annual policy review
Version 5	09/20/2024	Updated language regarding conservative treatment.
Version 6	04/17/2025	<p>Annual policy review & restructure:</p> <p>Removed requirement for “young, active patient” for medial or lateral meniscal tears</p> <p>Permitted osteochondritis dissecans to be visualized on plain radiograph per professional society guideline</p> <p>Removed qualifier for “due to patient's occupation” on ACL reconstruction per professional society guidelines</p> <p>Removed requirement for conservative management of greater than 3 months for ACL tear per professional society guidelines</p> <p>Modified indications for meniscus allograft transplantation for clarity and to better reflect the referenced source</p> <p>Removed redundant synovectomy indication for Lyme arthritis, which was already captured in prior indication</p>

		<p>Removed redundant synovectomy indication for tenosynovial giant cell tumor, which was already captured in prior indication</p> <p>Added indications for synovectomy: synovial hemangioma, degenerative synovitis, plica syndrome</p> <p>Added an “out” for conservative management requirement for plica syndrome and PVNS</p> <p>Adjusted chondral lesion size for optimal ACI treatment to reflect appropriate arthroscopic management versus open arthrotomy</p> <p>Added minimum age of 12 for optimal ACI treatment</p> <p>Removed requirement for grade III or grade IV osteochondral defect for optimal arthroscopic ACI treatment versus open arthrotomy</p> <p>Moved debridement of articular cartilage lesion indication beneath existing indications for debridement</p> <p>Removed hemangioma indication under diagnostic arthroscopy as this indication was better covered elsewhere</p> <p>Added indication for arthroscopic management of certain fracture types</p> <p>Removed BMI requirement for meniscus allograft as it is not an absolute contraindication</p> <p>Consolidated all indications which require three months of conservative care for policy’s ease of use.</p>
--	--	--

		<p>Updated conservative care to current internal standard language.</p> <p>Conservative care language modified to reflect non-opioid pain control.</p> <p>Modified steroid injection language for clarity.</p> <p>Added references.</p> <p>Updated medical evidence section.</p>
Version 7	5/22/2025	<p>Added CPT codes: 27405, 27407, 27409, 27427</p> <p>Slight modification to ligamentous injury indication to allow approval for lateral collateral ligament injury - changed "cruciate ligament injury" to "ligament injury" and added bullet points: presence of LCL tear, presence of PLC injury</p>
Version 8	9/11/2025	<p>Added "unstable lesion/unstable defect on imaging" as indication under osteochondritis dissecans to allow approval for patients who may not have mechanical symptoms</p> <p>Added requirement that underlying malalignment be addressed with osteotomy in the same procedure as meniscal transplant or autologous chondrocyte implantation per current medical guidelines</p> <p>Added definition of advanced osteoarthritis to "definition" section</p>